

Chapter 7

REFERENCES

1. Lagasse RS. Anesthesia safety: model or myth? A review of the published literature and analysis of current original data. *Anesthesiology*. 2002;97(6):1609-1617.
2. American Society of Anesthesiologists. *Practice Advisory for Preoperative Evaluation*. <http://www.asahq.org/publications-AndServices/preeval.pdf>.
3. American Society of Anesthesiologists. New classification of physical status. *Anesthesiology*. 1963;24:111.
4. Menke H, Klein A, John KD, Junginger T. Predictive value of ASA classification for the assessment of the perioperative risk. *Int Surg*. 1993;78(3):266-270.
5. Gilbert K, Larocque BJ, Patrick LT. Prospective evaluation of cardiac risk indices for patients undergoing noncardiac surgery. *Ann Intern Med*. 2000;133(5):356-359.
6. Fleisher LA, Beckman JA, Brown KA, et al. ACC/AHA 2007 guidelines on perioperative cardiovascular evaluation and care for noncardiac surgery: Executive summary: A report of the American College of Cardiology/American Heart Association Task Force on practice guidelines (writing committee to revise the 2002 guidelines on perioperative cardiovascular evaluation for noncardiac surgery). *Anesth Analg*. March 2008;106:685-712.
7. Apfel CC, Korttila K, Abdalla M, et al. A factorial trial of six interventions for the prevention of postoperative nausea and vomiting. *N Engl J Med*. 2004;350(24):2441-2451.
8. Golembiewski J, Chernin E, Chopra T. Prevention and treatment of postoperative nausea and vomiting. *Am J Health Syst Pharm*. 2005;62(12):1247-1260; quiz 1261-1242.
9. Apfel CC, Stoocklein K, Lipfert P. PONV: a problem of inhalational anaesthesia? *Best Pract Res Clin Anaesthesiol*. 2005;19(3):485-500.
10. Gan TJ. "Black box" warning on droperidol: report of the FDA convened expert panel. *Anesth Analg*. 2004;98(6):1809.
11. Dershwitz M. Droperidol: should the black box be light gray? *J Clin Anesth*. 2002;14(8):598-603.
12. Roth S, Thisted RA, Erickson JP, Black S, Schreider BD. Eye injuries after nonocular surgery. A study of 60,965 anesthetics from 1988 to 1992. *Anesthesiology*. 1996;85(5):1020-1027.
13. Roth S, Gillesberg I. *Injuries to the Visual System and Other Sense Organs*. St. Louis, MO: Mosby; 1999.
14. Rosenstock C, Gillesberg I, Gatke MR, Levin D, Kristensen MS, Rasmussen LS. Inter-observer agreement of tests used for prediction of difficult laryngoscopy/tracheal intubation. *Acta Anaesthesiol Scand*. 2005;49(8):1057-1062.
15. Mashour GA, Wang LY, Turner CR, Vandervest JC, Shanks A, Tremper KK. A Retrospective Study of Intraoperative Awareness with Methodological Implications. *Anesth Analg*. 2009;108(2):521-526.
16. Sebel PS, Boedle TA, Ghoneim MM, Rampil IJ, Padilla RE, Gan TJ, Domino KB. The incidence of awareness during anesthesia: A multicenter United States study. *Anesth Analg*. 2004;99:833-9.
17. Ording H. Incidence of malignant hyperthermia in Denmark. *Anesth Analg*. 1985;64(7):700-704.
18. Bachand M, Vachon N, Boisvert M, Mayer FM, Chartrand D. Clinical reassessment of malignant hyperthermia in Abitibi-Temiscamingue. *Can J Anaesth*. 1997;44(7):696-701.
19. Ali SZ, Taguchi A, Rosenberg H. Malignant hyperthermia. *Best Pract Res Clin Anaesthesiol*. 2003;17(4):519-533.
20. Bandom BW, Muldoon SM. The practicality and need for genetic testing for malignant hyperthermia. *Anesthesiology*. 2005;103(5):1100.
21. Bryson GL, Chung F, Cox RG, et al. Patient selection in ambulatory anesthesia—an evidence-based review: part II. *Can J Anaesth*. 2004;51(8):782-794.
22. Malignant Hyperthermia Association of the United States. Medical FAQs. <http://www.mhaus.org/index.cfm/fuseaction/Content.Display/PagePK/MedicalFAQs.cfm>.
23. Yen T, Nightingale BN, Burns JC, Sullivan DR, Stewart PM. Butyrylcholinesterase (BCHE) genotyping for post-succinylcholine apnea in an Australian population. *Clin Chem*. 2003;49(8):1297-1308.
24. Nilsson E. Case of prolonged apnea after succinylcholine administration [article in undetermined language]. *Sven Lakartidn*. 1953;50(42):2167-2170.
25. Cerf C, Mesguish M, Gabriel I, Amselem S, Duvaldestin P. Screening patients with prolonged neuromuscular blockade after succinylcholine and mivacurium. *Anesth Analg*. 2002;94(2):461-466; table of contents.
26. Osterman JE, Hopper J, Heran WJ, Keane TM, van der Kolk BA. Awareness under anesthesia and the development of posttraumatic stress disorder. *Gen Hosp Psychiatry*. 2001;23(4):198-204.
27. Domino KB, Posner KL, Caplan RA, Cheney FW. Awareness during anesthesia: a closed claims analysis. *Anesthesiology*. 1999;90(4):1053-1061.
28. Leslie K, Myles PS, Forbes A, Chan MT, Short TG, Swallow SK. Recovery from bispectral index-guided anaesthesia in a large randomized controlled trial of patients at high risk of awareness. *Anaesth Intensive Care*. 2005;33(4):443-451.

29. Mychaskiw G 2nd, Horowitz M, Sachdev V, Heath BJ. Explicit intraoperative recall at a Bispectral Index of 47. *Anesth Analg*. 2001;92(4):808-809.
30. Rampersad SE, Mulroy MF. A case of awareness despite an "adequate depth of anesthesia" as indicated by a Bispectral Index monitor. *Anesth Analg*. 2005;100(5):1363-1364; table of contents.
31. Kazanjian P. Anesthesia Complications. In Mulholland W, Doherty G, eds. *Complications in Surgery*, Philadelphia, PA: Lippincott Williams & Wilkins; 2006.
32. American Society of Anesthesiologists Practice advisory for intraoperative awareness and brain function monitoring. *Anesthesiology*. 2006;104(4):847-864.
33. Sebel PS, Bowdle TA, Ghoneim MM, et al. The incidence of awareness during anesthesia: a multicenter United States study. *Anesth Analg*. 2004;99(3):833-839; table of contents.
34. Mertes PM, Laxenaire MC. Allergy and anaphylaxis in anaesthesia. *Minerva Anesthesiol*. 2004;70(5):285-291.
35. Yilmaz E, Ozcan S, Basar M, Basar H, Batislam E, Ferhat M. Music decreases anxiety and provides sedation in extracorporeal shock wave lithotripsy. *Urology*. 2003;61(2):282-286.
36. Halaszynski TM, Juda R, Silverman DG. Optimizing postoperative outcomes with efficient preoperative assessment and management. *Crit Care Med*. 2004;32(4 Suppl):S76-S86.
37. American Society of Anesthesiologists. *What You Should Know About Herbal and Dietary Supplement Use*. <http://www.asahq.org/patientEducation/herbPhysician.pdf>, 2003.
38. Hyatt B, Bensky KP. Illicit drugs and anesthesia. *CRNA*. 1999;10(1):15-23.
39. Angeli F, Verdecchia P, Gattobigio R, Sardone M, Reboldi G. White-coat hypertension in adults. *Blood Press Monit*. 2005;10(6):301-305.
40. Caplan RA, Posner KL, Ward RJ, Cheney FW. Adverse respiratory events in anesthesia: a closed claims analysis. *Anesthesiology*. 1990;72(5):828-833.
41. Domino KB, Posner KL, Caplan RA, Cheney FW. Airway injury during anesthesia: a closed claims analysis. *Anesthesiology*. 1999;91(6):1703-1711.
42. Langeron O, Masso E, Huraux C, et al. Prediction of difficult mask ventilation. *Anesthesiology*. 2000;92(5):1229-1236.
43. Khan ZH, Kashfi A, Ebrahimkhani E. A comparison of the upper lip bite test (a simple new technique) with modified Mallampati classification in predicting difficulty in endotracheal intubation: a prospective blinded study. *Anesth Analg*. 2003;96(2):595-599; table of contents.
44. Samssoon GL, Young JR. Difficult tracheal intubation: a retrospective study. *Anaesthesia*. 1987;42(5):487-490.
45. Cattano D, Panicucci E, Paolicchi A, Forfori F, Giunta F, Hagberg C. Risk factors assessment of the difficult airway: an Italian survey of 1956 patients. *Anesth Analg*. 2004;99(6):1774-1779; table of contents.
46. Farmery AD. Sternomental distance as predictor of difficult laryngoscopy. *Br J Anaesth*. 1997;78(5):626-627.
47. Juvin P, Lavaut E, Dupont H, et al. Difficult tracheal intubation is more common in obese than in lean patients. *Anesth Analg*. 2003;97(2):595-600; table of contents.
48. Arné J, Descoins P, Fuscuardi J, et al. Preoperative assessment for difficult intubation in general and ENT surgery: predictive value of a clinical multivariate risk index. *Br J Anaesth*. 1998;80(2):140-146.
49. Pearce A. Evaluation of the airway and preparation for difficulty. *Best Pract Res Clin Anaesthesiol*. 2005;19(4):559-579.
50. Han R, Tremper KK, Kheterpal S, O'Reilly M. Grading scale for mask ventilation. *Anesthesiology*. 2004;101(1):267.
51. Kheterpal S HR, Shanks A, O'Reilly M, Tremper KK. Incidence and predictors of difficult and impossible mask ventilation. *Anesthesiology*. 2005;103:A1415.
52. American Society of Anesthesiologists. *Practice Advisory for Preoperative Evaluation*. <http://www.asahq.org/publicationsAndServices/preeval.pdf>.
53. Tremper KK, Benedict P. Paper "preoperative computer." *Anesthesiology*. 2000;92(4):1212-1213.
54. Ramachandran SK, Kheterpal S, Consens F, Shanks A, Doherty T, Morris M, Tremper KK. Derivation and Validation of a Simple Perioperative Sleep Apnea Prediction Score. *Anesth Analg*. 2010;110:1007-15.
55. Bray GA. Risks of obesity. *Endocrinol Metab Clin North Am*. 2003;32(4):787-804, viii.
56. Adams JP, Murphy PG. Obesity in anaesthesia and intensive care. *Br J Anaesth*. 2000;85(1):91-108.
57. Coe AJ, Saleh T, Samuel T, Edwards R. The management of patients with morbid obesity in the anaesthetic assessment clinic. *Anaesthesia*. 2004;59(6):570-573.
58. Alchanatis M, Tourkohoriti G, Kakouros S, Kosmas E, Podaras S, Jordanoglou JB. Daytime pulmonary hypertension in patients with obstructive sleep apnea: the effect of continuous positive airway pressure on pulmonary hemodynamics. *Respiration*. 2001;68(6):566-572.
59. American Society of Anesthesiologists. *Practice Guidelines for the Perioperative Management of Patients With Sleep Apnea*. <http://www.asahq.org/publicationsandservices/sleepapnea103105.pdf>.
60. Tiret L, Desmonts JM, Hatton F, Vourc'h G. Complications associated with anaesthesia—a prospective survey in France. *Can Anaesth Soc J*. 1986;33(3 Pt 1):336-344.
61. Warner MA, Saletel RA, Schroeder DR, Warner DO, Offord KP, Gray DT. Outcomes of anesthesia and surgery in people 100 years of age and older. *J Am Geriatr Soc*. 1998;46(8):988-993.
62. McLean AJ, Le Couteur DG. Aging biology and geriatric clinical pharmacology. *Pharmacol Rev*. 2004;56(2):163-184.
63. Kim JP, Kim SJ, Lee JH, Kim SW, Choi MG, Yu HJ. Surgery in the aged in Korea. *Arch Surg*. 1998;133(1):18-23.
64. Brooks-Brunn JA. Predictors of postoperative pulmonary complications following abdominal surgery. *Chest*. 1997;111(3):564-571.
65. Myles PS, Iacono GA, Hunt JO, et al. Risk of respiratory complications and wound infection in patients undergoing ambulatory surgery: smokers versus nonsmokers. *Anesthesiology*. 2002;97(4):842-847.
66. A clinical practice guideline for treating tobacco use and dependence: A US Public Health Service Report. *JAMA*. 2000;283:3244-3254.
67. Warner DO. Helping surgical patients quit smoking: why, when, and how. *Anesth Analg*. 2005;101(2):481-487; table of contents.
68. Warner DO, Patten CA, Ames SC, Offord K, Schroeder D. Smoking behavior and perceived stress in cigarette smokers undergoing elective surgery. *Anesthesiology*. 2004;100(5):1125-1137.
69. Schein OD, Katz J, Bass EB, et al. The value of routine preoperative medical testing before cataract surgery. Study of medical testing for cataract surgery. *N Engl J Med*. 2000;342(3):168-175.
70. Turnbull JM, Buck C. The value of preoperative screening investigations in otherwise healthy individuals. *Arch Intern Med*. 1987;147(6):1101-1105.

71. Olson RP, Stone A, Lubarsky D. The prevalence and significance of low preoperative hemoglobin in ASA I or 2 outpatient surgery candidates. *Anesth Analg*. 2005;101(5):1337-1340.
72. Correll DJ, Hepner DL, Chang C, Tsen L, Hevelone ND, Bader AM. Preoperative electrocardiograms: patient factors predictive of abnormalities. *Anesthesiology*. 2009;110(6):1217-1222.
73. Candido KD, Sukhani R, Doty R Jr, et al. Neurologic sequelae after interscalene brachial plexus block for shoulder/upper arm surgery: the association of patient, anesthetic, and surgical factors to the incidence and clinical course. *Anesth Analg*. 2005;100(5):1489-1495, table of contents.
74. Rodgers A, Walker N, Schug S, et al. Reduction of postoperative mortality and morbidity with epidural or spinal anaesthesia: results from overview of randomised trials. *BMJ*. 2000;321(7275):1493.
75. Farag E, Dilger J, Brooks P, Tetzlaff JE. Epidural analgesia improves early rehabilitation after total knee replacement. *J Clin Anesth*. 2005;17(4):281-285.
76. Liu SS, Strodtbeck WM, Richman JM, Wu CL. A comparison of regional versus general anesthesia for ambulatory anesthesia: a meta-analysis of randomized controlled trials. *Anesth Analg*. 2005;101(6):1634-1642.
77. Hug CC Jr. MAC should stand for maximum anesthesia caution, not minimal anesthesiology care. *Anesthesiology*. 2006;104(2):221-223.
78. Bhananker SM, Posner KL, Cheney FW, Caplan RA, Lee LA, Domino KB. Injury and liability associated with monitored anesthesia care: a closed claims analysis. *Anesthesiology*. 2006;104(2):228-234.
79. Practice guidelines for preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration: application to healthy patients undergoing elective procedures: a report by the American Society of Anesthesiologist Task Force on Preoperative Fasting. *Anesthesiology*. 1999;90(3):896-905.
80. Comfere T, Sprung J, Kumar MM, et al. Angiotensin system inhibitors in a general surgical population. *Anesth Analg*. 2005;100(3):636-644; table of contents.
81. Ang-Lee MK, Moss J, Yuan CS. Herbal medicines and perioperative care. *JAMA*. 2001;286(2):208-216.
82. Leak J. Herbal medicines: What do we need to know? *ASA Newsletter* 2000;64.
83. Jenkins K, Grady D, Wong J, Correa R, Armanious S, Chung F. Post-operative recovery: day surgery patients' preferences. *Br J Anaesth*. 2001;86(2):272-274.
84. Bauer KP, Dom PM, Ramirez AM, O'Flaherty JE. Preoperative intravenous midazolam: benefits beyond anxiolysis. *J Clin Anesth*. 2004;16(3):177-183.
85. Acil M, Basgul E, Celiker V, Karagöz AH, Demir B, Aypar U. Perioperative effects of melatonin and midazolam premedication on sedation, orientation, anxiety scores and psychomotor performance. *Eur J Anaesthesiol*. 2004;21(7):553-557.
86. Ong CK, Lirk P, Seymour RA, Jenkins BJ. The efficacy of preemptive analgesia for acute postoperative pain management: a meta-analysis. *Anesth Analg*. 2005;100(3):757-773; table of contents.
87. Grass JA. Patient-controlled analgesia. *Anesth Analg*. 2005;101(5 Suppl):S44-S61.
88. Liu S. Update in use of continuous perineural catheters for postoperative analgesia. *Anesth Analg*. 2006;(Suppl). <http://www.iars.org/2006ReviewCourseLectures/pdf/text/liu2006.pdf>. Accessed August 17, 2011.
89. Arbous MS, Meursing AE, van Kleef JW, et al. Impact of anesthesia management characteristics on severe morbidity and mortality. *Anesthesiology*. 2005;102(2):257-268; quiz 491-252.
90. Lienhart A, Auroy Y, Pequignot F, et al. Preliminary results from the SFAR-iNSERM inquiry on anaesthesia-related deaths in France: mortality rates have fallen ten-fold over the past two decades [in French]. *Bull Acad Natl Med*. 2004;188(8):1429-1437; discussion 1437-1441.
91. Pedersen T. Complications and death following anaesthesia. A prospective study with special reference to the influence of patient-, anaesthesia-, and surgery-related risk factors. *Dan Med Bull*. 1994;41(3):319-331.
92. Eichhorn JH, Cooper JB, Cullen DJ, Maier WR, Philip JH, Seeman RG. Standards for patient monitoring during anesthesia at Harvard Medical School. *JAMA*. 1986;256(8):1017-1020.
93. American Society of Anesthesiologists. Standards for Basic Intraoperative Monitoring, 1986. <http://www.asahq.org/For-Healthcare-Professionals/~media/For%20Members/documents/Standards%20Guidelines%20Stmts/Basic%20Anesthetic%20Monitoring%202005.ashx>. Accessed August 17, 2011.
94. O'Toole JD, Wurtzbacher JJ, Wearner NE, Jain AC. Pulmonary-valve injury and insufficiency during pulmonary-artery catheterization. *N Engl J Med*. 1979;301(21):1167-1168.
95. Yamada M, Tsuda K, Nagai S, Tadokoro M, Ishibe Y. A case of crush syndrome resulting from continuous compression of the upper arm by automatically cycled blood pressure cuff [in Japanese]. *Masui*. 1997;46(1):119-123.
96. Swedlow DB, Irving SM. *Monitoring and patient safety* chapter in Blitt CD, ed.: *Monitoring in anesthesia and critical care medicine*, ed. 2, New York, 1990, Churchill-Livingstone.
97. Dorje P, Tremper K. Systolic pressure variation: a dynamic measure of the adequacy of intravascular volume. *Semin Anesth Periop Med Pain*. 2005;24:147-153.
98. Perel A, Pizov R, Cotev S. Systolic blood pressure variation is a sensitive indicator of hypovolemia in ventilated dogs subjected to graded hemorrhage. *Anesthesiology*. 1987;67(4):498-502.
99. Michard F. Changes in arterial pressure during mechanical ventilation. *Anesthesiology*. 2005;103(2):419-428.
100. Mahenthiran J, Bangalore S, Yao SS, Chaudhry FA. Comparison of prognostic value of stress echocardiography versus stress electrocardiography in patients with suspected coronary artery disease. *Am J Cardiol*. 2005;96(5):628-634.
101. Kline JA, Kubin AK, Patel MM, Easton EJ, Seupal RA. Alveolar dead space as a predictor of severity of pulmonary embolism. *Acad Emerg Med*. 2000;7(6):611-617.
102. March PA, Muir WW. Bispectral analysis of the electroencephalogram: a review of its development and use in anesthesia. *Vet Anaesth Analg*. 2005;32(5):241-255.
103. Peñuelas-Acuña J, Oriol-López SA, Castelazo-Arredondo JA, Hernández-Bernal CE. Usefulness of bispectral index in pharmaceutical cost reduction for anesthesia [in Spanish]. *Cir Cir*. 2003;71(4):300-303.
104. Messner M, Beese U, Romstock J, Dinkel M, Tschaikowsky K. The bispectral index declines during neuromuscular block in fully awake persons. *Anesth Analg*. 2003;97(2):488-491, table of contents.
105. Dahaba AA. Different conditions that could result in the bispectral index indicating an incorrect hypnotic state. *Anesth Analg*. 2005;101(3):765-773.
106. Fiedler MA. Maintaining intraoperative normothermia: meta-analysis of outcomes with costs. *AANA J*. 1999;67(4):307-308.