

## Chapter 30

### REFERENCES

1. Cope DK. Monitoring in the 19th century: from blood-letting to blood-flow measurements. *ASA Newsletter*. 2002;66(9):6-8.
2. Young D, Griffiths J. Clinical trials of monitoring in anaesthesia, critical care and acute ward care: a review. *Br J Anaesth*. 2006;97(1):39-45.
3. Jain M, Canham M, Upadhyay D, Corbridge T. Variability in interventions with pulmonary artery catheter data. *Intensive Care Med*. 2003;29(11):2059-2062.
4. Cohn JN. Blood pressure measurement in shock. Mechanism of inaccuracy in auscultatory and palpatory methods. *JAMA*. 1967;199:972-976.
5. Williams JS, Brown SM, Conlin PR. Videos in clinical medicine. Blood-pressure measurement. *N Engl J Med*. 2009;360(5):e6.
6. Borow KM, Newburger JW. Noninvasive estimation of central aortic pressure using the oscillometric method for analyzing systemic artery pulsatile blood flow: comparative study of indirect systolic, diastolic, and mean brachial artery pressure with simultaneous direct ascending aortic pressure measurements. *Am Heart J*. 1982;103:879-886.
7. Lakhil K, Ehrmann S, Runge I, et al. Tracking hypotension and dynamic changes in arterial blood pressure with brachial cuff measurements. *Anesth Analg*. 2009;109(2):494-501.
8. Gardner RM. Direct blood pressure measurement—dynamic response requirements. *Anesthesiology*. 1981;54(3):227-236.
9. Courtois M, Fattal PG, Kovacs SJ, et al. Anatomically and physiologically based reference level for measurement of intracardiac pressures. *Circulation*. 1995;92:1994-2000.
10. O'Rourke MF, Gallagher DE. Pulse wave analysis. *J Hypertension*. 1996;14(5):S147-157.
11. Chauhan S, Saxena N, Mehrotra S, et al. Femoral artery pressures are more reliable than radial artery pressures on initiation of cardiopulmonary bypass. *J Cardiothorac Vasc Anesth*. 2000;14(3):274-276.
12. Dorman T, Breslow MJ, Lipsett PA, et al. Radial artery pressure monitoring underestimates central arterial pressure during vasopressor therapy in critically ill surgical patients. *Crit Care Med*. 1998;26(10):1646-1649.
13. Tegtmeier K, Brady G, Lai S, et al. Videos in clinical medicine. Placement of an arterial line. *N Engl J Med*. 2006;354(15):e13.
14. Barnes RW, Foster EJ, Janssen GA, et al. Safety of brachial arterial catheters as monitors in the intensive care unit—prospective evaluation with the Doppler ultrasonic velocity detector. *Anesthesiology*. 1976;44(3):260-264.
15. Anonymous. Standards for basic anesthesia monitoring. In: *ASA Standards, Guidelines and Statements*. Park Ridge, IL: American Society of Anesthesiologists; 1993:4-5.
16. Charlson ME, MacKenzie CR, Gold JP, et al. Intraoperative blood pressure. What patterns identify patients at risk for post-operative complications? *Ann Surg*. 1990;212(5):567-580.
17. Reich DL, Bennett-Guerrero E, Bodian CA, et al. Intraoperative tachycardia and hypertension are independently associated with adverse outcome in noncardiac surgery of long duration. *Anesth Analg*. 2002;95(2):273-277.
18. Aronson S, Stafford-Smith M, Phillips-Bute B, et al. Intraoperative systolic blood pressure variability predicts 30-day mortality in aortocoronary bypass surgery patients. *Anesthesiology*. 2010;113(2):305-312.
19. Alford JW, Palumbo MA, Barnum MJ. Compartment syndrome of the arm: a complication of noninvasive blood pressure monitoring during thrombolytic therapy for myocardial infarction. *J Clin Monit Comput*. 2002;17(3-4):163-166.
20. Bhananker SM, Liau DW, Kooner PK, et al. Liability related to peripheral venous and arterial catheterization: a closed claims analysis. *Anesth Analg*. 2009;109(1):124-129.
21. Brzezinski M, Luisetti T, London MJ. Radial artery cannulation: a comprehensive review of recent anatomic and physiologic investigations. *Anesth Analg*. 2009;109(6):1763-1781.
22. Sinha S, Jayaram R, Hargreaves CG. Fatal neuroglycopenia after accidental use of a glucose 5% solution in a peripheral arterial cannula flush system. *Anaesthesia*. 2007;62(6):615-620.
23. Mark JB. Central venous pressure monitoring: clinical insights beyond the numbers. *J Cardiothorac Vasc Anesth*. 1991;5:163-173.
24. O'Grady NP, Alexander M, Dellinger EP, et al. Guidelines for the prevention of intravascular catheter-related infections. Centers for Disease Control and Prevention. *MMWR Recomm Rep*. 2002;51(RR-10):1-29.
25. Sulek CA, Gravenstein N, Blackshear RH, et al. Head rotation during internal jugular vein cannulation and the risk of carotid artery puncture. *Anesth Analg*. 1996;82(1):125-128.
26. Daily PO, Griep RB, Shumway NE. Percutaneous internal jugular vein cannulation. *Arch Surg*. 1970;101(4):534-536.
27. Ezaru CS, Mangione MP, Oravitz TM, et al. Eliminating arterial injury during central venous catheterization using manometry. *Anesth Analg*. 2009;109(1):130-134.
28. Statement on recommendations for uniform use of real-time ultrasound guidance for placement of central venous catheters. *Statements of the American College of Surgeons*. 2008; ST-60.

29. Stone MB, Nagdev A, Murphy MC, et al. Ultrasound detection of guidewire position during central venous catheterization. *Am J Emerg Med.* 2010;28(1):82-84.
30. Kumar A, Chuan A. Ultrasound guided vascular access: efficacy and safety. *Best Pract Res Clin Anaesthesiol.* 2009;23(3):299-311.
31. Hind D, Calvert N, McWilliams R, et al. Ultrasonic locating devices for central venous cannulation: meta-analysis. *BMJ.* 2003;327(7411):361.
32. The National Institute for Clinical Excellence. *The Effectiveness and Cost-Effectiveness of Ultrasound Locating Devices for Central Venous Access.* Sheffield, UK: Trent Institute for Health Services Research; 2002:1-98.
33. Karakitsos D, Labropoulos N, De Groot E, et al. Real-time ultrasound-guided catheterisation of the internal jugular vein: a prospective comparison with the landmark technique in critical care patients. *Crit Care.* 2006;10(6):R162.
34. Girard TD, Schectman JM. Ultrasound guidance during central venous catheterization: a survey of use by house staff physicians. *J Crit Care.* 2005;20(3):224-229.
35. Parsons AJ, Alfa J. Carotid dissection: a complication of internal jugular vein cannulation with the use of ultrasound. *Anesth Analg.* 2009;109(1):135-136.
36. Blaiwas M. Video analysis of accidental arterial cannulation with dynamic ultrasound guidance for central venous access. *J Ultrasound Med.* 2009;28(9):1239-1244.
37. Albrecht K, Nave H, Breitmeier D, et al. Applied anatomy of the superior vena cava—the carina as a landmark to guide central venous catheter placement. *Br J Anaesth.* 2004;92(1):75-77.
38. Pronovost PJ, Goeschel CA, Colantuoni E, et al. Sustaining reductions in catheter related bloodstream infections in Michigan intensive care units: observational study. *BMJ.* 2010;340:c309.
39. Swan HJ, Ganz W, Forrester J, et al. Catheterization of the heart in man with use of a flow-directed balloon-tipped catheter. *N Engl J Med* 1970;283(9):447-451.
40. Mimoz O, Rauss A, Reikik N, et al. Pulmonary artery catheterization in critically ill patients: a prospective analysis of outcome changes associated with catheter-prompted changes in therapy. *Crit Care Med.* 1994;22(4):573-579.
41. Iberti TJ, Fischer EP, Leibowitz AB, et al. A multicenter study of physicians' knowledge of the pulmonary artery catheter. Pulmonary Artery Catheter Study Group *JAMA.* 1990;264(22):2928-2932.
42. Squara P, Bennett D, Perret C. Pulmonary artery catheter: does the problem lie in the users? *Chest.* 2002;121(6):2009-2015.
43. Roizen M, Berger D, Gabel R, et al. Practice guidelines for pulmonary artery catheterization: an updated report by the American Society of Anesthesiologists Task Force on Pulmonary Artery Catheterization. *Anesthesiology.* 2003;99(4):988-1014.
44. Society of Critical Care Medicine. Pulmonary Artery Catheter Consensus conference: consensus statement. *Crit Care Med.* 1997;25(6):910-925.
45. Bernard GR, Sopko G, Cerra F, et al. Pulmonary artery catheterization and clinical outcomes: National Heart, Lung, and Blood Institute and Food and Drug Administration Workshop Report. Consensus Statement. *JAMA.* 2000;283(19):2568-2572.
46. De Backer D. Hemodynamic assessment: the technique or the physician at fault? *Intensive Care Med.* 2003;29(11):1865-1867.
47. Parlow JL, Milne B, Cervenka FW. Balloon flotation is more important than flow direction in determining the position of flow-directed pulmonary artery catheters. *J Cardiothorac Vasc Anesth.* 1992;6:20-23.
48. Szabo Z. A simple method to pass a pulmonary artery flotation catheter rapidly into the pulmonary artery in anaesthetized patients. *Br J Anaesth.* 2003;90(6):794-796.
49. Tempe DK, Datt V, Banerjee A, et al. Case 5—2004: Transesophageal echocardiography-guided insertion of a pulmonary artery catheter. *J Cardiothorac Vasc Anesth.* 2004;18(5):657-662.
50. Pinsky MR. Pulmonary artery occlusion pressure. *Intensive Care Med.* 2003;29(1):19-22.
51. Falicov RE, Resnekov L. Relationship of the pulmonary artery end-diastolic pressure to the left ventricular end-diastolic and mean filling pressures in patients with and without left ventricular dysfunction. *Circulation.* 1970;42:65-73.
52. O'Quin R, Marini JJ. Pulmonary artery occlusion pressure: clinical physiology, measurement, and interpretation. *Am Rev Respir Dis.* 1983;128(2):319-326.
53. West JB, Dollery CT, Naimark A. Distribution of blood flow in isolated lung: relation to vascular and alveolar pressures. *J Appl Physiol.* 1964;19:713-724.
54. Teboul JL, Pinsky MR, Mercat A, et al. Estimating cardiac filling pressure in mechanically ventilated patients with hyperinflation. *Crit Care Med.* 2000;28(11):3631-3636.
55. Marini JJ, O'Quin R, Culver BH, et al. Estimation of transmural cardiac pressures during ventilation with PEEP. *J Appl Physiol.* 1982;53(2):384-391.
56. Jardin F, Delorme G, Hardy A, et al. Reevaluation of hemodynamic consequences of positive pressure ventilation: emphasis on cyclic right ventricular afterloading by mechanical lung inflation. *Anesthesiology.* 1990;72(6):966-970.
57. Sharkey SW. Beyond the wedge: clinical physiology and the Swan-Ganz catheter. *Am J Med.* 1987;83(1):111-122.
58. Shah KB, Rao TL, Laughlin S, et al. A review of pulmonary artery catheterization in 6,245 patients. *Anesthesiology.* 1984;61(3):271-275.
59. Procaccini B, Clementi G. Pulmonary artery catheterization in 9071 cardiac surgery patients: a review of complications [in Italian]. *Ital Heart J.* 2004;5(12):891-899.
60. Harvey S, Harrison DA, Singer M, et al. Assessment of the clinical effectiveness of pulmonary artery catheters in management of patients in intensive care (PAC-Man): a randomised controlled trial. *Lancet.* 2005;366(9484):472-477.
61. Luck JC, Engel TR. Transient right bundle branch block with "Swan-Ganz" catheterization. *Am Heart J.* 1976;92(2):263-264.
62. Sprung CL, Elser B, Schein RM, et al. Risk of right bundle-branch block and complete heart block during pulmonary artery catheterization *Crit Care Med.* 1989;17(1):1-3.
63. Fibuch EE, Tuohy GF. Intracardiac knotting of a flow-directed balloon-tipped catheter. *Anesth Analg.* 1980;59(3):217-212.
64. Kranz A, Mundigler G, Bankier A, et al. Knotting of two central venous catheters: a rare complication of pulmonary artery catheterization. *Wien Klin Wochenschr.* 1996;108(13):404-406.
65. Schwartz KV, Garcia FG. Entanglement of Swan-Ganz catheter around an intracardiac structure *JAMA.* 1977;237(12):1198-1199.
66. Dumesnil JG, Proulx G. A new nonsurgical technique for untying tight knots in flow-directed balloon catheters. *Am J Cardiol.* 1984;53(2):395-396.

67. Randolph AG, Cook DJ, Gonzales CA, et al. Benefit of heparin in central venous and pulmonary artery catheters: a meta-analysis of randomized controlled trials *Chest*. 1998;113(1):165-171.
68. Sandham JD, Hull RD, Brant RF, et al. A randomized, controlled trial of the use of pulmonary-artery catheters in high-risk surgical patients. *N Engl J Med*. 2003;348(1):5-14.
69. Foote GA, Schabel SI, Hodges M. Pulmonary complications of the flow-directed balloon-tipped catheter. *N Engl J Med* 1974;290(17):927-931.
70. Boscoe MJ, de Lange S. Damage to the tricuspid valve with a Swann-Ganz catheter. *Br Med J (Clin Res Ed)*. 1981;283(6287):346-347.
71. O'Toole JD, Wurtzbacher JJ, Wearner NE, et al. Pulmonary-valve injury and insufficiency during pulmonary-artery catheterization. *N Engl J Med*. 1979;301(21):1167-1168.
72. Ehrie M, Morgan AP, Moore FD, et al. Endocarditis with the indwelling balloon-tipped pulmonary artery catheter in burn patients. *J Trauma*. 1978;18(9):664-666.
73. Hartmann G, Steib A, Ludes B, Ravanello J. Perforation of the pulmonary artery following Swan-Ganz catheterization [in French]. *Ann Fr Anesth Reanim*. 1988;7(6):486-493.
74. Kearney TJ, Shabot MM. Pulmonary artery rupture associated with the Swan-Ganz catheter *Chest*. 1995;108(5):1349-1352.
75. Klafta JM, Olson JP. Emergent lung separation for management of pulmonary artery rupture. *Anesthesiology*. 1997;87(5):1248-1250.
76. Tayoro J, Dequin PF, Delhommais A, et al. Rupture of pulmonary artery induced by Swan-Ganz catheter: success of coil embolization. *Intensive Care Med*. 1997;23(2):198-200.
77. Rapoport J, Teres D, Steingrub J, et al. Patient characteristics and ICU organizational factors that influence frequency of pulmonary artery catheterization. *JAMA*. 2000;283(19):2559-2567.
78. Vincent JL, Dhainaut JF, Perret C, et al. Is the pulmonary artery catheter misused? A European view. *Crit Care Med* 1998;26(7):1283-1287.
79. Connors AF, Jr., Speroff T, Dawson NV, et al. The effectiveness of right heart catheterization in the initial care of critically ill patients. SUPPORT Investigators *JAMA*. 1996;276(11):889-897.
80. Dalen JE, Bone RC. Is it time to pull the pulmonary artery catheter? *JAMA*. 1996;276:916-918.
81. Polanczyk CA, Rohde LE, Goldman L, et al. Right heart catheterization and cardiac complications in patients undergoing noncardiac surgery: an observational study. *JAMA*. 2001;286(3):309-314.
82. Valentine RJ, Duke ML, Inman MH, et al. Effectiveness of pulmonary artery catheters in aortic surgery: a randomized trial. *J Vasc Surg*. 1998;27(2):203-211; discussion 211-202.
83. Barone JE, Tucker JB, Rassias D, et al. Routine perioperative pulmonary artery catheterization has no effect on rate of complications in vascular surgery: a meta-analysis. *Am Surg*. 2001;67(7):674-679.
84. Tuman KJ, McCarthy RJ, Spiess BD, et al. Effect of pulmonary artery catheterization on outcome in patients undergoing coronary artery surgery. *Anesthesiology*. 1989;70(2):199-206.
85. Binanay C, Califf RM, Hasselblad V, et al. Evaluation study of congestive heart failure and pulmonary artery catheterization effectiveness: the ESCAPE trial. *JAMA*. 2005;294(13):1625-1633.
86. Wheeler AP, Bernard GR, Thompson BT, et al. Pulmonary-artery versus central venous catheter to guide treatment of acute lung injury. *N Engl J Med*. 2006;354(21):2213-2224.
87. Rhodes A, Cusack RJ, Newman PJ, et al. A randomised, controlled trial of the pulmonary artery catheter in critically ill patients. *Intensive Care Med*. 2002;28(3):256-264.
88. Richard C, Warszawski J, Anguel N, et al. Early use of the pulmonary artery catheter and outcomes in patients with shock and acute respiratory distress syndrome: a randomized controlled trial. *JAMA*. 2003;290(20):2713-2720.
89. De Backer D, Creteur J, Vincent JL. Perioperative optimization and right heart catheterization: what technique in which patient? *Crit Care*. 2003;7(3):201-202.
90. Shah MR, Hasselblad V, Stevenson LW, et al. Impact of the pulmonary artery catheter in critically ill patients: meta-analysis of randomized clinical trials. *JAMA*. 2005;294(13):1664-1670.
91. Kavarana MN, Azimuddin K, Agarwal A, et al. Hemodynamic monitoring in the elderly undergoing elective colon resection for cancer. *Am Surg*. 2003;69(5):411-415.
92. Chittock DR, Dhingra VK, Ronco JJ, et al. Severity of illness and risk of death associated with pulmonary artery catheter use. *Crit Care Med*. 2004;32(4):911-915.
93. Friese RS, Shafi S, Gentilello LM. Pulmonary artery catheter use is associated with reduced mortality in severely injured patients: A National Trauma Data Bank analysis of 53,312 patients. *Crit Care Med*. 2006;34(6):1597-1601.
94. Zaidan JR, Freniere S. Use of a pacing pulmonary artery catheter during cardiac surgery. *Ann Thorac Surg*. 1983;35(6):633-636.
95. Bloos F, Reinhart K. Venous oximetry. *Intensive Care Med*. 2005;31(7):911-913.
96. Inomata S, Nishikawa T, Taguchi M. Continuous monitoring of mixed venous oxygen saturation for detecting alterations in cardiac output after discontinuation of cardiopulmonary bypass. *Br J Anaesth*. 1994;72(1):11-16.
97. Spiess BD, Tuman KJ, McCarthy RJ, et al. Oxygen consumption and mixed venous oxygen saturation monitoring during orthotopic liver transplantation. *J Clin Monit*. 1992;8(1):7-11.
98. Powelson JA, Maini BS, Bishop RL, et al. Continuous monitoring of mixed venous oxygen saturation during aortic operations. *Crit Care Med*. 1992;20(3):332-336.
99. Teboul JL, Graini L, Boujdaria R, et al. Cardiac index vs oxygen-derived parameters for rational use of dobutamine in patients with congestive heart failure. *Chest*. 1993;103(1):81-85.
100. Rivers E, Nguyen B, Havstad S, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med*. 2001;345(19):1368-1377.
101. Polonen P, Ruokonen E, Hippelainen M, et al. A prospective, randomized study of goal-oriented hemodynamic therapy in cardiac surgical patients. *Anesth Analg*. 2000;90(5):1052-1059.
102. Donati A, Loggi S, Preiser J-C, et al. Goal-directed intraoperative therapy reduces morbidity and length of hospital stay in high-risk surgical patients. *Chest*. 2007;132(6):1817-1824.
103. Gattinoni L, Brazzi L, Pelosi P, et al. A trial of goal-oriented hemodynamic therapy in critically ill patients. SvO2 Collaborative Group. *N Engl J Med*. 1995;333(16):1025-1032.
104. Reinhart K, Kuhn HJ, Hartog C, et al. Continuous central venous and pulmonary artery oxygen saturation monitoring in the critically ill. *Intensive Care Med*. 2004;30(8):1572-1578.

105. Scalea TM, Hartnett RW, Duncan AO, et al. Central venous oxygen saturation: a useful clinical tool in trauma patients. *J Trauma*. 1990;30(12):1539-1543.
106. Pearse R, Dawson D, Fawcett J, et al. Changes in central venous saturation after major surgery, and association with outcome. *Crit Care*. 2005;9(6):R694-R699.
107. Smetkin AA, Kirov MY, Kuzkov VV, et al. Single transpulmonary thermodilution and continuous monitoring of central venous oxygen saturation during off-pump coronary surgery. *Acta Anaesthesiol Scand*. 2009;53(4):505-514.
108. Vincent JL, Thirion M, Brimiouille S, et al. Thermodilution measurement of right ventricular ejection fraction with a modified pulmonary artery catheter. *Intensive Care Med*. 1986;12(1):33-38.
109. Kay HR, Afshari M, Barash P, et al. Measurement of ejection fraction by thermal dilution techniques. *J Surg Res*. 1983;34(4):337-346.
110. Boldt J, Kling D, Moosdorf R, et al. Influence of acute volume loading on right ventricular function after cardiopulmonary bypass. *Crit Care Med*. 1989;17(6):518-522.
111. Vincent JL, Gris P, Coffernils M, et al. Myocardial depression characterizes the fatal course of septic shock. *Surgery*. 1992;111(6):660-667.
112. Redl G, Germann P, Plattner H, et al. Right ventricular function in early septic shock states. *Intensive Care Med*. 1993;19(1):3-7.
113. Dhingra VK, Fenwick JC, Walley KR, et al. Lack of agreement between thermodilution and fick cardiac output in critically ill patients. *Chest*. 2002;122(3):990-997.
114. Singer M. What's in a beat? *Intensive Care Med*. 2003;29(10):1617-1620.
115. Chaney JC, Derdak S. Minimally invasive hemodynamic monitoring for the intensivist: current and emerging technology. *Crit Care Med*. 2002;30(10):2338-2345.
116. Reuter DA, Huang C, Edrich T, et al. Cardiac output monitoring using indicator-dilution techniques: basics, limits, and perspectives. *Anesth Analg*. 2010;110(3):799-811.
117. Wyse SD, Pfitzner J, Rees A, et al. Measurement of cardiac output by thermal dilution in infants and children. *Thorax*. 1975;30(3):262-265.
118. Nishikawa T, Dohi S. Errors in the measurement of cardiac output by thermodilution. *Can J Anaesth*. 1993;40(2):142-153.
119. Moise SF, Sinclair CJ, Scott DH. Pulmonary artery blood temperature and the measurement of cardiac output by thermodilution. *Anaesthesia*. 2002;57(6):562-566.
120. Jansen JR, Versprille A. Improvement of cardiac output estimation by the thermodilution method during mechanical ventilation. *Intensive Care Med*. 1986;12(2):71-79.
121. Wetzel RC, Latson TW. Major errors in thermodilution cardiac output measurement during rapid volume infusion. *Anesthesiology*. 1985;62(5):684-687.
122. Bazaral MG, Petre J, Novoa R. Errors in thermodilution cardiac output measurements caused by rapid pulmonary artery temperature decreases after cardiopulmonary bypass. *Anesthesiology*. 1992;77(1):31-37.
123. Latson TW, Whitten CW, O'Flaherty D. Ventilation, thermal noise, and errors in cardiac output measurements after cardiopulmonary bypass. *Anesthesiology*. 1993;79(6):1233-1243.
124. Heerd PM, Blessios GA, Beach ML, et al. Flow dependency of error in thermodilution measurement of cardiac output during acute tricuspid regurgitation. *J Cardiothorac Vasc Anesth*. 2001;15(2):183-187.
125. Artucio H, Hurtado J, Zimet L, et al. PEEP-induced tricuspid regurgitation. *Intensive Care Med*. 1997;23(8):836.
126. van Grondelle A, Ditchey RV, Groves BM, et al. Thermodilution method overestimates low cardiac output in humans. *Am J Physiol*. 1983;245(4):H690-692.
127. Hoeft A. Transpulmonary indicator dilution: an alternative approach for hemodynamic monitoring. In: Vincent J, ed. *Yearbook of Intensive Care and Emergency Medicine*. Berlin, Germany: Springer; 1995:593-605.
128. von Spiegel T, Wietasch G, Bursch J, et al. Cardiac output determination with transpulmonary thermodilution. An alternative to pulmonary catheterization? [in German]. *Anaesthesist*. 1996;45(11):1045-1050.
129. Segal E, Katzenelson R, Berkenstadt H, et al. Transpulmonary thermodilution cardiac output measurement using the axillary artery in critically ill patients. *J Clin Anesth*. 2002;14(3):210-213.
130. Mielck F, Buhre W, Hanekop G, et al. Comparison of continuous cardiac output measurements in patients after cardiac surgery. *J Cardiothorac Vasc Anesth*. 2003;17(2):211-216.
131. Linton R, Band D, O'Brien T, et al. Lithium dilution cardiac output measurement: a comparison with thermodilution. *Crit Care Med*. 1997;25(11):1796-1800.
132. Garcia-Rodriguez C, Pittman J, Cassell CH, et al. Lithium dilution cardiac output measurement: a clinical assessment of central venous and peripheral venous indicator injection. *Crit Care Med*. 2002;30(10):2199-2204.
133. Linton RA, Jonas MM, Tibby SM, et al. Cardiac output measured by lithium dilution and transpulmonary thermodilution in patients in a paediatric intensive care unit. *Intensive Care Med*. 2000;26(10):1507-1511.
134. Jaffe MB. Partial CO<sub>2</sub> rebreathing cardiac output—operating principles of the NICO system. *J Clin Monit Comput*. 1999;15(6):387.
135. Osterlund B, Gedeon A, Krill P, et al. A new method of using gas exchange measurements for the noninvasive determination of cardiac output: clinical experiences in adults following cardiac surgery. *Acta Anaesthesiol Scand*. 1995;39(6):727-732.
136. van Heerden PV, Baker S, Lim SI, et al. Clinical evaluation of the non-invasive cardiac output (NICO) monitor in the intensive care unit. *Anaesth Intensive Care*. 2000;28(4):427-430.
137. Yelderman M. Continuous measurement of cardiac output with the use of stochastic system identification techniques. *J Clin Monit*. 1990;6(4):322-332.
138. Bottiger BW, Rauch H, Bohrer H, et al. Continuous versus intermittent cardiac output measurement in cardiac surgical patients undergoing hypothermic cardiopulmonary bypass. *J Cardiothorac Vasc Anesth*. 1995;9(4):405-411.
139. Botero M, Kirby D, Lobato EB, et al. Measurement of cardiac output before and after cardiopulmonary bypass: comparison among aortic transit-time ultrasound, thermodilution, and noninvasive partial CO<sub>2</sub> rebreathing. *J Cardiothorac Vasc Anesth*. 2004;18(5):563.
140. Siegel LC, Hennessy MM, Pearl RG. Delayed time response of the continuous cardiac output pulmonary artery catheter. *Anesth Analg*. 1996;83(6):1173-1177.
141. Lichtenthal PR, Wade LD. Continuous cardiac output measurements. *J Cardiothorac Vasc Anesth*. 1994;8(6):668-670.

142. Wesseling KH, Jansen JR, Settels JJ, et al. Computation of aortic flow from pressure in humans using a nonlinear, three-element model. *J Appl Physiol*. 1993;74(5):2566-2573.
143. Linton NW, Linton RA. Estimation of changes in cardiac output from the arterial blood pressure waveform in the upper limb. *Br J Anaesth*. 2001;86(4):486-496.
144. van Lieshout JJ, Wesseling KH. Continuous cardiac output by pulse contour analysis? *Br J Anaesth*. 2001;86(4):467-469.
145. Camporota L, Beale R. Pitfalls in haemodynamic monitoring based on the arterial pressure waveform. *Crit Care*. 2010;14(2):124.
146. Pittman J, Bar-Yosef S, SumPing J, et al. Continuous cardiac output monitoring with pulse contour analysis: a comparison with lithium indicator dilution cardiac output measurement. *Crit Care Med*. 2005;33(9):2015-2021.
147. Weissman C, Ornstein EJ, Young WL. Arterial pulse contour analysis trending of cardiac output: hemodynamic manipulations during cerebral arteriovenous malformation resection. *J Clin Monit*. 1993;9(5):347-353.
148. Singer M. Esophageal Doppler monitoring of aortic blood flow: beat-by-beat cardiac output monitoring. *Int Anesthesiol Clin*. 1993;31(3):99-125.
149. Odenstedt H, Aneman A, Oi Y, et al. Descending aortic blood flow and cardiac output: a clinical and experimental study of continuous oesophageal echo-Doppler flowmetry. *Acta Anaesthesiol Scand*. 2001;45(2):180-187.
150. Monnet X, Rienzo M, Osman D, et al. Esophageal Doppler monitoring predicts fluid responsiveness in critically ill ventilated patients. *Intensive Care Med*. 2005;31(9):1195-1201.
151. Dark PM, Singer M. The validity of trans-esophageal Doppler ultrasonography as a measure of cardiac output in critically ill adults. *Intensive Care Med*. 2004;30(11):2060-2066.
152. Phan TD, Ismail H, Heriot AG, et al. Improving perioperative outcomes: fluid optimization with the esophageal Doppler monitor, a metaanalysis and review. *J Am Coll Surg*. 2008;207(6):935-941.
153. Schlichtig R, Bowles SA. Distinguishing between aerobic and anaerobic appearance of dissolved CO<sub>2</sub> in intestine during low flow. *J Appl Physiol*. 1994;76(6):2443-2451.
154. Kolkman JJ, Otte JA, Groeneveld AB. Gastrointestinal luminal PCO<sub>2</sub> tonometry: an update on physiology, methodology and clinical applications. *Br J Anaesth*. 2000;84(1):74-86.
155. Schlichtig R, Mehta N, Gayowski TJ. Tissue-arterial PCO<sub>2</sub> difference is a better marker of ischemia than intramural pH (pHi) or arterial pH-pHi difference. *J Crit Care*. 1996;11(2):51-56.
156. Adar R, Franklin A, Spark RF, et al. Effect of dehydration and cardiac tamponade on superior mesenteric artery flow: role of vasoactive substances. *Surgery*. 1976;79(5):534-543.
157. Hamilton-Davies C, Mythen MG, Salmon JB, et al. Comparison of commonly used clinical indicators of hypovolaemia with gastrointestinal tonometry. *Intensive Care Med*. 1997;23(3):276-281.
158. Ivatury RR, Simon RJ, Havriliak D, et al. Gastric mucosal pH and oxygen delivery and oxygen consumption indices in the assessment of adequacy of resuscitation after trauma: a prospective, randomized study. *J Trauma*. 1995;39(1):128-134; discussion 134-126.
159. Mythen MG, Webb AR. The role of gut mucosal hypoperfusion in the pathogenesis of post-operative organ dysfunction. *Intensive Care Med*. 1994;20(3):203-209.
160. Bennett-Guerrero E, Panah MH, Bodian CA, et al. Automated detection of gastric luminal partial pressure of carbon dioxide during cardiovascular surgery using the Tonocap. *Anesthesiology*. 2000;92(1):38-45.
161. Lebuffe G, Vallet B, Takala J, et al. A European, multicenter, observational study to assess the value of gastric-to-end tidal PCO<sub>2</sub> difference in predicting postoperative complications. *Anesth Analg*. 2004;99(1):166-172.
162. Gutierrez G, Palizas F, Doglio G, et al. Gastric intramucosal pH as a therapeutic index of tissue oxygenation in critically ill patients. *Lancet*. 1992;339(8787):195-199.
163. Poeze M, Takala J, Greve JW, et al. Pre-operative tonometry is predictive for mortality and morbidity in high-risk surgical patients. *Intensive Care Med*. 2000;26(9):1272-1281.
164. Pargger H, Hampl KF, Christen P, et al. Gastric intramucosal pH-guided therapy in patients after elective repair of infrarenal abdominal aneurysms: is it beneficial? *Intensive Care Med*. 1998;24(8):769-776.
165. Gomersall CD, Joynt GM, Freebairn RC, et al. Resuscitation of critically ill patients based on the results of gastric tonometry: a prospective, randomized, controlled trial. *Crit Care Med*. 2000;28(3):607-614.
166. Landesberg G, Mosseri M, Zahger D, et al. Myocardial infarction after vascular surgery: the role of prolonged stress-induced, ST depression-type ischemia. *J Am Coll Cardiol*. 2001;37(7):1839-1845.
167. Mangano DT, Browner WS, Hollenberg M, et al. Association of perioperative myocardial ischemia with cardiac morbidity and mortality in men undergoing noncardiac surgery. The Study of Perioperative Ischemia Research Group *N Engl J Med*. 1990;323(26):1781-1788.
168. Kaplan JA, Wells PH. Early diagnosis of myocardial ischemia using the pulmonary arterial catheter. *Anesth Analg*. 1981;60:789-793.
169. McLaurin LP, Rolett EL, Grossman W. Impaired left ventricular relaxation during pacing-induced ischemia. *Am J Cardiol*. 1973;32:751-757.
170. van Daele MERM, Sutherland GR, Mitchell MM, et al. Do changes in pulmonary capillary wedge pressure adequately reflect myocardial ischemia during anesthesia: a correlative preoperative hemodynamic, electrocardiographic, and transesophageal echocardiographic study. *Circulation*. 1990;81:865-871.
171. Hall RI, O'Regan N, Gardner M. Detection of intraoperative myocardial ischaemia—a comparison among electrocardiographic, myocardial metabolic, and haemodynamic measurements in patients with reduced ventricular function. *Can J Anaesth*. 1995;42(6):487-494.
172. Pinsky MR. Clinical significance of pulmonary artery occlusion pressure. *Intensive Care Med*. 2003;29(2):175-178.
173. Bernard GR, Artigas A, Brigham KL, et al. The American-European Consensus Conference on ARDS. Definitions, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med*. 1994;149(3 Pt 1):818-824.
174. Nunes S, Ruokonen E, Takala J. Pulmonary capillary pressures during the acute respiratory distress syndrome. *Intensive Care Med*. 2003;29(12):2174-2179.
175. Mitchell JP, Schuller D, Calandrino FS. Improved outcome based on fluid management in critically ill patients requiring pulmonary artery catheterization. *Am Rev Respir Dis*. 1992;145(5):990-998.
176. Wiedemann HP, Wheeler AP, Bernard GR, et al. Comparison of two fluid-management strategies in acute lung injury. *N Engl J Med*. 2006;354(24):2564-2575.

177. Michard F. Bedside assessment of extravascular lung water by dilution methods: temptations and pitfalls. *Crit Care Med.* 2007;35(4):1186-1192.
178. Sakka SG, Klein M, Reinhart K, et al. Prognostic value of extravascular lung water in critically ill patients. *Chest.* 2002;122(6):2080-2086.
179. Matejovic M, Krouzicky A, Rokyta R Jr, et al. Fluid challenge in patients at risk for fluid loading-induced pulmonary edema. *Acta Anaesthesiol Scand.* 2004;48(1):69-73.
180. Kass DA, Maughan WL, Guo ZM, et al. Comparative influence of load versus inotropic states on indexes of ventricular contractility: experimental and theoretical analysis based on pressure-volume relationships. *Circulation.* 1987;76(6):1422-1436.
181. Forrester JS, Diamond G, Chatterjee K, et al. Medical therapy of acute myocardial infarction by application of hemodynamic subsets (first of two parts). *N Engl J Med.* 1976;295(24):1356-1362.
182. Michard F, Reuter DA. Assessing cardiac preload or fluid responsiveness? It depends on the question we want to answer. *Intensive Care Med.* 2003;29(8):1396.
183. Ramsey JD, Tisdale LA. Use of ventricular stroke work index and ventricular function curves in assessing myocardial contractility. *Crit Care Nurse.* 1995;15(1):61-67.
184. Ognibene FP, Parker MM, Natanson C, et al. Depressed left ventricular performance. Response to volume infusion in patients with sepsis and septic shock. *Chest.* 1988;93(5):903-910.
185. Poortmans G, Schupfer G, Roosens C, et al. Transesophageal echocardiographic evaluation of left ventricular function. *J Cardiothorac Vasc Anesth.* 2000;14(5):588-598.
186. Thys DM, Hillel Z. Left ventricular performance indices by transesophageal Doppler. *Anesthesiology.* 1988;69(5):728-737.
187. Wisner-Eutenier, Lichtwarck-Aschoff M, Zimmermann G, et al. Evaluation of the cardiac function index as a new bedside indicator of cardiac performance. *Intensive Care Med.* 1994;20(Suppl 2):S21.
188. Moore TD, Frenneaux MP, Sas R, et al. Ventricular interaction and external constraint account for decreased stroke work during volume loading in CHF. *Am J Physiol Heart Circ Physiol.* 2001;281(6):H2385-2391.
189. Holte K, Sharrock NE, Kehlet H. Pathophysiology and clinical implications of perioperative fluid excess. *Br J Anaesth.* 2002;89(4):622-632.
190. Grocott MP, Mythen MG, Gan TJ. Perioperative fluid management and clinical outcomes in adults. *Anesth Analg.* 2005;100(4):1093-1106.
191. Connolly CM, Kramer GC, Hahn RG, et al. Isoflurane but not mechanical ventilation promotes extravascular fluid accumulation during crystalloid volume loading. *Anesthesiology.* 2003;98(3):670-681.
192. Michard F, Teboul JL. Predicting fluid responsiveness in ICU patients: a critical analysis of the evidence. *Chest.* 2002;121(6):2000-2008.
193. Tavernier B, Makhotine O, Lebuffe G, et al. Systolic pressure variation as a guide to fluid therapy in patients with sepsis-induced hypotension. *Anesthesiology.* 1998;89(6):1313-1321.
194. McGee S, Abernethy WB 3rd, Simel DL. The rational clinical examination. Is this patient hypovolemic? *JAMA.* 1999;281(11):1022-1029.
195. Kellen M, Aronson S, Roizen MF, et al. Predictive and diagnostic tests of renal failure: a review. *Anesth Analg.* 1994;78(1):134-142.
196. Vincent J-L, Weil MH. Fluid challenge revisited. *Crit Care Med.* 2006;34(5):1333-1337.
197. Boulain T, Achard JM, Teboul JL, et al. Changes in BP induced by passive leg raising predict response to fluid loading in critically ill patients. *Chest.* 2002;121(4):1245-1252.
198. Monnet X, Rienzo M, Osman D, et al. Passive leg raising predicts fluid responsiveness in the critically ill. *Crit Care Med.* 2006;34(5):1402-1407.
199. Kumar A, Anel R, Bunnell E, et al. Pulmonary artery occlusion pressure and central venous pressure fail to predict ventricular filling volume, cardiac performance, or the response to volume infusion in normal subjects. *Crit Care Med.* 2004;32(3):691-699.
200. Venn R, Steele A, Richardson P, et al. Randomized controlled trial to investigate influence of the fluid challenge on duration of hospital stay and perioperative morbidity in patients with hip fractures. *Br J Anaesth.* 2002;88(1):65-71.
201. Reuse C, Vincent JL, Pinsky MR. Measurements of right ventricular volumes during fluid challenge. *Chest.* 1990;98(6):1450-1454.
202. Chang MC, Blinman TA, Rutherford EJ, et al. Preload assessment in trauma patients during large-volume shock resuscitation. *Arch Surg.* 1996;131(7):728-731.
203. Wagner JG, Leatherman JW. Right ventricular end-diastolic volume as a predictor of the hemodynamic response to a fluid challenge. *Chest.* 1998;113(4):1048-1054.
204. Cheatham ML, Nelson LD, Chang MC, et al. Right ventricular end-diastolic volume index as a predictor of preload status in patients on positive end-expiratory pressure. *Crit Care Med.* 1998;26(11):1801-1806.
205. Michard F, Alaya S, Zarka V, et al. Global end-diastolic volume as an indicator of cardiac preload in patients with septic shock. *Chest.* 2003;124(5):1900-1908.
206. Wiesenack C, Prasser C, Keyl C, et al. Assessment of intrathoracic blood volume as an indicator of cardiac preload: single transpulmonary thermodilution technique versus assessment of pressure preload parameters derived from a pulmonary artery catheter. *J Cardiothorac Vasc Anesth.* 2001;15(5):584-588.
207. Della Rocca G, Costa GM, Coccia C, et al. Preload index: pulmonary artery occlusion pressure versus intrathoracic blood volume monitoring during lung transplantation. *Anesth Analg.* 2002;95(4):835-843.
208. Marik PE, Cavallazzi R, Vasu T, et al. Dynamic changes in arterial waveform derived variables and fluid responsiveness in mechanically ventilated patients: a systematic review of the literature. *Crit Care Med.* 2009;37(9):2642-2647.
209. Thys DM, Hillel Z, Goldman ME, et al. A comparison of hemodynamic indices derived by invasive monitoring and two-dimensional echocardiography. *Anesthesiology.* 1987;67(5):630-634.
210. Vieillard-Baron A, Chergui K, Rabiller A, et al. Superior vena caval collapsibility as a gauge of volume status in ventilated septic patients. *Intensive Care Med.* 2004;30(9):1734-1739.
211. Barbier C, Loubieres Y, Schmit C, et al. Respiratory changes in inferior vena cava diameter are helpful in predicting fluid responsiveness in ventilated septic patients. *Intensive Care Med.* 2004;30(9):1740-1746.
212. Michard F. Changes in arterial pressure during mechanical ventilation. *Anesthesiology.* 2005;103(2):419-428.
213. Perel A, Pizov R, Coté S. Systolic blood pressure variation is a sensitive indicator of hypovolemia in ventilated dogs subjected to graded hemorrhage. *Anesthesiology.* 1987;67(4):498-502.

214. Coriat P, Vrillon M, Perel A, et al. A comparison of systolic blood pressure variations and echocardiographic estimates of end-diastolic left ventricular size in patients after aortic surgery *Anesth Analg*. 1994;78(1):46-53.
215. Vieillard-Baron A, Chergui K, Augarde R, et al. Cyclic changes in arterial pulse during respiratory support revisited by Doppler echocardiography. *Am J Respir Crit Care Med*. 2003;168(6):671-676.
216. Preisman S, DiSegni E, Vered Z, et al. Left ventricular preload and function during graded haemorrhage and retransfusion in pigs: analysis of arterial pressure waveform and correlation with echocardiography. *Br J Anaesth*. 2002;88(5):716-718.
217. Vieillard-Baron A, Augarde R, Prin S, et al. Influence of superior vena caval zone condition on cyclic changes in right ventricular outflow during respiratory support. *Anesthesiology*. 2001;95(5):1083-1088.
218. Kramer A, Zygun D, Hawes H, et al. Pulse pressure variation predicts fluid responsiveness following coronary artery bypass surgery. *Chest*. 2004;126(5):1563-1568.
219. Michard F, Boussat S, Chemla D, et al. Relation between respiratory changes in arterial pulse pressure and fluid responsiveness in septic patients with acute circulatory failure. *Am J Respir Crit Care Med*. 2000;162(1):134-138.
220. Berkenstadt H, Margalit N, Hadani M, et al. Stroke volume variation as a predictor of fluid responsiveness in patients undergoing brain surgery. *Anesth Analg*. 2001;92(4):984-989.
221. Hofer CK, Muller SM, Furrer L, et al. Stroke volume and pulse pressure variation for prediction of fluid responsiveness in patients undergoing off-pump coronary artery bypass grafting. *Chest*. 2005;128(2):848-854.
222. Natalini G, Rosano A, Franceschetti ME, et al. Variations in arterial blood pressure and photoplethysmography during mechanical ventilation. *Anesth Analg*. 2006;103(5):1182-1188.
223. Natalini G, Rosano A, Taranto M, et al. Arterial versus plethysmographic dynamic indices to test responsiveness for testing fluid administration in hypotensive patients: a clinical trial. *Anesth Analg*. 2006;103(6):1478-1484.
224. Landsverk SA, Hoiseth LO, Kvandal P, et al. Poor agreement between respiratory variations in pulse oximetry photoplethysmographic waveform amplitude and pulse pressure in intensive care unit patients. *Anesthesiology*. 2008;109(5):849-855.
225. Cannesson M, Delannoy B, Morand A, et al. Does the Pleth variability index indicate the respiratory-induced variation in the plethysmogram and arterial pressure waveforms? *Anesth Analg*. 2008;106(4):1189-1194.
226. Cannesson M, Desebbe O, Rosamel P, et al. Pleth variability index to monitor the respiratory variations in the pulse oximeter plethysmographic waveform amplitude and predict fluid responsiveness in the operating theatre. *Br J Anaesth*. 2008;101(2):200-206.
227. Rooke GA, Schwid HA, Shapira Y. The effect of graded hemorrhage and intravascular volume replacement on systolic pressure variation in humans during mechanical and spontaneous ventilation. *Anesth Analg*. 1995;80(5):925-932.
228. De Backer D, Heenen S, Piagnerelli M, et al. Pulse pressure variations to predict fluid responsiveness: influence of tidal volume. *Intensive Care Med*. 2005;31(4):517-523.
229. Perel A, Minkovich L, Preisman S, et al. Assessing fluid-responsiveness by a standardized ventilatory maneuver: the respiratory systolic variation test. *Anesth Analg*. 2005;100(4):942-945.
230. Michard F, Lopes M, Auler J. Pulse pressure variation: beyond the fluid management of patients with shock. *Crit Care*. 2007;11:131.
231. Kumar A, Anel R, Bunnell E, et al. Preload-independent mechanisms contribute to increased stroke volume following large volume saline infusion in normal volunteers: a prospective interventional study. *Crit Care*. 2004;8(3):R128-136.
232. Michard F. Volume management using dynamic parameters: the good, the bad, and the ugly. *Chest*. 2005;128(4):1902-1903.
233. Uusaro A, Ruokonen E, Takala J. Splanchnic oxygen transport after cardiac surgery: evidence for inadequate tissue perfusion after stabilization of hemodynamics. *Intensive Care Med*. 1996;22(1):26-33.
234. Rady MY, Rivers EP, Martin GB, et al. Continuous central venous oximetry and shock index in the emergency department: use in the evaluation of clinical shock. *Am J Emerg Med*. 1992;10(6):538-541.
235. Bland RD, Shoemaker WC, Abraham E, et al. Hemodynamic and oxygen transport patterns in surviving and nonsurviving postoperative patients. *Crit Care Med*. 1985;13(2):85-90.
236. Shoemaker WC, Appel PL, Kram HB. Role of oxygen debt in the development of organ failure sepsis, and death in high-risk surgical patients. *Chest*. 1992;102(1):208-215.
237. Kaplan LJ, Kellum JA. Initial pH, base deficit, lactate, anion gap, strong ion difference, and strong ion gap predict outcome from major vascular injury. *Crit Care Med*. 2004;32(5):1120-1124.
238. Cheifetz IM, Kern FH, Schulman SR, et al. Serum lactates correlate with mortality after operations for complex congenital heart disease. *Ann Thorac Surg*. 1997;64(3):735-738.
239. Rossi AF, Khan DM, Hannan R, et al. Goal-directed medical therapy and point-of-care testing improve outcomes after congenital heart surgery. *Intensive Care Med*. 2005;31(1):98-104.
240. De Backer D. Lactic acidosis. *Intensive Care Med*. 2003;29(5):699-702.
241. Ronco JJ, Fenwick JC, Tweeddale MG, et al. Identification of the critical oxygen delivery for anaerobic metabolism in critically ill septic and nonseptic humans. *JAMA*. 1993;270(14):1724-1730.
242. Polonen P, Hippelainen M, Takala R, et al. Relationship between intra- and postoperative oxygen transport and prolonged intensive care after cardiac surgery: a prospective study. *Acta Anaesthesiol Scand*. 1997;41(7):810-817.
243. Yu M, Burchell S, Hasaniya NW, et al. Relationship of mortality to increasing oxygen delivery in patients > or = 50 years of age: a prospective, randomized trial. *Crit Care Med*. 1998;26(6):1011-1019.
244. Vincent JL, De Backer D. Cardiac output measurement: is least invasive always the best? *Crit Care Med*. 2002;30(10):2380-2382.
245. Siegemund M, van Bommel J, Ince C. Assessment of regional tissue oxygenation. *Intensive Care Med*. 1999;25(10):1044.
246. Toledo Maciel A, Creteur J, Vincent J-L. Tissue capnometry: does the answer lie under the tongue? *Intensive Care Med*. 2004;30(12):2157.
247. Lees N, Hamilton M, Rhodes A. Clinical review: goal-directed therapy in high risk surgical patients. *Crit Care*. 2009;13(5):231.
248. Shoemaker WC, Appel PL, Kram HB, et al. Prospective trial of supranormal values of survivors as therapeutic goals in high-risk surgical patients. *Chest*. 1988;94(6):1176-1186.

249. Giglio MT, Marucci M, Testini M, et al. Goal-directed haemodynamic therapy and gastrointestinal complications in major surgery: a meta-analysis of randomized controlled trials. *Br J Anaesth*. 2009;103(5):637-646.
250. Brienza N, Giglio MT, Marucci M, et al. Does perioperative hemodynamic optimization protect renal function in surgical patients? A meta-analytic study. *Crit Care Med*. 2009;37(6):2079-2090.
251. Boyd O, Bennett ED. Enhancement of perioperative tissue perfusion as a therapeutic strategy for major surgery. *New Horiz*. 1996;4(4):453-465.
252. Takala J, Meier-Hellmann A, Eddleston J, et al. Effect of dopexamine on outcome after major abdominal surgery: a prospective, randomized, controlled multicenter study. European Multicenter Study Group on Dopexamine in Major Abdominal Surgery. *Crit Care Med*. 2000;28(10):3417-3423.
253. Lobo S, Lobo F, Polachini C, et al. Prospective, randomized trial comparing fluids and dobutamine optimization of oxygen delivery in high-risk surgical patients. *Crit Care*. 2006;10:R72.
254. Alia I, Esteban A, Gordo F, et al. A randomized and controlled trial of the effect of treatment aimed at maximizing oxygen delivery in patients with severe sepsis or septic shock. *Chest*. 1999;115(2):453-461.
255. Hayes MA, Timmins AC, Yau EH, et al. Oxygen transport patterns in patients with sepsis syndrome or septic shock: influence of treatment and relationship to outcome *Crit Care Med*. 1997;25(6):926-936.
256. Balogh Z, McKinley BA, Cocanour CS, et al. Supranormal trauma resuscitation causes more cases of abdominal compartment syndrome. *Arch Surg*. 2003;138(6):637-642; discussion 642-633.
257. Brandstrup B, Tonnesen H, Beier-Holgersen R, et al. Effects of intravenous fluid restriction on postoperative complications: comparison of two perioperative fluid regimens. *Ann Surg*. 2003;238:641-648.
258. Nisanevich V, Felsenstein I, Almogy G, et al. Effect of intraoperative fluid management on outcome after intraabdominal surgery. *Anesthesiology*. 2005;103:25-32.
259. Senagore AJ, Emery T, Luchtefeld M, et al. Fluid management for laparoscopic colectomy: a prospective, randomized assessment of goal-directed administration of balanced salt solution or hetastarch coupled with an enhanced recovery program. *Dis Colon Rectum*. 2009;52(12):1935-1940.
260. Kimberger O, Arnberger M, Brandt S, et al. Goal-directed colloid administration improves the microcirculation of healthy and perianastomotic colon. *Anesthesiology*. 2009;110(3):496-504.
261. Kern JW, Shoemaker WC. Meta-analysis of hemodynamic optimization in high-risk patients. *Crit Care Med*. 2002;30(8):1686-1692.
262. Cook TM, Day CJ. Hospital mortality after urgent and emergency laparotomy in patients aged 65 yr and over. Risk and prediction of risk using multiple logistic regression analysis. *Br J Anaesth*. 1998;80(6):776-781.
263. Older P, Hall A, Hader R. Cardiopulmonary exercise testing as a screening test for perioperative management of major surgery in the elderly. *Chest*. 1999;116(2):355-362.
264. Girish M, Trayner E, Jr., Dammann O, et al. Symptom-limited stair climbing as a predictor of postoperative cardiopulmonary complications after high-risk surgery. *Chest*. 2001;120(4):1147-1151.
265. Fenwick E, Wilson J, Sculpher M, et al. Pre-operative optimisation employing dopexamine or adrenaline for patients undergoing major elective surgery: a cost-effectiveness analysis. *Intensive Care Med*. 2002;28(5):599-608.
266. Mark JB. *Atlas of Cardiovascular Monitoring*. New York, NY: Churchill Livingstone; 1998.
267. Boyd O, Grounds RM, Bennett ED. A randomized clinical trial of the effect of deliberate perioperative increase of oxygen delivery on mortality in high-risk surgical patients. *JAMA*. 1993;270(22):2699-2707.
268. Wilson J, Woods I, Fawcett J, et al. Reducing the risk of major elective surgery: randomised controlled trial of preoperative optimisation of oxygen delivery. *B Med J*. 1999;318:1099-1103.
269. Pearse R, Dawson D, Fawcett J, et al. Early goal directed therapy reduces morbidity and length of hospital stay following high-risk surgery. *Crit Care*. 2005;9(Suppl 1):P45.
270. Mythen MG, Webb AR. Perioperative plasma volume expansion reduces the incidence of gut mucosal hypoperfusion during cardiac surgery. *Arch Surg*. 1995;130(4):423-429.
271. Gan TJ, Soppitt A, Maroof M, et al. Goal-directed intraoperative fluid administration reduces length of hospital stay after major surgery. *Anesthesiology*. 2002;97(4):820-826.
272. Wakeling HG, McFall MR, Jenkins CS, et al. Intraoperative oesophageal Doppler guided fluid management shortens postoperative hospital stay after major bowel surgery. *Br J Anaesth*. 2005;95(5):634-642.
273. Lopes M, Oliveira M, Pereira V, et al. Goal-directed fluid management based on pulse pressure variation monitoring during high-risk surgery: a pilot randomized controlled trial. *Crit Care*. 2007;11(5):R100.
274. Mayer J, Boldt J, Mengistu A, et al. Goal-directed intraoperative therapy based on autocalibrated arterial pressure waveform analysis reduces hospital stay in high-risk surgical patients: a randomized, controlled trial. *Crit Care*. 2010;14(1):R18.