

Chapter 33

REFERENCES

1. Martin JH. The collective electrical behavior of cortical neurons: The electroencephalogram and the mechanisms of epilepsy. In: Kandel ER, Schwartz JH, Jessel TM, eds. *Principles of Neural Science*. 3rd ed. New York, NY: Elsevier; 1991:777-791.
2. Rampil IJ. A primer for EEG signal processing in anesthesia. *Anesthesiology*. 1998;89:980-1002.
3. Lukatch H, Greenwald S. Cerebral cortex-anesthetic action on the electroencephalogram. In: Antognini J, Carstens E, Raines D, eds. *Neural Mechanisms of Anesthesia*. Humana: Totowa, NJ; 2003:93-112.
4. McPherson R. Intraoperative neurologic monitoring. In: Longnecker D, Tinker J, Morgan E, eds. *Principles and Practice of Anesthesiology*. St. Louis, MO: Mosby; 1998:883-906.
5. Amzica F, Steriade M. Electrophysiological correlates of sleep delta waves. *Electroencephalogr Clin Neurophysiol* 1998;107:69-83.
6. Sebel PS, Lang E, Rampil IJ, et al. A multicenter study of bispectral electroencephalogram analysis for monitoring anesthetic effect. *Anesth Analg* 1997;84:891-899.
7. Rabinovici GD, Lukatch HS, MacIver MB. Hypoglycemic and hypoxic modulation of cortical micro-EEG activity in rat brain slices. *Clin Neurophysiol* 2000;111:112-121.
8. Luu P, Tucker DM, Englander R, et al. Localizing acute stroke-related EEG changes: assessing the effects of spatial undersampling. *J Clin Neurophysiol* 2001;18:302-317.
9. Chiappa KH, Burke SR, Young RR. Results of electroencephalographic monitoring during 367 carotid endarterectomies. Use of a dedicated minicomputer. *Stroke* 1979;10:381-388.
10. Bond R, Rerkasem K, Rothwell PM. Routine or selective carotid artery shunting for carotid endarterectomy (and different methods of monitoring in selective shunting). *Stroke* 2003;34:824-825.
11. Lavine SD, Masri LS, Levy ML, Giannotta SL. Temporary occlusion of the middle cerebral artery in intracranial aneurysm surgery: time limitation and advantage of brain protection. *J Neurosurg* 1997;87:817-824.
12. Melgar MA, Mariwalla N, Madhusudan H, Weinand M. Carotid endarterectomy without shunt: the role of cerebral metabolic protection. *Neurol Res* 2005;27:850-856.
13. Bush RL, Kougiass P, Guerrero MA, et al. A comparison of carotid artery stenting with neuroprotection versus carotid endarterectomy under local anesthesia. *Am J Surg* 2005;190:696-700.
14. Roach GW, Newman MF, Murkin JM, et al. Ineffectiveness of burst suppression therapy in mitigating perioperative cerebrovascular dysfunction. Multicenter Study of Perioperative Ischemia (McSPI) Research Group. *Anesthesiology* 1999;90:1255-1264.
15. Newman MF, Croughwell ND, White WD, et al. Pharmacologic electroencephalographic suppression during cardiopulmonary bypass: a comparison of thiopental and isoflurane. *Anesth Analg* 1998;86:246-251.
16. Sigl JC, Chamoun NG. An introduction to bispectral analysis for the electroencephalogram. *J Clin Monit* 1994;10:392-404.
17. Vanluchene AL, Vereecke H, Thas O, et al. Spectral entropy as an electroencephalographic measure of anesthetic drug effect: a comparison with bispectral index and processed midlatency auditory evoked response. *Anesthesiology* 2004;101:34-42.
18. Viertio-Oja H, Maja V, Sarkela M, et al. Description of the Entropy algorithm as applied in the Datex-Ohmeda S/5 Entropy Module. *Acta Anaesthesiol Scand* 2004;48:154-161.
19. Sleight JW, Andrzejewski J, Steyn-Ross A, Steyn-Ross M: The bispectral index: a measure of depth of sleep? *Anesth Analg* 1999;88:659-661.
20. Miller A, Sleight JW, Barnard J, Steyn-Ross DA. Does bispectral analysis of the electroencephalogram add anything but complexity? *Br J Anaesth* 2004;92:8-13.
21. Shannon CE. A mathematical theory of communication. *Bell System Techn J* 1948; 27 (379-423): 623-56.
22. Johnson RW, Shore JE. Which is the better entropy expression for speech processing: $-S \log S$ or $\log S$? *IEEE Trans Acoust* 1984; ASSP-32: 129-37.
23. White PF, Tang J, Romero GF, et al. A comparison of state and response entropy versus bispectral index values during the perioperative period. *Anesth Analg* 2006;102:160-167.
24. Wheeler P, Hoffman WE, Baughman VL, Koenig H. Response entropy increases during painful stimulation. *J Neurosurg Anesthesiol* 2005;17:86-90.
25. Anonymous. National Institutes of Health Consensus Development Conference Statement on Acoustic Neuroma, December 11-13, 1991. The Consensus Development Panel. *Arch Neurol* 1994;51:201-207.
26. Toleikis JR. Intraoperative monitoring using somatosensory evoked potentials. A position statement by the American Society of Neurophysiological Monitoring. *J Clin Monit Comput* 2005;19:241-258.
27. York DH. Somatosensory evoked potentials in man: differentiation of spinal pathways responsible for conduction from the forelimb vs hindlimb. *Prog Neurobiol* 1985;25:1-25.

28. Powers SK, Bolger CA, Edwards MS. Spinal cord pathways mediating somatosensory evoked potentials. *J Neurosurg* 1982;57:472-482.
29. Banoub M, Tetzlaff JE, Schubert A. Pharmacologic and physiologic influences affecting sensory evoked potentials: implications for perioperative monitoring. *Anesthesiology* 2003;99:716-737.
30. Sloan TB. Anesthetic effects on electrophysiologic recordings. *J Clin Neurophysiol* 1998;15:217-226.
31. Huotari AM, Koskinen M, Suominen K, et al. Evoked EEG patterns during burst suppression with propofol. *Br J Anaesth* 2004;92:18-24.
32. Drummond JC, Todd MM, Schubert A, Sang H. Effect of the acute administration of high dose pentobarbital on human brain stem auditory and median nerve somatosensory evoked responses. *Neurosurgery* 1987;20:830-835.
33. Hall JW. Anatomy and physiology. In: Hall J, ed. *Handbook of Auditory Evoked Responses*. Boston, MA: Allyn and Bacon; 1992:41-69.
34. Morimoto Y, Sakabe T. [Auditory evoked potentials]. *Masui* 2006;55:314-321.
35. Schneider G, Nahm W, Kochs EF, et al. Quality of perioperative AEP—variability of expert ratings. *Br J Anaesth* 2003;91:905-908.
36. Smith WD, Dutton RC, Smith NT. Measuring the performance of anesthetic depth indicators. *Anesthesiology* 1996;84:38-51.
37. Alpiger S, Helbo-Hansen HS, Vach W, Ording H. Efficacy of A-line AEP Monitor as a tool for predicting acceptable tracheal intubation conditions during sevoflurane anaesthesia. *Br J Anaesth* 2005;94:601-606.
38. Hall JW. Overview of auditory evoked responses: past, present and future. In: Hall J, ed. *Handbook of Auditory Evoked Responses*. Boston, MA: Allyn and Bacon; 1992:3-40.
39. Drouot X, Nguyen JP, Peschanski M, Lefaucheur JP. The antalgic efficacy of chronic motor cortex stimulation is related to sensory changes in the painful zone. *Brain* 2002;125:1660-1664.
40. Lefaucheur JP. Transcranial magnetic stimulation in the management of pain. *Suppl Clin Neurophysiol* 2004;57:737-748.
41. Nguyen JP, Lefaucheur JP, Le Guerinel C, et al. Motor cortex stimulation in the treatment of central and neuropathic pain. *Arch Med Res* 2000;31:263-265.
42. Brown JA, Pilitsis JG. Motor cortex stimulation for central and neuropathic facial pain: a prospective study of 10 patients and observations of enhanced sensory and motor function during stimulation. *Neurosurgery* 2005;56:290-297; discussion 290-297.
43. Sala F, Krzan MJ, Deletis V. Intraoperative neurophysiological monitoring in pediatric neurosurgery: why, when, how? *Childs Nerv Syst* 2002;18:264-287.
44. Nguyen JP, Lefaucheur JP, Decq P, et al. Chronic motor cortex stimulation in the treatment of central and neuropathic pain. Correlations between clinical, electrophysiological and anatomical data. *Pain* 1999;82:245-251.
45. Quinones-Hinojosa A, Alam M, Lyon R, et al. Transcranial motor evoked potentials during basilar artery aneurysm surgery: technique application for 30 consecutive patients. *Neurosurgery* 2004;54:916-924; discussion 924.
46. McElveen JT Jr, Belmonte RG, Fukushima T, Bullard DE. A review of facial nerve outcome in 100 consecutive cases of acoustic tumor surgery. *Laryngoscope* 2000;110:1667-1672.
47. Belzberg AJ, Dorsi MJ, Storm PB, Moriarity JL. Surgical repair of brachial plexus injury: a multinational survey of experienced peripheral nerve surgeons. *J Neurosurg* 2004;101:365-376.
48. Xu JG, Gu YD, Hu SN, et al. Electrophysiologic study of influence on C7 innervated muscles after transection of different C7 fascicles. *Microsurgery* 2004;24:147-150.
49. Balakrishnan G, Kadadi BK. Clinical examination versus routine and paraspinale electromyographic studies in predicting the site of lesion in brachial plexus injury. *J Hand Surg [Am]* 2004;29:140-143.
50. Pring ME, Trousdale RT, Cabanela ME, Harper CM. Intraoperative electromyographic monitoring during periacetabular osteotomy. *Clin Orthop Relat Res* 2002; (400):158-164.
51. Toleikis JR, Skelly JP, Calvin AO, Burkus JK. Spinally elicited peripheral nerve responses are sensory rather than motor. *Clin Neurophysiol* 2000;111:736-742.
52. Rose RD. Removing the antidromically driven sensory component from cervically evoked motor potentials. *Med Hypotheses* 1998;50:147-154.
53. Scheufler KM, Zentner J. Total intravenous anesthesia for intraoperative monitoring of the motor pathways: an integral view combining clinical and experimental data. *J Neurosurg* 2002;96:571-579.
54. Wang AC, Than KD, Etame AB, et al. Impact of anesthesia on transcranial electric motor evoked potential monitoring during spine surgery: a review of the literature. *Neurosurg Focus* 2009;27:E7.
55. Zentner J, Albrecht T, Heuser D. Influence of halothane, enflurane, and isoflurane on motor evoked potentials. *Neurosurgery* 1992;31:298-305.
56. Nathan N, Tabaraud F, Lacroix F, et al. Influence of propofol concentrations on multipulse transcranial motor evoked potentials. *Br J Anaesth* 2003;91:493-497.
57. Sakamoto T, Kawaguchi M, Inoue S, Furuya H. Suppressing effect of nitrous oxide on motor evoked potentials can be reversed by train stimulation in rabbits under ketamine/fentanyl anaesthesia, but not with additional propofol. *Br J Anaesth* 2001;86:395-402.
58. Hargreaves SJ, Watt JW. Intravenous anaesthesia and repetitive transcranial magnetic stimulation monitoring in spinal column surgery. *Br J Anaesth* 2005;94:70-73.
59. Costa P, Bruno A, Bonzanino M, Massaro F, et al. Somatosensory- and motor-evoked potential monitoring during spine and spinal cord surgery. *Spinal Cord* 2007;45(1):86-91.
60. Lo YL, Dan YF, Tan YE, et al. Intraoperative motor-evoked potential monitoring in scoliosis surgery: comparison of desflurane/nitrous oxide with propofol total intravenous anesthetic regimens. *J Neurosurg Anesthesiol* 2006;18:211-214.
61. Tanaka S, Kobayashi I, Sagiuchi T, et al. Compensation of intraoperative transcranial motor-evoked potential monitoring by compound muscle action potential after peripheral nerve stimulation. *J Clin Neurophysiol* 2005;22:271-274.
62. Sala F, Lanteri P, Bricolo A. Motor evoked potential monitoring for spinal cord and brain stem surgery. *Adv Tech Stand Neurosurg* 2004;29:133-169.
63. Vauzelle C, Stagnara P, Jouvinroux P. Functional monitoring of spinal cord activity during spinal surgery. *Clin Orthop Relat Res* 1973;(93):173-178.
64. Owen JH. The application of intraoperative monitoring during surgery for spinal deformity. *Spine* 1999;24:2649-2662.

65. Keifer J. Anesthesia for craniotomy with intraoperative speech testing. *Curr Rev Clin Anesthesi* 2001;21:283-290.
66. Keifer JC, Dentchev D, Little K, et al. A retrospective analysis of a remifentanyl/propofol general anesthetic for craniotomy before awake functional brain mapping. *Anesth Analg* 2005;101:502-508, table of contents.
67. Lind G, Meyerson BA, Winter J, Linderoth B. Implantation of laminotomy electrodes for spinal cord stimulation in spinal anesthesia with intraoperative dorsal column activation. *Neurosurgery* 2003;53:1150-1153; discussion 1153-1154.