

Chapter 35

REFERENCES

1. Geissler PL, Dellago C, Chandler D, et al. Autoionization in liquid water. *Science*. 2001;291:2121-2124.
2. Haskins SC, Hopper K, Rezende ML. The acid-base impact of free water removal from, and addition to, plasma. *J Lab Clin Med*. 2006;147:114-120.
3. Scheingraber S, Rehm M, Sehmisch C, Finsterer U. Rapid saline infusion produces hyperchloremic acidosis in patients undergoing gynecologic surgery. *Anesthesiology*. 1999;90:1265-1270.
4. Fencel V, Leith DE. Stewart's quantitative acid-base chemistry: applications in biology and medicine. *Respir Physiol*. 1993;91:1-16.
5. Cuthbertson D, Tilstone WJ. Metabolism during the postinjury period. *Adv Clin Chem*. 1969;12:1-55.
6. Moore FD. Some observations on the metabolic requirement in surgical patients. *Bull N Engl Med Cent*. 1949;11:193-201.
7. Moore FD. Common patterns of water and electrolyte change in injury, surgery and disease. *N Engl J Med*. 1958;258:277-285.
8. Drucker WR, Chadwick CD, Gann DS. Transcapillary refill in hemorrhage and shock. *Arch Surg*. 1981;116:1344-1153.
9. Marty AT, Zweifach BW. Splanchnic contribution to transcapillary refill after hemorrhagic shock. *Ann Surg*. 1971;174:131-136.
10. Moss G. Fluid distribution in prevention of hypovolemic shock. *Arch Surg*. 1969;98:281-286.
11. Holte K, Kehlet H. Compensatory fluid administration for preoperative dehydration—does it improve outcome? *Acta Anaesthesiol Scand*. 2002;46:1089-1093.
12. Maharaj CH, Kallam SR, Malik A, et al. preoperative intravenous fluid therapy decreases postoperative nausea and pain in high risk patients. *Anesth Analg*. 2005;100:675-682.
13. Magner JJ, McCaul C, Carton E, et al. Effect of intraoperative intravenous crystalloid infusion on postoperative nausea and vomiting after gynaecological laparoscopy: comparison of 30 and 10 ml kg(-1). *Br J Anaesth*. 2004;93:381-385.
14. McCaul C, Moran C, O'Cronin D, et al. Intravenous fluid loading with or without supplementary dextrose does not prevent nausea, vomiting and pain after laparoscopy. *Can J Anaesth*. 2003;50:440-444.
15. Brandstrup B. Fluid therapy for the surgical patient. *Best Pract Res Clin Anaesthesiol*. 2006;20:265-283.
16. Holte K, Sharrock NE, Kehlet H. Pathophysiology and clinical implications of perioperative fluid excess. *Br J Anaesth*. 2002;89:622-632.
17. Shires T, Williams J, Brown F. Acute change in extracellular fluids associated with major surgical procedures. *Ann Surg*. 1961;154:803-810.
18. Brandstrup B, Svendsen C, Engquist A. Hemorrhage and operation cause a contraction of the extracellular space needing replacement—evidence and implications? A systematic review. *Surgery*. 2006;139:419-432.
19. Brandstrup B, Tonnesen H, Beier-Holgersen R, et al. Effects of intravenous fluid restriction on postoperative complications: comparison of two perioperative fluid regimens: a randomized assessor-blinded multicenter trial. *Ann Surg*. 2003;238:641-648.
20. Holte K, Jensen P, Kehlet H. Physiologic effects of intravenous fluid administration in healthy volunteers. *Anesth Analg*. 2003;96:1504-1509.
21. Adrogué HJ, Madias NE. Hyponatremia. *N Engl J Med*. 2000;342:1581-1589.
22. Adrogué HJ, Madias NE. Hypernatremia. *N Engl J Med*. 2000;342:1493-1499.
23. Fawcett WJ, Haxby EJ, Male DA. Magnesium: physiology and pharmacology. *Br J Anaesth*. 1999;83:302-320.
24. Gupta K, Vohra V, Sood J. The role of magnesium as an adjuvant during general anaesthesia. *Anaesthesia*. 2006;61:1058-1063.
25. Blitz M, Blitz S, Beasley R, et al. Inhaled magnesium sulfate in the treatment of acute asthma. *Cochrane Database Syst Rev*. 2005;CD003898.
26. Dube L, Granry JC. The therapeutic use of magnesium in anesthesiology, intensive care and emergency medicine: a review: [L'usage thérapeutique du magnésium en anesthésiologie, réanimation et médecine d'urgence]. *Can J Anesth*. 2003;50:732-746.
27. Agustí AG, Torres A, Estopa R, Agustívidal A. Hypophosphatemia as a cause of failed weaning: the importance of metabolic factors. *Crit Care Med*. 1984;12:142-143.
28. Aubier M, Murciano D, Lecocguic Y, et al. Effect of hypophosphatemia on diaphragmatic contractility in patients with acute respiratory failure. *N Engl J Med*. 1985;313:420-424.
29. Zazzo JF, Troche G, Ruel P, Maintenant J. High incidence of hypophosphatemia in surgical intensive care patients: efficacy of phosphorus therapy on myocardial function. *Intensive Care Med*. 1995;21:826-831.
30. Nilius B, Droogmans G. Amazing chloride channels: an overview. *Acta Physiol Scand*. 2003;177:119-147.

31. Jentsch TJ. Chloride channels are different. *Nature*. 2002;415:276-277.
32. Thakker RV. Chloride channels in renal disease. *Adv Nephrol Necker Hosp*. 1999;29:289-298.
33. Wilcox CS. Regulation of renal blood flow by plasma chloride. *J Clin Invest* 1983;71:726-735.
34. Wilkes NJ, Woolf R, Mutch M, et al. The effects of balanced versus saline-based hetastarch and crystalloid solutions on acid-base and electrolyte status and gastric mucosal perfusion in elderly surgical patients. *Anesth Analg*. 2001;93:811-816.
35. Nicholson JP, Wolmarans MR, Park GR. The role of albumin in critical illness. *Br J Anaesth*. 2000;85:599-610.
36. Soni N. Human albumin administration in critically ill patients. Validity of review methods must be assessed. *BMJ* 1998;317:883-884.
37. Perioperative total parenteral nutrition in surgical patients. The Veterans Affairs Total Parenteral Nutrition Cooperative Study Group. *N Engl J Med*. 1991;325:525-532.
38. Goldwasser P, Feldman J. Association of serum albumin and mortality risk. *J Clin Epidemiol*. 1997;50:693-703.
39. Blunt MC, Nicholson JP, Park GR. Serum albumin and colloid osmotic pressure in survivors and nonsurvivors of prolonged critical illness. *Anaesthesia*. 1998;53:755-761.
40. Narins R, Emmett M. Simple and mixed acid-base disorders: a practical approach. *Medicine (Baltimore)*. 1980;59:161-187.
41. Stewart PA. Independent and dependent variables of acid-base control. *Respir Physiol*. 1978;33:9-26.
42. Fencel V, Jabor A, Kazda A, Figge J. Diagnosis of metabolic acid-base disturbances in critically ill patients. *Am J Respir Crit Care Med*. 2000;162:2246-2251.
43. Figge J, Rossing TH, Fencel V. The role of serum proteins in acid-base equilibria. *J Lab Clin Med*. 1991;117:453-467.
44. Wilkes P. Hypoproteinemia, strong-ion difference, and acid-base status in critically ill patients. *J Appl Physiol*. 1998;84:1740-1748.
45. Stewart PA. Modern quantitative acid-base chemistry. *Can J Physiol Pharmacol*. 1983;61:1444-1461.
46. Kellum JA. In: Shoemaker W, ed. *Diagnosis and Treatment of Acid Base Disorders, Textbook of Critical Care Medicine*. 4th ed. Philadelphia, PA: Saunders; 2000:839-853.
47. Geissler PL, Dellago C, Chandler D, et al. Autoionization in liquid water. *Science*. 2001;291:2121-2124.
48. Chaplin MF. A proposal for the structuring of water. *Biophys Chem*. 2000;83:211-221.
49. Rini M, Magnes BZ, Pines E, Nibbering ETJ. Real-time observation of bimodal proton transfer in acid-base pairs in water. *Science*. 2003;301:349-352.
50. Marx D, Tuckerman ME, Hutter J, Parrinello M. The nature of the hydrated excess proton in water. *Nature*. 1999;397:601-604.
51. Chaplin MF. A proposal for the structuring of water. *Biophys Chem*. 2000;83:211-221.
52. Singer RB, Hastings AB. An improved clinical method for the estimation of disturbances of the acid-base balance of human blood. *Medicine*. 1948;10:242.
53. Rossing TH, Maffeo N, Fencel V. Acid-base effects of altering plasma protein concentration in human blood in vitro. *J Appl Physiol*. 1986;61:2260-2265.
54. Corey HE. Stewart and beyond: new models of acid-base balance. *Kidney Int*. 2003;64:777-787.
55. Goldwasser P, Feldman J. Association of serum albumin and mortality risk. *J Clin Epidemiol*. 1997;50:693-703.
56. Alfaro V, Torras R, Ibanez J, Palacios L. A physical-chemical analysis of the acid-base response to chronic obstructive pulmonary disease. *Can J Physiol Pharmacol*. 1996;11(74):1229-1235.
57. Rehm MO, Orth V, Scheingraber S, et al. Acid-base changes caused by 5% albumin versus 6% HES in patients undergoing acute normovolemic hemodilution: a randomized prospective study. *Anesthesiology*. 2000;93:1174-1183.
58. Waters J, Gottlieb A, Schoenwald P, Popovich M. Normal saline versus lactated Ringer's solution for intraoperative fluid management in patients undergoing abdominal aortic aneurysm repair: an outcome study. *Anesth Analg*. 2001;93:817-822.
59. Kellum JA. Fluid resuscitation and hyperchloremic acidosis in experimental sepsis: improved short-term survival and acid-base balance with Hextend compared with saline. *Crit Care Med*. 2002;30:300-305.
60. Brill SA, Stewart TR, Brundage SI, Schreiber MA. Base deficit does not predict mortality when secondary to hyperchloremic acidosis. *Shock*. 2002;17:459-462.
61. Tournadre JP, Allaouchiche B, Malbert CH, Chassard D. Metabolic acidosis and respiratory acidosis impair gastropyloric motility in anesthetized pigs. *Anesth Analg*. 2000;90:74-79.
62. Wilcox CS. Regulation of renal blood flow by plasma chloride. *J Clin Invest*. 1983;71:726-735.
63. Hansen PB, Jensen BL, Skott O. Chloride regulates afferent arteriolar contraction in response to depolarization. *Hypertension*. 1998;32:1066-1070.
64. Williams EL, Hildebrand KL, McCormick SA, Bedel MJ. The effect of intravenous lactated Ringer's solution versus 0.9% sodium chloride solution on serum osmolality in human volunteers. *Anesth Analg*. 1999;88:999-1003.
65. Merten GJ, Burgess WP, Gray LV, et al. Prevention of contrast-induced nephropathy with sodium bicarbonate: a randomized controlled trial. *JAMA*. 2004;291:2328-2334.
66. Haase M, Haase-Fielitz A, Bellomo R, et al. Sodium bicarbonate to prevent increases in serum creatinine after cardiac surgery: a pilot double-blind, randomized controlled trial. *Crit Care Med*. 2009;37:39-47.
67. Story DA, Poustie S, Bellomo R. Quantitative physical chemistry analysis of acid-base disorders in critically ill patients. *Anaesthesia*. 2001;56:530-533.
68. Tayar J, Jabbour G, Saggi SJ. Severe hyperosmolar metabolic acidosis due to a large dose of intravenous lorazepam. *N Engl J Med*. 2002;346:1253-1254.
69. Rocktäschel J, Morimatsu H, Uchino S, et al. Impact of continuous veno-venous hemofiltration on acid-base balance. *Int J Artif Organs*. 2003;26:19-25.
70. Bruegger D, Bauer A, Rehm M, et al. Effect of hypertonic saline dextran on acid-base balance in patients undergoing surgery of abdominal aortic aneurysm. *Crit Care Med*. 2005;33:556-563.
71. Kellum JA. In: Shoemaker W, ed. *Diagnosis and Treatment of Acid Base Disorders, Textbook of Critical Care Medicine*. 4th ed. Philadelphia, PA: Saunders; 2000:839-853.
72. Rodriguez-Soriano J. New insights into the pathogenesis of renal tubular acidosis—from functional to molecular studies. *Pediatr Nephrol*. 2000;14:1121-1136.

73. Choate KA, Kahle KT, Wilson FH, et al. WNK1, a kinase mutated in inherited hypertension with hyperkalemia, localizes to diverse Cl⁻ transporting epithelia. *Proc Natl Acad Sci U S A*. 2003;100:663-668.
74. Shaer AJ. Inherited primary renal tubular hypokalemic alkalosis: a review of Gitelman and Bartter syndromes. *Am J Med Sci*. 2001;322:316-332.
75. Severinghaus JW. Acid-base balance nomogram—a Boston-Copenhagen detente. *Anesthesiology*. 1976;45:539-541.
76. Schwartz WB, Relman AS. A critique of the parameters used in the evaluation of acid-base disorders. “Whole-blood buffer base” and “standard bicarbonate” compared with blood pH and plasma bicarbonate concentration. *N Engl J Med*. 1963;268:1382-1388.
77. Siggaard-Anderson O. *The Acid Base Status of the Blood*. Copenhagen Denmark: Munksgard; 1963:134.
78. Siggaard-Andersen O. The van Slyke equation. *Scand J Clin Lab Invest Suppl*. 1977;37:15-20.
79. Schlichtig R, Grogono AW, Severinghaus JW. Human PaCO₂ and standard base excess compensation for acid-base imbalance. *Crit Care Med*. 1998;26:1173-1179.
80. Morgan TJ, Clark C, Endre ZH. Accuracy of base excess—an in vitro evaluation of the Van Slyke equation. *Crit Care Med*. 2003;28:2932-2936.
81. Emmett M, Narins RG. Clinical use of the anion gap. *Medicine (Baltimore)*. 1977;56:38-54.
82. Salem MM, Mujais SK. Gaps in the anion gap. *Arch Intern Med*. 1992;152:1625-1629.
83. Figge J, Jabor A, Kazda A, Fencl V. Anion gap and hypoalbuminemia. *Crit Care Med*. 1998;26:1807-1810.
84. Wooten EW. Analytic calculation of physiological acid-base parameters in plasma. *J Appl Physiol*. 1999;86:326-334.
85. Figge J, Mydosh T, Fencl V. Serum proteins and acid-base equilibria: a follow-up. *J Lab Clin Med*. 1992;120:713-719.
86. Vitek V, Cowley RA. Blood lactate in the prognosis of various forms of shock. *Ann Surg*. 1971;173:308-313.
87. Eddy VA, Morris JA Jr, Cullinane DC. Hypothermia, coagulopathy, and acidosis. *Surg Clin North Am*. 2000;80:845-854.
88. Husain FA, Martin MJ, Mullenix PS, et al. Serum lactate and base deficit as predictors of mortality and morbidity. *Am J Surg*. 2003;185:485-491.
89. Abramson D, Scalea TM, Hitchcock R, et al. Lactate clearance and survival following injury. *J Trauma*. 1993;35:584-588.
90. McNelis J, Marini CP, Jurkiewicz A, et al. Prolonged lactate clearance is associated with increased mortality in the surgical intensive care unit. *Am J Surg*. 2001;182:481-485.
91. Nguyen HB, Rivers EP, Knoblich BP, et al. Early lactate clearance is associated with improved outcome in severe sepsis and septic shock. *Crit Care Med*. 2004;32:1637-1642.
92. Baranov D, Neligan P. Trauma and aggressive homeostasis management. *Anesthesiol Clin*. 2007;25:49-63, viii.
93. Brill SA, Stewart TR, Brundage SI, Schreiber MA. Base deficit does not predict mortality when secondary to hyperchloremic acidosis. *Shock*. 2002;17:459-462.
94. Kaplan LJ, Kellum JA. Initial pH, base deficit, lactate, anion gap, strong ion difference, and strong ion gap predict outcome from major vascular injury. *Crit Care Med*. 2004;32:1120-1124.
95. Martin MJ, FitzSullivan E, Salim A, et al. Discordance between lactate and base deficit in the surgical intensive care unit: which one do you trust? *Am J Surg*. 2006;191:625-630.
96. Kaplan LJ, Kellum JA. Comparison of acid-base models for prediction of hospital mortality after trauma. *Shock*. 2008;29:662-666.
97. Zehtabchi S, Soghoian S, Sinert R. Utility of Stewart’s strong ion difference as a predictor of major injury after trauma in the ED. *Am J Emerg Med*. 2007;25:938-941.
98. Omron EM. Comparative quantitative acid-base analysis in coronary artery bypass, severe sepsis, and diabetic ketoacidosis. *J Intensive Care Med*. 2005;20:317-326.
99. Martin M, Murray J, Berne T, et al. Diagnosis of acid-base derangements and mortality prediction in the trauma intensive care unit: the physicochemical approach. *J Trauma*. 2005;58:238-243.
100. Rocktaeschel J, Morimatsu H, Uchino S, Bellomo R. Unmeasured anions in critically ill patients: can they predict mortality? *Crit Care Med*. 2003;31:2131-2136.
101. Cusack RJ, Rhodes A, Lochhead P, et al. The strong ion gap does not have prognostic value in critically ill patients in a mixed medical/surgical adult ICU. *Intensive Care Med*. 2002;28:864-869.
102. Moviat M, van HF, van der HH. Conventional or physicochemical approach in intensive care unit patients with metabolic acidosis. *Crit Care*. 2003;7:R41-R45.
103. Esche V, Ru M, Melzer S, Gromann B, et al. Buffer capacity of 4% succinylated gelatin does not provide any advantages over acidic 6% hydroxyethyl starch 130/0.4 for acid-base balance during experimental mixed acidemia in a porcine model. *Eur J Anaesthesiol*. 2008;25:905-916.
104. Gilfix BM, Bique M, Magder S. A physical chemical approach to the analysis of acid-base balance in the clinical setting. *J Crit Care*. 1993;8:187-197.
105. Balasubramanian N, Havens PL, Hoffman GM. Unmeasured anions identified by the Fencl-Stewart method predict mortality better than base excess, anion gap, and lactate in patients in the pediatric intensive care unit. *Crit Care Med*. 1999;27:1577-1581.
106. Story DA, Morimatsu H, Bellomo R. Strong ions, weak acids and base excess: a simplified Fencl-Stewart approach to clinical acid-base disorders. *Br J Anaesth*. 2004;92:54-60.
107. Forsythe SM, Schmidt GA. Sodium bicarbonate for the treatment of lactic acidosis. *Chest*. 2000;117:260-267.
108. Rehm M, Finsterer U. Treating intraoperative hyperchloremic acidosis with sodium bicarbonate or tris-hydroxymethyl aminomethane: a randomized prospective study. *Anesth Analg*. 2003;96:1201-1208, table.
109. Goldsmith DJ, Forni LG, Hilton PJ. Bicarbonate therapy and intracellular acidosis. *Clin Sci (Lond)*. 1997;93:593-598.
110. Nielsen HB, Hein L, Svendsen LB, et al. Bicarbonate attenuates intracellular acidosis. *Acta Anaesthesiol Scand*. 2002;46:579-584.
111. Hickling KG. Permissive hypercapnia. *Respir Care Clin N Am*. 2002;8:155-169, v.
112. Costello J, Higgins B, Contreras M, et al. Hypercapnic acidosis attenuates shock and lung injury in early and prolonged systemic sepsis. *Crit Care Med*. 2009;37:2412-2420.
113. Curley G, Laffey J, Kavanagh B. Bench-to-bedside review: carbon dioxide. *Crit Care*. 2010;14:220.
114. Laffey JG, Engelberts DORE, Kavanagh BP. Buffering hypercapnic acidosis worsens Acute lung injury. *Am J Respir Crit Care Med*. 2000;161:141-146.
115. Holmdahl MH, Wiklund L, Wetterberg T, et al. The place of THAM in the management of acidemia in clinical practice. *Acta Anaesthesiol Scand*. 2000;44:524-527.

116. Reeves RB. An imidazole alaphastat hypothesis for vertebrate acid-base regulation: tissue carbon dioxide content and body temperature in bullfrogs. *Respir Physiol.* 1972;14:219-236.
117. Rahn H, Reeves RB, Howell BJ. Hydrogen ion regulation, temperature, and evolution. *Am Rev Respir Dis.* 1975;112:165-172.
118. Laffey JG, Kavanagh BP. Carbon dioxide and the critically ill—too little of a good thing? *Lancet.* 1999;354:1283-1286.
119. Murkin JM, Farrar JK, Tweed WA, et al. Cerebral autoregulation and flow/metabolism coupling during cardiopulmonary bypass: the influence of PaCO₂. *Anesth Analg.* 1987;66:825-832.
120. Hogue CW Jr, Palin CA, Arrowsmith JE. Cardiopulmonary bypass management and neurologic outcomes: an evidence-based appraisal of current practices. *Anesth Analg.* 2006;103:21-37.
121. Practice guidelines for preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration: application to healthy patients undergoing elective procedures: a report by the American Society of Anesthesiologist Task Force on Preoperative Fasting. *Anesthesiology.* 1999;90:896-905.
122. Dillon J, Lynch LJ Jr, Myers R, Butcher HR Jr. The treatment of hemorrhagic shock. *Surg Gynecol Obstet.* 1966;122:967-978.
123. Skillman JJ, Awwad HK, Moore FD. Plasma protein kinetics of the early transcapillary refill after hemorrhage in man. *Surg Gynecol Obstet.* 1967;125:983-996.
124. Fleck A, Raines G, Hawker F, et al. Increased vascular permeability: a major cause of hypoalbuminaemia in disease and injury. *Lancet.* 1985;1:781-784.
125. Fisher M. Postoperative intravenous therapy. *Anaesth Intensive Care.* 1977;5:339-346.
126. Wan L, Bellomo R, May CN. The effects of normal and hypertonic saline on regional Blood flow and oxygen delivery. *Anesth Analg.* 2007;105:141-147.
127. Van den Berghe G, Wouters P, Weekers F, et al. Intensive Insulin therapy in critically ill patients. *N Engl J Med.* 2001;345:1359-1367.
128. Capes SE, Hunt D, Malmberg K, Gerstein HC. Stress hyperglycaemia and increased risk of death after myocardial infarction in patients with and without diabetes: a systematic overview. *Lancet.* 2000;355:773-778.
129. Bochicchio GV, Sung J, Joshi M, et al. Persistent hyperglycemia is predictive of outcome in critically ill trauma patients. *J Trauma.* 2005;58:921-924.
130. O'Malley CM, Frumento RJ, Hardy MA, et al. A randomized, double-blind comparison of lactated Ringer's solution and 0.9% NaCl during renal transplantation. *Anesth Analg.* 2005;100:1518-1524, table.
131. Hadimioglu N, Saadawy I, Saglam T, et al. The effect of different crystalloid solutions on acid-base balance and early kidney function after kidney transplantation. *Anesth Analg.* 2008;107:264-269.
132. Hansen PB, Jensen BL, Skott O. Chloride regulates afferent arteriolar contraction in response to depolarization. *Hypertension.* 1998;32:1066-1070.
133. Wilcox CS. Regulation of renal blood flow by plasma chloride. *J Clin Invest.* 1983;71:726-735.
134. Lobo DN, Bostock KA, Neal KR, et al. Effect of salt and water balance on recovery of gastrointestinal function after elective colonic resection: a randomised controlled trial. *Lancet.* 2002;359:1812-1818.
135. Merten GJ, Burgess WP, Gray LV, et al. Prevention of contrast-induced nephropathy with sodium bicarbonate: a randomized controlled trial. *JAMA.* 2004;291:2328-2334.
136. Boldt J. New light on intravascular volume replacement regimens: what did we learn from the past three years? *Anesth Analg.* 2003;97:1595-1604.
137. Ruttman TG, James MFM, Finlayson J. Effects on coagulation of intravenous crystalloid or colloid in patients undergoing peripheral vascular surgery. *Br J Anaesth.* 2002;89:226-230.
138. Ng KF, Lam CC, Chan LC. In vivo effect of haemodilution with saline on coagulation: a randomized controlled trial. *Br J Anaesth.* 2002;88:475-480.
139. Roche AM, James MFM, Nett-Guerrero E, Mythen MG. A head-to-head comparison of the in vitro coagulation effects of saline-based and balanced electrolyte crystalloid and colloid intravenous fluids. *Anesth Analg.* 2006;102:1274-1279.
140. Ruttman TG, Jamest MF, Lombard EH. Haemodilution-induced enhancement of coagulation is attenuated in vitro by restoring antithrombin III to pre-dilution concentrations. *Anaesth Intensive Care.* 2001;29:489-493.
141. Healey MA, Davis RE, Liu FC, et al. Lactated Ringer's is superior to normal saline in a model of massive hemorrhage and resuscitation. *J Trauma.* 1998;45:894-899.
142. Waters JH, Gottlieb A, Schoenwald P, et al. Normal saline versus lactated Ringer's solution for intraoperative fluid management in patients undergoing abdominal aortic aneurysm repair: an outcome study. *Anesth Analg.* 2001;93:817-822.
143. Tataru T, Nagao Y, Tashiro C. The effect of duration of surgery on fluid balance during abdominal surgery: a mathematical model. *Anesth Analg.* 2009;109:211-216.
144. Mazzoni MC, Borgstrom P, Intaglietta M, Arfors KE. Capillary narrowing in hemorrhagic shock is rectified by hyperosmotic saline-dextran reinfusion. *Circ Shock.* 1990;31:407-418.
145. Bruegger D, Bauer A, Rehm M, et al. Effect of hypertonic saline dextran on acid-base balance in patients undergoing surgery of abdominal aortic aneurysm. *Crit Care Med.* 2005;33:556-563.
146. Oliveira RP, Weingartner R, Ribas EO, et al. Acute haemodynamic effects of a hypertonic saline/dextran solution in stable patients with severe sepsis. *Intensive Care Med.* 2002;28:1574-1581.
147. Nolan J. Fluid resuscitation for the trauma patient. *Resuscitation.* 2001;48:57-69.
148. Mattox KL, Maningas PA, Moore EE, et al. Prehospital hypertonic saline/dextran infusion for post-traumatic hypotension. The U.S.A. Multicenter Trial. *Ann Surg.* 1991;213:482-491.
149. Wade CE, Grady JJ, Kramer GC. Efficacy of hypertonic saline dextran fluid resuscitation for patients with hypotension from penetrating trauma. *J Trauma.* 2003;54(Suppl):S144-S148.
150. Wade CE, Kramer GC, Grady JJ, et al. Efficacy of hypertonic 7.5% saline and 6% dextran-70 in treating trauma: a meta-analysis of controlled clinical studies. *Surgery.* 1997;122:609-616.
151. Cooper DJ, Myles PS, McDermott FT, et al. Prehospital hypertonic saline resuscitation of patients with hypotension and severe traumatic brain injury: a randomized controlled trial. *JAMA.* 2004;291:1350-1357.
152. Jarvela K, Kaukinen S. Hypertonic saline (7.5%) decreases perioperative weight gain following cardiac surgery. *J Cardiothorac Vasc Anesth.* 2002;16:43-46.
153. White H, Cook D, Venkatesh B. The use of hypertonic saline for treating intracranial hypertension after traumatic brain injury. *Anesth Analg.* 2006;102:1836-1846.

154. Klatzo I. Presidential address. Neuropathological aspects of brain edema. *J Neuropathol Exp Neurol*. 1967;26:1-14.
155. Schmoker JD, Shackford SR, Wald SL, Pietropaoli JA. An analysis of the relationship between fluid and sodium administration and intracranial pressure after head injury. *J Trauma*. 1992;33:476-481.
156. Grocott MPW, Mythen MG, Gan TJ. Perioperative fluid management and clinical outcomes in adults. *Anesth Analg*. 2005;100:1093-1106.
157. Blunt MC, Nicholson JP, Park GR. Serum albumin and colloid osmotic pressure in survivors and nonsurvivors of prolonged critical illness. *Anaesthesia*. 1998;53:755-761.
158. Jacob M, Bruegger D, Rehm M, et al. Contrasting effects of colloid and crystalloid resuscitation fluids on cardiac vascular permeability. *Anesthesiology*. 2006;104: 1223-1231.
159. Rittoo D, Gosling P, Burnley S, et al. Randomized study comparing the effects of hydroxyethyl starch solution with Gelofusine on pulmonary function in patients undergoing abdominal aortic aneurysm surgery. *Br J Anaesth*. 2004;92:61-66.
160. Zikria BA, King TC, Stanford J, Freeman HP. A biophysical approach to capillary permeability. *Surgery*. 1989;105:625-631.
161. Choi PT, Yip G, Quinonez LG, Cook DJ. Crystalloids vs. colloids in fluid resuscitation: a systematic review. *Crit Care Med*. 1999;27:200-210.
162. Roberts I, Alderson P, Bunn F, et al. Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database Syst Rev*. 2004;CD000567.
163. Velanovich V. Crystalloid versus colloid fluid resuscitation: a meta-analysis of mortality. *Surgery*. 1989;105:65-71.
164. Dubois MJ, Orellana-Jimenez C, Melot C, et al. Albumin administration improves organ function in critically ill hypoalbuminemic patients: a prospective, randomized, controlled, pilot study. *Crit Care Med*. 2006;34:2536-2540.
165. Human albumin administration in critically ill patients: systematic review of randomised controlled trials. Cochrane Injuries Group Albumin Reviewers. *BMJ*. 1998;317:235-240.
166. The SAFE Investigators. A comparison of albumin and saline for fluid resuscitation in the intensive care unit. *N Engl J Med*. 2004;350:2247-2256.
167. Boldt J. Use of albumin: an update. *Br J Anaesth*. 2010;104: 276-284.
168. Boldt J, Schöhlhorn T, Mayer J, et al. The value of an albumin-based intravascular volume replacement strategy in elderly patients undergoing major abdominal surgery. *Anesth Analg*. 2006;103:191-199.
169. Zinderman CE, Landow L, Wise RP. Anaphylactoid reactions to dextran 40 and 70: reports to the United States Food and Drug Administration, 1969 to 2004. *J Vasc Surg*. 2006;43:1004-1009.
170. Ljungstrom KG, Renck H, Strandberg K, et al. Adverse reactions to dextran in Sweden 1970-1979. *Acta Chirurgica Scand*. 1983;149:253-262.
171. Yacobi A, Stoll RG, Sum CY, et al. Pharmacokinetics of hydroxyethyl starch in normal subjects. *J Clin Pharmacol*. 1982;22:206-212.
172. Feng X, Liu J, Yu M, et al. Hydroxyethyl starch, but not modified fluid gelatin, affects inflammatory response in a rat model of polymicrobial sepsis with capillary leakage. *Anesth Analg*. 2007;104:624-630.
173. Treib J, Haass A, Pindur G, et al. Influence of low molecular weight hydroxyethyl starch (HES 40/0.5-0.55) on hemostasis and hemorheology. *Haemostasis*. 1996;26:258-265.
174. Gan TJ, Bennett-Guerrero E, Phillips-Bute B, et al. Hextend(R), a physiologically balanced plasma expander for large volume use in major surgery: a randomized phase III clinical trial. *Anesth Analg*. 1999;88:992-998.
175. Westphal M, James MFM, Kozek-Langenecker S, et al. Hydroxyethyl starches: different products—different effects. *Anesthesiology*. 2009;111:187-202.
176. Zaar M, Lauritzen B, Secher NH, et al. Initial administration of hydroxyethyl starch vs lactated Ringer after liver trauma in the pig. *Br J Anaesth*. 2009;102:221-226.
177. Bedforth NM, Hardman JG. Mad colloid disease? *Anaesthesia*. 1997;52:389-390.
178. Chappell D, Jacob M, Hofmann-Kiefer K, et al. A rational approach to perioperative fluid management. *Anesthesiology*. 2008;109:723-740.
179. Funk W, Baldinger V. Microcirculatory perfusion during volume therapy. A comparative study using crystalloid or colloid in awake animals. *Anesthesiology*. 1995;82:975-982.
180. Lang K, Boldt J, Suttner S, Haisch G. Colloids versus crystalloids and tissue oxygen tension in patients undergoing major abdominal surgery. *Anesth Analg*. 2001;93:405-409.
181. Ernest D, Belzberg AS, Dodek PM. Distribution of normal saline and 5% albumin infusions in septic patients. *Crit Care Med*. 1999;27:46-50.
182. Tataru T, Tsuneto T, Tashiro C. Crystalloid infusion rate during fluid resuscitation from acute haemorrhage. *Br J Anaesth*. 2007;99:212-217.
183. Rhee P, Burris D, Kaufmann C, et al. Lactated Ringer's solution resuscitation causes neutrophil activation after hemorrhagic shock. *J Trauma*. 1998;44:313-319.
184. Alam HB, Stanton K, Koustova E, et al. Effect of different resuscitation strategies on neutrophil activation in a swine model of hemorrhagic shock. *Resuscitation*. 2004;60:91-99.
185. Deb S, Martin B, Sun L, et al. Resuscitation with lactated Ringer's solution in rats with hemorrhagic shock induces immediate apoptosis. *J Trauma*. 1999;46:582-588.
186. Deb S, Martin B, Sun L, et al. Resuscitation with lactated Ringer's solution in rats with hemorrhagic shock induces immediate apoptosis. *J Trauma*. 1999;46:582-588.
187. Alam HB, Stanton K, Koustova E, et al. Effect of different resuscitation strategies on neutrophil activation in a swine model of hemorrhagic shock. *Resuscitation*. 2004;60:91-99.
188. Shoemaker WC, Montgomery ES, Kaplan E, Elwyn DH. Physiologic patterns in surviving and nonsurviving shock patients. Use of sequential cardiorespiratory variables in defining criteria for therapeutic goals and early warning of death. *Arch Surg*. 1973;106:630-636.
189. Giglio MT, Marucci M, Testini M, Brienza N. Goal-directed haemodynamic therapy and gastrointestinal complications in major surgery: a meta-analysis of randomized controlled trials. *Br J Anaesth*. 2009;103:637-646.
190. Moretti EW, Robertson KM, el Moalem H, Gan TJ. Intraoperative colloid administration reduces postoperative nausea and vomiting and improves postoperative outcomes compared with crystalloid administration. *Anesth Analg*. 2003;96:611-617, table.
191. Mythen MG, Webb AR. Perioperative plasma volume expansion reduces the incidence of gut mucosal hypoperfusion during cardiac surgery. *Arch Surg*. 1995;130:423-429.

192. Gan TJ, Soppitt A, Maroof M, et al. Goal-directed intraoperative fluid administration reduces length of hospital stay after major surgery. *Anesthesiology*. 2002;97:820-826.
193. Venn R, Steele A, Richardson P, et al. Randomized controlled trial to investigate influence of the fluid challenge on duration of hospital stay and perioperative morbidity in patients with hip fractures. *Br J Anaesth*. 2002;88:65-71.
194. Sinclair S, James S, Singer M. Intraoperative intravascular volume optimisation and length of hospital stay after repair of proximal femoral fracture: randomised controlled trial. *BMJ*. 1997;315:909-912.
195. Wakeling HG, McFall MR, Jenkins CS, et al. Intraoperative oesophageal Doppler guided fluid management shortens postoperative hospital stay after major bowel surgery. *Br J Anaesth*. 2005;95:634-642.
196. Noblett SE, Snowden CP, Shenton BK, Horgan AF. Randomized clinical trial assessing the effect of Doppler-optimized fluid management on outcome after elective colorectal resection. *Br J Surg*. 2006;93:1069-1076.
197. Shoemaker WC, Appel PL, Kram HB. Tissue oxygen debt as a determinant of lethal and nonlethal postoperative organ failure. *Crit Care Med*. 1988;16:1117-1120.
198. Rivers E, Nguyen B, Havstad S, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med*. 2001;345:1368-1377.
199. Brandstrup B, Tonnesen H, Beier-Holgersen R, et al. Effects of intravenous fluid restriction on postoperative complications: comparison of two perioperative fluid regimens: a randomized assessor-blinded multicenter trial. *Ann Surg*. 2003;238:641-648.
200. Nisanevich V, Felsenstein I, Almogy G, et al. Effect of intraoperative fluid management on outcome after intraabdominal surgery. *Anesthesiology*. 2005;103:25-32.
201. Lobo DN, Bostock KA, Neal KR, et al. Effect of salt and water balance on recovery of gastrointestinal function after elective colonic resection: a randomised controlled trial. *Lancet*. 2002;359:1812-1818.
202. Tambyraja AL, Sengupta F, MacGregor AB, et al. Patterns and clinical outcomes associated with routine intravenous sodium and fluid administration after colorectal resection. *World J Surg*. 2004;28:1046-1051.
203. Solus-Biguenet H, Fleyfel M, Tavernier B, et al. Non-invasive prediction of fluid responsiveness during major hepatic surgery. *Br J Anaesth*. 2006;97:808-816.