

Chapter 48

REFERENCES

1. Macintosh R, Bryce-Smith R. *Local Analgesia—Abdominal Surgery*. 2nd ed. Edinburgh, Scotland: Livingstone; 1962.
2. Karmakar MK. Thoracic paravertebral block. *Anesthesiology*. 2001;95:771-780.
3. Eason MJ, Wyatt R. Paravertebral thoracic block—a reappraisal. *Anaesthesia*. 1979;34:638-642.
4. Richardson J. Fin-de-siecle renaissance of paravertebral analgesia. *Pain Rev*. 1997;4:159-171.
5. Conacher I. Resin injection of thoracic paravertebral spaces. *Br J Anaesth*. 1988;61:657-661.
6. Karmakar MK. Paravertebral somatic nerve block for outpatient inguinal herniorrhaphy [comment]. *Reg Anesth Pain Med*. 1999;24:96-97.
7. Lonnqvist PA, Hesser U. Radiological and clinical distribution of thoracic paravertebral blockade in infants and children. *Paediatr Anaesth*. 1993;3:83-87.
8. Purcell-Jones G, Pither CE, Justins DM. Paravertebral somatic nerve block: a clinical, radiographic, and computed tomographic study in chronic pain patients. *Anesth Analg*. 1989;68:32-39.
9. Tenicela R, Pollan SB. Paravertebral-peridural block technique: a unilateral thoracic block. *Clin J Pain*. 1990;6:227-234.
10. Karmakar MK, Kwok WH, Kew J. Thoracic paravertebral block: radiological evidence of contralateral spread anterior to the vertebral bodies [see comment]. *Br J Anaesth*. 2000;84:263-265.
11. Lonnqvist PA, Hildingsson U. The caudal boundary of the thoracic paravertebral space. A study in human cadavers. *Anaesthesia*. 1992;47:1051-1052.
12. Cheema SP, Ilsley D, Richardson J, Sabanathan S. A thermographic study of paravertebral analgesia. *Anaesthesia*. 1995;50:118-121.
13. Saito T, Den S, Tanuma K, Tanuma Y, Carney E, Carlsson C. Anatomical bases for paravertebral anesthetic block: fluid communication between the thoracic and lumbar paravertebral regions. *Surg Radiol Anat*. 1999;21:359-363.
14. Dugan DJ, Samson PC. Surgical significance of the endothoracic fascia. The anatomic basis for empyectomy and other extrapleural technics. *Am J Surg*. 1975;130:151-158.
15. Im J, Webb WR, Rosen A, Gamsu G. Costal pleura: appearances at high-resolution CT. *Radiology*. 1989;171:125-131.
16. Saito T, Den S, Cheema SP, et al. A single-injection, multi-segmental paravertebral block—extension of somatosensory and sympathetic block in volunteers [see comment]. *Acta Anaesthesiol Scand*. 2001;45:30-33.
17. Nunn JF, Slavin G. Posterior intercostal nerve block for pain relief after cholecystectomy. Anatomical basis and efficacy. *Br J Anaesth*. 1980;52:253-260.
18. Klein SM, Nielsen KC, Ahmed N, Buckenmaier CC, Steele SM. In situ images of the thoracic paravertebral space. *Reg Anesth Pain Med*. 2004;29:596-599.
19. Kittredge RD. Computed tomographic evaluation of the thoracic prevertebral and paravertebral spaces. *J Comput Tomogr*. 1983;7:239-250.
20. Nel L, Conacher ID, Shanahan D. Lymphatic drainage of the thoracic paravertebral space [comment]. *Br J Anaesth*. 2001;86:453-454.
21. Naja MZ, Gustafsson AC, Ziade MF, et al. Distance between the skin and the thoracic paravertebral space. *Anaesthesia*. 2005;60:680-684.
22. Karmakar MK, Gin T, Ho AM. Ipsilateral thoraco-lumbar anaesthesia and paravertebral spread after low thoracic paravertebral injection. *Br J Anaesth*. 2001;87:312-316.
23. Richardson J, Lonnqvist PA. Thoracic paravertebral block. *Br J Anaesth*. 1998;81:230-238.
24. Richardson J, Cheema SP, Hawkins J, Sabanathan S. Thoracic paravertebral space location. A new method using pressure measurement. *Anaesthesia*. 1996;51:137-139.
25. Greengrass R, O'Brien F, Lyerly K, et al. Paravertebral block for breast cancer surgery. *Can J Anaesth*. 1996;43:858-861.
26. Lang SA. The use of a nerve stimulator for thoracic paravertebral block [comment]. *Anesthesiology*. 2002;97:521; author reply 522.
27. Hara K, Sakura S, Nomura T, Saito Y. Ultrasound guided thoracic paravertebral block in breast surgery. *Anaesthesia*. 2009;64:223-225.
28. Pusch F, Wildling E, Klimscha W, Weinstabl C. Sonographic measurement of needle insertion depth in paravertebral blocks in women [see comment]. *Br J Anaesth*. 2000;85:841-843.
29. Karmakar MK. Ultrasound-guided thoracic paravertebral block. *Tech Reg Anesth Pain Manage*. 2009;13:142-149.
30. Naja MZ, Ziade MF, El Rajab M, El Tayara K, Lonnqvist PA. Varying anatomical injection points within the thoracic paravertebral space: effect on spread of solution and nerve blockade. *Anaesthesia*. 2004;59:459-463.
31. Luyet C, Eichenberger U, Greif R, Vogt A, Szucs Farkas Z, Moriggl B. Ultrasound-guided paravertebral puncture and placement of catheters in human cadavers: an imaging study. *Br J Anaesth*. 2009;102:534-539.

32. Riain SCi, Donnell BO, Cuffe T, Harmon DC, Fraher JP, Shorten G. Thoracic paravertebral block using real-time ultrasound guidance. *Anesth Analg*. 2010;110:248-251.
33. Cowie B, McGlade D, Ivanusic J, Barrington MJ. Ultrasound-guided thoracic paravertebral blockade: a cadaveric study. *Anesth Analg*. 2010;110:1735-1739.
34. Renes SH, Bruhn J, Gielen MJ, Scheffer GJ, van Geffen GJ. In-plane ultrasound-guided thoracic paravertebral block: a preliminary report of 36 cases with radiologic confirmation of catheter position. *Reg Anesth Pain Med*. 2010;35:212-216.
35. Burns DA, Ben-David B, Chelly JE, Greensmith JE. Intercostally placed paravertebral catheterization: an alternative approach to continuous paravertebral blockade. *Anesth Analg*. 2008;107:339-341.
36. Gulbahar G, Kocer B, Muratli SN, et al. A comparison of epidural and paravertebral catheterisation techniques in post-thoracotomy pain management. *Eur J Cardiothorac Surg*. 2010;37:467-472.
37. Garutti I, Gonzalez-Aragoneses F, Biencinto MT, et al. Thoracic paravertebral block after thoracotomy: comparison of three different approaches. *Eur J Cardiothorac Surg*. 2009;35:829-832.
38. Dauphin A, Lubanska-Hubert E, Young EM, Miller JD, Bennett WF, Fuller HD. Comparative study of continuous extrapleural intercostal nerve block and lumbar epidural morphine in post thoracotomy pain. *Can J Surg*. 1997;40:431-436.
39. Sabanathan S, Richardson J, Shah R. 1988: continuous intercostal nerve block for pain relief after thoracotomy. Updated in 1995. *Ann Thorac Surg*. 1995;59:1261-1263.
40. Bimston DN, McGee JP, Liptay MJ, Fry WA. Continuous paravertebral extrapleural infusion for post-thoracotomy pain management. *Surgery*. 1999;126:650-656; discussion 656-657.
41. Soni AK, Conacher ID, Waller DA, Hilton CJ. Video-assisted thoracoscopic placement of paravertebral catheters: a technique for postoperative analgesia for bilateral thoracoscopic surgery. *Br J Anaesth*. 1994;72:462-464.
42. Feltracco P, Ori C. A new look at the paravertebral block: a percutaneous video-assisted technique. *Reg Anesth Pain Med*. 2007;32:538-539.
43. Klein SM, Bergh A, Steele SM, Georgiade GS, Greengrass RA. Thoracic paravertebral block for breast surgery. *Anesth Analg*. 2000;90:1402-1405.
44. Kairaluoma PM, Bachmann MS, Korpinen AK, Rosenberg PH, Pere PJ. Single-injection paravertebral block before general anesthesia enhances analgesia after breast cancer surgery with and without associated lymph node biopsy. *Anesth Analg*. 2004;99:1837-1843.
45. Terheggen MA, Wille F, Borel Rinkes IH, Ionescu TI, Knape JT. Paravertebral blockade for minor breast surgery. *Anesth Analg*. 2002;94:355-359.
46. Naja MZ, Ziade MF, Lonnqvist PA. Nerve-stimulator guided paravertebral blockade vs. general anaesthesia for breast surgery: a prospective randomized trial. *Eur J Anaesthesiol*. 2003;20:897-903.
47. Coveney E, Weltz CR, Greengrass R, et al. Use of paravertebral block anesthesia in the surgical management of breast cancer: experience in 156 cases. *Ann Surg*. 1998;227:496-501.
48. Cheema S, Richardson J, McGurgan P. Factors affecting the spread of bupivacaine in the adult thoracic paravertebral space. *Anaesthesia*. 2003;58:684-687.
49. Greengrass R, Buckenmaier CC 3rd. Paravertebral anaesthesia/analgesia for ambulatory surgery. *Best Pract Res Clin Anaesthesiol*. 2002;16:271-283.
50. Naja ZM, El-Rajab M, Al-Tannir MA, et al. Thoracic paravertebral block: influence of the number of injections. *Reg Anesth Pain Med*. 2006;31:196-201.
51. Saito T, Gallagher ET, Cutler S, et al. Extended unilateral anesthesia. New technique or paravertebral anesthesia? *Reg Anesth*. 1996;21:304-307.
52. Hura G, Knapik P, Misiolek H, Krakus A, Karpe J. Sensory blockade after thoracic paravertebral injection of ropivacaine or bupivacaine. *Eur J Anaesthesiol*. 2006;23:658-664.
53. Navlet MG, Garutti I, Olmedilla L, et al. Paravertebral ropivacaine, 0.3%, and bupivacaine, 0.25%, provide similar pain relief after thoracotomy. *J Cardiothorac Vasc Anesth*. 2006;20:644-647.
54. Kotze A, Scally A, Howell S. Efficacy and safety of different techniques of paravertebral block for analgesia after thoracotomy: a systematic review and metaregression. *Br J Anaesth*. 2009;103:626-636.
55. Karmakar MK, Ho AM, Law BK, Wong ASY, Shafer SL, Gin T. Arterial and venous pharmacokinetics of ropivacaine with and without epinephrine after thoracic paravertebral block. *Anesthesiology*. 2005;103:704-711.
56. Naja Z, Ziade MF, Lonnqvist PA. Bilateral paravertebral somatic nerve block for ventral hernia repair. *Eur J Anaesthesiol*. 2002;19:197-202.
57. Richardson J, Jones J, Atkinson R. The effect of thoracic paravertebral blockade on intercostal somatosensory evoked potentials. *Anesth Analg*. 1998;87:373-376.
58. Dahl JB, Rosenberg J, Lund C, Kehlet H. Effect of thoracic epidural bupivacaine 0.75% on somatosensory evoked potentials after dermatomal stimulation. *Reg Anesth*. 1990;15:73-75.
59. Iohom G, Abdalla H, O'Brien J, et al. The associations between severity of early postoperative pain, chronic postsurgical pain and plasma concentration of stable nitric oxide products after breast surgery. *Anesth Analg*. 2006;103:995-1000.
60. Kairaluoma PM, Bachmann MS, Rosenberg PH, Pere PJ. Preincisional paravertebral block reduces the prevalence of chronic pain after breast surgery. *Anesth Analg*. 2006;103:703-708.
61. Richardson J, Sabanathan S, Mearns AJ, Sides C, Goulden C. Post-thoracotomy neuralgia. *Pain Clinic*. 1994;7:87-97.
62. Giesecke K, Hamberger B, Jarnberg PO, Klingstedt D. Paravertebral block during cholecystectomy: effects on circulatory and hormonal responses. *Br J Anaesth*. 1988;61:652-656.
63. Berrisford RG, Sabanathan S, Mearns AJ, Bickford-Smith PJ. Pulmonary complications after lung resection: the effect of continuous extrapleural intercostal nerve block. *Eur J Cardiothorac Surg*. 1990;4:407-410.
64. Sabanathan S, Bibby SR, Berrisford RG. Efficacy of continuous intercostal nerve blockade on postoperative analgesia and pulmonary mechanics. *Br J Surg*. 1989;77:221-225.
65. Sabanathan S, Mearns AJ, Bickford-Smith PJ, et al. Efficacy of continuous extrapleural intercostal nerve block on post-thoracotomy pain and pulmonary mechanics. *Br J Surg*. 1990;77:221-225.
66. Eng J, Sabanathan S. Continuous extrapleural intercostal nerve block and post-thoracotomy pulmonary complications. *Scand J Thorac Cardiovasc Surg*. 1992;26:219-223.

67. Barron DJ, Tolan MJ, Lea RE. A randomized controlled trial of continuous extra-pleural analgesia post-thoracotomy: efficacy and choice of local anaesthetic. *Eur J Anaesthesiol*. 1999;16:236-245.
68. Pusch F, Freitag H, Weinstabl C, Obwegeser R, Huber E, Wildling E. Single-injection paravertebral block compared to general anaesthesia in breast surgery. *Acta Anaesthesiol Scand*. 1999;43:770-774.
69. Matthews PJ, Govenden V. Comparison of continuous paravertebral and extradural infusions of bupivacaine for pain relief after thoracotomy. *Br J Anaesth*. 1989;62:204-205.
70. Exadaktylos AK, Buggy DJ, Moriarty DC, Mascha E, Sessler DI. Can anesthetic technique for primary breast cancer surgery affect recurrence or metastasis? *Anesthesiology*. 2006;105:660-664.
71. Richardson J, Sabanathan S. Thoracic paravertebral analgesia [see comment]. *Acta Anaesthesiol Scand*. 1995;39:1005-1015.
72. Naja Z, Lonnqvist PA. Somatic paravertebral nerve blockade. Incidence of failed block and complications. *Anaesthesia*. 2001;56:1184-1188.
73. Lonnqvist PA, MacKenzie J, Soni AK, Conacher ID. Paravertebral blockade. Failure rate and complications. *Anaesthesia*. 1995;50:813-815.
74. Thomas PW, Sanders DJ, Berrisford RG. Pulmonary haemorrhage after percutaneous paravertebral block [see comment]. *Br J Anaesth*. 1999;83:668-669.
75. Lekhak B, Bartley C, Conacher ID, Nouraei SM. Total spinal anaesthesia in association with insertion of a paravertebral catheter. *Br J Anaesth*. 2001;86:280-282.
76. Sharrock NE. Postural headache following thoracic somatic paravertebral nerve block. *Anesthesiology*. 1980;52:360-362.
77. Richardson J, Lonnqvist PA. Thoracic paravertebral block [see comment]. *Br J Anaesth*. 1998;81:230-238.
78. Snowden CP, Bower S, Conacher I. Plasma bupivacaine levels in paravertebral blockade in adults [comment]. *Anaesthesia*. 1994;49:546.
79. Berrisford RG, Sabanathan S, Mearns AJ, Clarke BJ, Hamdi A. Plasma concentrations of bupivacaine and its enantiomers during continuous extrapleural intercostal nerve block. *Br J Anaesth*. 1993;70:201-204.
80. Karmakar MK, Booker PD, Franks R, Pozzi M. Continuous extrapleural paravertebral infusion of bupivacaine for post-thoracotomy analgesia in young infants. *Br J Anaesth*. 1996;76:811-815.
81. Cheung SLW, Booker PD, Franks R, Pozzi M. Serum concentrations of bupivacaine during prolonged continuous paravertebral infusion in young infants. *Br J Anaesth*. 1997;79:9-13.
82. Karmakar MK, Ho AM-H, Law BK, Wong ASY, Shafer SL, Gin T. Arterial and venous pharmacokinetics of ropivacaine with and without epinephrine after thoracic paravertebral block. *Anesthesiology*. 2005;103:704-711.
83. Dauphin A, Gupta RN, Young JEM, Morton WD. Serum bupivacaine concentrations during continuous extrapleural infusion. *Can J Anaesth*. 1997;44:367-370.
84. Clark BJ, Hamdi A, Berrisford RG, Sabanathan S, Mearns AJ. Reversed-phase and chiral high-performance liquid chromatographic assay of bupivacaine and its enantiomers in clinical samples after continuous extrapleural infusion. *J Chromatogr*. 1991;553:383-390.
85. Brittingham TE, Berlin LN, Wolff HG. Nervous system damage following paravertebral block with efocaine: report of three cases. *JAMA*. 1954;154:329-330.
86. Molitch M, Wilson G. Brown-sequard paralysis following a paravertebral alcohol injection for angina pectoris. *JAMA*. 1931;97:247.
87. Bigler D, Dirkes W, Hansen R, Rosenberg J, Kehlet H. Effects of thoracic paravertebral block with bupivacaine versus combined thoracic epidural block with bupivacaine and morphine on pain and pulmonary function after cholecystectomy. *Acta Anaesthesiol Scand*. 1989;33:561-564.
88. Thavaneswaran P, Rudkin GE, Cooter RD, Moyes DG, Perera CL, Maddern GJ. Paravertebral block for anesthesia: a systematic review. *Anesth Analg*. 2010;110:1740-1744.
89. Joshi GP, Bonnet F, Shah R, et al. A systematic review of randomized trials evaluating regional techniques for postthoracotomy analgesia. *Anesth Analg*. 2008;107:1026-1040.
90. Messina M, Boroli F, Landoni G, et al. A comparison of epidural vs. paravertebral blockade in thoracic surgery. *Minerva Anestesiol*. 2009;75:616-621.
91. Marret E, Bazelly B, Taylor G, et al. Paravertebral block with ropivacaine 0.5% versus systemic analgesia for pain relief after thoracotomy. *Ann Thorac Surg*. 2005;79:2109-2113.
92. Hill SE, Keller RA, Stafford-Smith M, et al. Efficacy of single-dose, multilevel paravertebral nerve blockade for analgesia after thoracoscopic procedures. *Anesthesiology*. 2006;104:1047-1053.
93. Scarci M, Joshi A, Attia R. In patients undergoing thoracic surgery is paravertebral block as effective as epidural analgesia for pain management? *Interact Cardiovasc Thorac Surg*. 2010;10:92-96.
94. Vogt A, Stieger DS, Theurillat C, Curatolo M. Single-injection thoracic paravertebral block for postoperative pain treatment after thoracoscopic surgery. *Br J Anaesth*. 2005;95:816-821.
95. Kaya FN, Turker G, Basagan-Mogol E, Goren S, Bayram S, Gebitekin C. Preoperative multiple-injection thoracic paravertebral blocks reduce postoperative pain and analgesic requirements after video-assisted thoracic surgery. *J Cardiothorac Vasc Anesth*. 2006;20:639-643.
96. Shah R, Sabanathan S, Richardson J, Mearns AJ, Bembridge J. Continuous paravertebral block for post thoracotomy analgesia in children. *J Cardiovasc Surg*. 1997;38:543-546.
97. Fibla JJ, Molins L, Mier JM, Sierra A, Vidal G. A prospective study of analgesic quality after a thoracotomy: paravertebral block with ropivacaine before and after rib spreading. *Eur J Cardiothorac Surg*. 2009;36:901-905.
98. Perttunen K, Nilsson E, Heinonen J, Hirvisalo EL, Salo JA, Kalso E. Extradural, paravertebral and intercostal nerve blocks for post-thoracotomy pain. *Br J Anaesth*. 1995;75:541-547.
99. Richardson J, Sabanathan S, Mearns AJ, Shah RD, Goulden C. A prospective, randomized comparison of interpleural and paravertebral analgesia in thoracic surgery. *Br J Anaesth*. 1995;75:405-408.
100. Richardson J, Sabanathan S, Shah RD, Clarke BJ, Cheema S, Mearns AJ. Pleural bupivacaine placement for optimal post-thoracotomy pulmonary function: a prospective, randomized study. *J Cardiothorac Vasc Anesth*. 1998;12:166-169.
101. Richardson J, Sabanathan S, Jones J, Shah RD, Cheema S, Mearns AJ. A prospective, randomized comparison of preoperative and continuous balanced epidural or paravertebral bupivacaine on post-thoracotomy pain, pulmonary function and stress responses. *Br J Anaesth*. 1999;83:387-392.
102. Kaiser AM, Zollinger A, De Lorenzi D, Largiader F, Weder W. Prospective, randomized comparison of extrapleural versus epidural analgesia for postthoracotomy pain. *Ann Thorac Surg*. 1998;66:367-372.

103. Mozell EJ, Sabanathan S, Mearns AJ, Bickford-Smith PJ, Majid MR, Zografos G. Continuous extrapleural intercostal nerve block after pleurectomy. *Thorax*. 1991;46:21-24.
104. Patel AN, Finlay KU, Schyra KC, et al. Use of general anesthetic only vs general anesthetic combined with paravertebral block for perioperative pain management after first rib resection. *Proc (Bayl Univ Med Cent)*. 2002;15:374-375.
105. Mohta M, Verma P, Saxena AK, Sethi AK, Tyagi A, Girotra G. Prospective, randomized comparison of continuous thoracic epidural and thoracic paravertebral infusion in patients with unilateral multiple fractured ribs—a pilot study. *J Trauma*. 2009;66:1096-1101.
106. Williamson S, Kumar CM. Paravertebral block in head injured patient with chest trauma. *Anaesthesia*. 1997;52:284-285.
107. Karmakar MK, Chui PT, Joynt GM, Ho AM. Thoracic paravertebral block for management of pain associated with multiple fractured ribs in patients with concomitant lumbar spinal trauma. *Reg Anesth Pain Med*. 2001;26:169-173.
108. Gilbert J, Hultman J. Thoracic paravertebral block: a method of pain control. *Acta Anaesthesiol Scand*. 1989;33:142-145.
109. Paniagua P, Catala E, Villar Landeira JM. Successful management of pleuritic pain with thoracic paravertebral block. *Reg Anesth Pain Med*. 2000;25:651-653.
110. Weltz CR, Greengrass RA, Lyerly HK. Ambulatory surgical management of breast carcinoma using paravertebral block. *Ann Surg*. 1995;222:19-26.
111. Boughey JC, Goravanchi F, Parris RN, et al. Prospective randomized trial of paravertebral block for patients undergoing breast cancer surgery. *Am J Surg*. 2009;198:720-725.
112. Moller JF, Nikolajsen L, Rodt SA, Ronning H, Carlsson PS. Thoracic paravertebral block for breast cancer surgery: a randomized double-blind study. *Anesth Analg*. 2007;105:1848-1851.
113. Sidiropoulou T, Buonomo O, Fabbi E, et al. A prospective comparison of continuous wound infiltration with ropivacaine versus single-injection paravertebral block after modified radical mastectomy. *Anesth Analg*. 2008;106:997-1001.
114. Klein SM, Greengrass RA, Weltz C, Warner DS. Paravertebral somatic nerve block for outpatient inguinal herniorrhaphy: an expanded case report of 22 patients. *Reg Anesth Pain Med*. 1998;23:306-310.
115. Weltz CR, Klein SM, Arbo JE, Greengrass RA. Paravertebral block anesthesia for inguinal hernia repair. *World J Surg*. 2003;27:425-429.
116. Hadzic A, Kerimoglu B, Loreio D, et al. Paravertebral blocks provide superior same-day recovery over general anesthesia for patients undergoing inguinal hernia repair. *Anesth Analg*. 2006;102:1076-1081.
117. Bhattacharya P, Mandal MC, Mukhopadhyay S, Das S, Pal PP, Basu SR. Unilateral paravertebral block: an alternative to conventional spinal anaesthesia for inguinal hernia repair. *Acta Anaesthesiol Scand*. 2010;54:246-251.
118. Akcaboy EY, Akcaboy ZN, Gogus N. Ambulatory inguinal herniorrhaphy: paravertebral block versus spinal anesthesia. *Minerva Anesthesiol*. 2009;75:684-691.
119. Naja MZ, el Hassan MJ, Oweidat M, Zbibo R, Ziade MF, Lonnqvist PA. Paravertebral blockade vs general anesthesia or spinal anesthesia for inguinal hernia repair. *Middle East J Anesthesiol*. 2001;16:201-210.
120. Wassef MR, Randazzo T, Ward W. The paravertebral nerve root block for inguinal herniorrhaphy—a comparison with the field block approach. *Reg Anesth Pain Med*. 1998;23:451-456.
121. Klein SM, Pietrobon R, Nielsen KC, et al. Paravertebral somatic nerve block compared with peripheral nerve blocks for outpatient inguinal herniorrhaphy. *Reg Anesth Pain Med*. 2002;27:476-480.
122. Naja ZM, Raf M, El Rajab M, Ziade FM, Al Tannir MA, Lonnqvist PA. Nerve stimulator-guided paravertebral blockade combined with sevoflurane sedation versus general anesthesia with systemic analgesia for postherniorrhaphy pain relief in children: a prospective randomized trial. *Anesthesiology*. 2005;103:600-605.
123. Naja ZM, Raf M, Rajab ME, et al. A comparison of nerve stimulator guided paravertebral block and ilio-inguinal nerve block for analgesia after inguinal herniorrhaphy in children. *Anaesthesia*. 2006;61:1064-1068.
124. Ozkan D, Akkaya T, Comert A, et al. Paravertebral block in inguinal hernia surgeries: two segments or 4 segments? *Reg Anesth Pain Med*. 2009;34:312-315.
125. Kalady MF, Fields RC, Klein S, et al. Loop ileostomy closure at an ambulatory surgery facility: a safe and cost-effective alternative to routine hospitalization. *Dis Colon Rectum*. 2003;46:486-490.
126. Awwad ZM, Atiyat BA. Pain relief using continuous bupivacaine infusion in the paravertebral space after loin incision. *Saudi Med J*. 2004;25:1369-1373.
127. Lonnqvist PA. Continuous paravertebral block in children. Initial experience. *Anaesthesia*. 1992;47:607-609.
128. Lonnqvist PA, Olsson GL. Paravertebral vs epidural block in children. Effects on postoperative morphine requirement after renal surgery. *Acta Anaesthesiol Scand*. 1994;38:346-349.
129. Wolf JS Jr, Marcovich R, Merion RM, Konnak JW. Prospective, case matched comparison of hand assisted laparoscopic and open surgical live donor nephrectomy. *J Urol*. 2000;163:1650-1653.
130. Bachmann A, Wolff T, Giannini O, et al. How painful is donor nephrectomy? Retrospective analysis of early pain and pain management in open versus laparoscopic versus retroperitoneoscopic nephrectomy. *Transplantation*. 2006;81:1735-1738.
131. Clendenen SR, Wehle MJ, Rodriguez GA, Greengrass RA. Paravertebral block provides significant opioid sparing after hand-assisted laparoscopic nephrectomy: an expanded case report of 30 patients. *J Endourol*. 2009;23:1979-1983.
132. Jamieson BD, Mariano ER. Thoracic and lumbar paravertebral blocks for outpatient lithotripsy. *J Clin Anesth*. 2007;19:149-151.
133. Naja MZ, Ziade MF, Lonnqvist PA. General anesthesia combined with bilateral paravertebral blockade (T5-6) vs. general anesthesia for laparoscopic cholecystectomy: a prospective, randomized controlled trial. *Eur J Anaesthesiol*. 2004;21:489-495.
134. Paleczny J, Zipser P, Pysz M. [Paravertebral block for open cholecystectomy]. *Anestezjol Intens Ter*. 2009;41:89-93.
135. Splinter WM, Thomson ME. Somatic paravertebral block decreases opioid requirements in children undergoing appendectomy. *Can J Anaesth*. 2010;57:206-210.
136. Bogoch ER, Henke M, Mackenzie T, Olschewski E, Mahomed NN. Lumbar paravertebral nerve block in the management of pain after total hip and knee arthroplasty: a randomized controlled clinical trial. *J Arthroplasty*. 2002;17:398-401.
137. Canto M, Sanchez MJ, Casas MA, Bataller ML. Bilateral paravertebral blockade for conventional cardiac surgery. *Anaesthesia*. 2003;58:365-370.

138. Ganapathy S, Murkin JM, Boyd DW, Dobkowski W, Morgan J. Continuous percutaneous paravertebral block for minimally invasive cardiac surgery. *J Cardiothorac Vasc Anesth.* 1999;13:594-596.
139. Lynch JJ, Mauermann WJ, Pulido JN, Rehfeldt KH, Torres NE. Use of paravertebral blockade to facilitate early extubation after minimally invasive cardiac surgery. *Semin Cardiothorac Vasc Anesth.* 2010;14:47-48.
140. Dhole S, Mehta Y, Saxena H, Juneja R, Trehan N. Comparison of continuous thoracic epidural and paravertebral blocks for postoperative analgesia after minimally invasive direct coronary artery bypass surgery. *J Cardiothorac Vasc Anesth.* 2001;15:288-292.
141. Mehta Y, Arora D, Sharma K, Mishra Y, Wasir H, Trehan N. Comparison of continuous thoracic epidural and paravertebral block for postoperative analgesia after robotic-assisted coronary artery bypass surgery. *Ann Card Anaesth.* 2008; 11:91-96.
142. Tsai T, Rodriguez-Diaz C, Deschner B, Thomas K, Wasnick JD. Thoracic paravertebral block for implantable cardioverter-defibrillator and laser lead extraction. *J Clin Anesth.* 2008;20:379-82.
143. Richardson J, Vowden P, Sabanathan S. Bilateral paravertebral analgesia for major abdominal vascular surgery: a preliminary report. *Anaesthesia.* 1995;50:995-998.
144. Falkensammer J, Hakaim AG, Klocker J, et al. Paravertebral blockade with propofol sedation versus general anesthesia for elective endovascular abdominal aortic aneurysm repair. *Vascular.* 2006;14:17-22.
145. Basagan-Mogol E, Turker G, Yilmaz M, Goren S. Combination of a psoas compartment, sciatic nerve, and T12-L1 paravertebral blocks for femoropopliteal bypass surgery in a high-risk patient. *J Cardiothorac Vasc Anesth.* 2008;22: 337-339.
146. Ho AM, Karmakar MK, Cheung M, Lam GC. Right thoracic paravertebral analgesia for hepatectomy. *Br J Anaesth.* 2004;93:458-461.
147. Hall H, Leach A. Paravertebral block in the management of liver capsule pain after blunt trauma. *Br J Anaesth.* 1999;83: 819-821.
148. Culp WC, McCowan TC, DeValdenebro M, Wright LB, Workman JL, Culp WC Jr. Paravertebral block: an improved method of pain control in percutaneous transhepatic biliary drainage. *Cardiovasc Intervent Radiol.* 2006;29:1015-1021.
149. Culp WC, Payne MN, Montgomery ML. Thoracic paravertebral block for analgesia following liver mass radiofrequency ablation. *Br J Radiol.* 2008;81:e23-e25.
150. Nair V, Henry R. Bilateral paravertebral block: a satisfactory alternative for labour analgesia. *Can J Anaesth.* 2001;48: 179-184.
151. Suelto MD. Paravertebral lumbar sympathetic block for labor analgesia [comment]. *Anesthesiology.* 2000;93:580.
152. Kirvela O, Antila H. Thoracic paravertebral block in chronic postoperative pain [see comment]. *Reg Anesth.* 1992;17: 348-350.
153. Antila H, Kirvela O. Neurolytic thoracic paravertebral block in cancer pain. A clinical report. *Acta Anaesthesiol Scand.* 1998;42:581-585.
154. Naja ZM, Al-Tannir MA, Zeidan A, El-Rajab M, Ziade F, Baraka A. Nerve stimulator-guided repetitive paravertebral block for thoracic myofascial pain syndrome. *Pain Pract.* 2007;7:348-351.
155. Johnson LR, Rocco AG, Ferrante FM. Continuous subpleural-paravertebral block in acute thoracic herpes zoster. *Anesth Analg.* 1988;67:1105-1108.
156. Desai PM. Thoracic paravertebral block [comment]. *Br J Anaesth.* 1999;82:149-150.
157. Greengrass R, Steele S. Paravertebral blocks for breast surgery. *Techniques in Regional Anesthesia and Pain Management* 1998;2:8-12.
158. Berrisford RG, Sabanathan SS. Direct access to the paravertebral space at thoracotomy.[comment]. *Annals of Thoracic Surgery* 1990;49:854.