

Chapter 49

REFERENCES

1. Brown A, Weiss R, Greenberg C, Flatow EL, Bigliani LU. Interscalene block for shoulder arthroscopy: comparison with general anesthesia. *Arthroscopy*. 1993;9(3):295-300.
2. Borgeat A, Ekatodramis G, Schenker CA. Postoperative nausea and vomiting in regional anesthesia. *Anesthesiology*. 2003;98(2):530-547.
3. Gregory SR, Cole TR. The changing role of dissection in medical education. *JAMA*. 2002;287:1180-1181.
4. Greenberg SA. The history of dermatome mapping. *Arch Neurol*. 2003;60:126-131.
5. Deschner B, Robards C, et al. In: Hadzić A. *Textbook of Regional Anesthesia and Acute Pain Management*. New York, NY: McGraw-Hill; 2007:3-18.
6. Fink B. Leaves and needles: the introduction of surgical local anesthesia. *Anesthesiology*. 1985;63(1):77.
7. Labat G. *Regional Anesthesia: Its Technic and Clinical Application*. Philadelphia, PA: WB Saunders; 1922.
8. Wedel DJ, Horlocker TT. In: Longnecker DE. *Anesthesiology*. New York, NY: McGraw-Hill; 2008:1025-1052.
9. Mandabach MG, Wright AJ. The American Society of Regional Anesthesia: a concise history of the original group—its birth, growth, and eventual dissolution. *Reg Anesth Pain Med*. 2006;31(1):53-65.
10. Ilfeld BM, Le LT, Ramjohn J, et al. The effects of local anesthetic concentration and dose on continuous infraclavicular nerve blocks: a multicenter, randomized, observer-masked, controlled study. *Anesth Analg*. 2009;108(1):345-350.
11. Ilfeld BM, Moeller LK, Mariano ER, et al. Continuous peripheral nerve blocks. Is local anesthetic dose the only factor, or do concentration and volume influence infusion effects as well? *Anesthesiology*. 2010;112:347-354.
12. Seet E, Leong WL, Yeo AS, Fook-Chong S. Effectiveness of 3-in-1 continuous femoral block of differing concentrations compared to patient controlled intravenous morphine for post total knee arthroplasty analgesia and knee rehabilitation. *Anaesth Intensive Care*. 2006;34:25-30.
13. Mulroy MF. Systemic toxicity and cardiotoxicity from local anesthetics: incidence and preventive measures. *Reg Anesth Pain Med*. 2002;27(6):556-561.
14. Weinberg G, Ripper R, Feinstein DL, et al. Lipid emulsion infusion rescues dogs from bupivacaine-induced cardiac toxicity. *Reg Anesth Pain Med*. 2003;28(3):198-202.
15. Rosenblatt MA, Abel M, Fischer GW, et al. Successful use of a 20% lipid emulsion to resuscitate a patient after a presumed bupivacaine-related cardiac arrest. *Anesthesiology*. 2006;105(1):217-218.
16. McGlade DP, Kalpokas MV, Mooney PH, et al. A comparison of 0.5% ropivacaine and 0.5% bupivacaine for axillary brachial plexus anaesthesia. *Anaesth Intensive Care*. 1998;26(5):515-520.
17. Dony P, Dewinde V, Vanderick B, et al. The comparative toxicity of ropivacaine and bupivacaine at equipotent doses in rats. *Anesth Analg*. 2000;91(6):1489-1492.
18. Singelyn FJ, Dangoisse M, Bartholome S, et al. Adding clonidine to mepivacaine prolongs the duration of anesthesia and analgesia after axillary brachial plexus block. *Reg Anesth Pain Med*. 1992;17(3):148-150.
19. Culebras X, Van Gessel E, Hoffmeyer P, et al. Clonidine combined with a long acting local anesthetic does not prolong postoperative analgesia after brachial plexus block but does induce hemodynamic changes. *Anesth Analg*. 2001;92(1):199-204.
20. Movafegh A, Razazian M, Hajimaohamadi F, et al. Dexamethasone added to lidocaine prolongs axillary brachial plexus blockade. *Anesth Analg*. 2006;102(1):263-267.
21. Rose JS, Neal JM, Kopacz D. Extended-duration analgesia: update on microspheres and liposomes. *Reg Anesth Pain Med*. 2005;30(3):275-285.
22. Neal JM, Hebl JR, Geranchar JC, et al. Brachial plexus anesthesia: essentials of our current understanding. *Reg Anesth Pain Med*. 2002;27(4):402-428.
23. Enneking FK, Chan V, Greger J. Lower-extremity nerve blockade: essentials of our current understanding. *Reg Anesth Pain Med*. 2005;30(1):4-35.
24. Murphy DB, McCartney CJ, Chan VW. Novel analgesic adjuncts for brachial plexus block: a systematic review. *Anesth Analg*. 2000;90(5):1122-1128.
25. Reinhart DJ, Wang W, Stagg KS, et al. Postoperative analgesia after peripheral nerve block for podiatric surgery: clinical efficacy and chemical stability of lidocaine alone versus lidocaine plus ketorolac. *Reg Anesth Pain Med*. 2000;25(5):506-513.
26. Moore DC. "No paresthesia—no analgesia," the nerve stimulator or neither? *Reg Anesth Pain Med*. 1997;22(4):388-390.
27. Beach ML, Sites BD, Gallagher JD. Use of a nerve stimulator does not improve the efficiency of ultrasound-guided supraclavicular nerve blocks. *J Clin Anesth*. 2006;18(8):580-584.

28. La Grange P, Foster P, Pretorius L. Application of the Doppler ultrasound blood flow detector in supraclavicular brachial plexus block. *Br J Anaesth.* 1978;50:965-967.
29. Ting PL, Sivagnanaratnam V. Ultrasonographic study of the spread of local anesthetic during axillary brachial plexus block. *Br J Anaesth.* 1989;63:326-329.
30. Neal JM, Brull R, Chan V, et al. The ASRA evidence-based medicine assessment of ultrasound-guided regional anesthesia and pain medicine. *Reg Anesth Pain Med.* 2010;35(2):S1-S9.
31. McCartney C, Lin L, Shastri U. Evidence basis for the use of ultrasound for upper-extremity blocks. *Reg Anesth Pain Med.* 2010;35(2):S10-S15.
32. Liu SS, Ngeow J, John RS. Evidence basis for ultrasound-guided block characteristics: onset, quality, and duration. *Reg Anesth Pain Med.* 2010;35(2):S26-S35.
33. Marhofer P, Chan VW. Ultrasound-guided regional anesthesia: current concepts and future trends. *Anesth Analg.* 2007;104:1265-1269.
34. Marhofer P, Sitzwohl C, Gerher M, Kapral S. Ultrasound guidance for infraclavicular brachial plexus anaesthesia in children. *Anaesthesia.* 2004;59(7):642-646.
35. Willschke H, Marhofer P, Bösenberg A, et al. Ultrasonography for ilioinguinal/iliohypogastric nerve blocks in children. *Br J Anaesth.* 2005;95(2):226-230.
36. Oberndorfer U, Marhofer P, Bösenberg A, et al. Ultrasonographic guidance for sciatic and femoral nerve blocks in children. *Br J Anaesth.* 2007;98(6):797-801.
37. Capdevila X, Bringuier S, Borgeat A. Infectious risk of continuous peripheral nerve blocks. *Anesthesiology.* 2009;110(1):182-188.
38. Borgeat A, Blumenthal S, Lambert M, et al. The feasibility and complications of the continuous popliteal nerve block: a 1001-case survey. *Anesth Analg.* 2006;103(1):229-233.
39. Capdevila X, Pirat P, Bringuier, et al. Continuous peripheral nerve blocks in hospital wards after orthopedic surgery: a multicenter prospective analysis of the quality of postoperative analgesia and complications in 1,416 patients. *Anesthesiology.* 2005;103(5):1035-1045.
40. Bickler P, Brandes J, Lee M, et al. Bleeding complications from femoral and sciatic nerve catheters in patients receiving low molecular weight heparin. *Anesth Analg.* 2006;103(4):1036-1037.
41. Horlocker TT, Wedel DJ, Benzon H, et al. Regional anesthesia in the anticoagulated patient: defining the risks (the second ASRA Consensus Conference on Neuraxial Anesthesia and Anticoagulation). *Reg Anesth Pain Med.* 2003;28(3):172-197.
42. Horlocker TT, Wedel DJ, Rowlingson JC, et al. Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American Society of Regional Anesthesia and Pain Medicine Evidence-Based Guidelines (Third Edition). *Reg Anesth Pain Med.* 2010;35(1):64-101.
43. Ilfeld BM, Le LT, Meyer RS, et al. Ambulatory continuous nerve blocks decrease time to discharge readiness after tricompartiment total knee arthroplasty. *Anesthesiology.* 2008;108(4):703-713.
44. Brodner G, Buerkle H, Van Aken H, et al. Postoperative analgesia after knee surgery: a comparison of three different concentrations of ropivacaine for continuous femoral nerve blockade. *Anesth Analg.* 2007;105(1):256-262.
45. Ilfeld BM, Le LT, Ramjohn J, et al. The effects of local anesthetic concentration and dose on continuous infraclavicular nerve blocks: a multicenter, randomized, observer-masked, controlled study. *Anesth Analg.* 2009;108(1):345-350.
46. Ilfeld BM, Morey TE, Enneking FK. Infraclavicular perineural local anesthetic infusion: a comparison of three dosing regimens for postoperative analgesia. *Anesthesiology.* 2004;100(2):395-402.
47. Ilfeld BM, Moeller LK, Mariano ER, et al. Continuous peripheral nerve blocks: is local anesthetic dose the only factor, or do concentration and volume influence infusion effects as well? *Anesthesiology.* 2010;112(2):347-354.
48. Cuvillon P, Nouvellon E, Ripart J, et al. A comparison of pharmacodynamics and pharmacokinetics of bupivacaine, ropivacaine (with epinephrine) and their equal volume mixtures with lidocaine used for femoral and sciatic nerve blocks: a double-blind randomized study. *Anesth Analg.* 2009;108(2):641-649.
49. Usui Y, Kobayashi T, Kakinamu H, Watanabe K, Kitajima T, Matsuno K. An anatomical basis for blocking of the deep cervical plexus and cervical sympathetic tract using an ultrasound-guided technique. *Anesth Analg.* 2010;110(3):964-968.
50. Harry WG, Bennett JD, Guha SC. Scalene muscles and the brachial plexus: anatomical variations and their clinical significance. *Clin Anat.* 1997;10:250-252.
51. Winnie AP. Interscalene brachial plexus block. *Anesth Analg.* 1970;49:455-466.
52. Antonakakis JG, Sites BD, Shiffrin J. Ultrasound-guided posterior approach for the placement of a continuous interscalene catheter. *Reg Anesth Pain Med.* 2009;34(1):64-68.
53. Chan VW, Perlas A, Rawson R, Odukoya O. Ultrasound-guided supraclavicular brachial plexus block. *Anesth Analg.* 2003;97(5):1514-1517.
54. Soares LG, Brull R, Lai J, Chan VW. Eight ball, corner pocket: the optimal needle position for ultrasound-guided supraclavicular block. *Reg Anesth Pain Med.* 2007;32(1):94-95.
55. Kapral S, Jandrasits O, Schabernig, et al. Lateral infraclavicular plexus block vs. axillary block for hand and forearm surgery. *Acta Anaesthesiol Scand.* 1999;43:1047-1052.
56. Winnie AP. *Plexus Anesthesia: Volume 1 Perivascular Techniques of Brachial Plexus Blocks.* Philadelphia, PA: WB Saunders; 1983.
57. Sandhu NS, Capan LM. Ultrasound-guided infraclavicular brachial plexus block. *Br J Anaesth.* 2002;89(2):254-259.
58. Retzl G, Kapral S, Greher M, Mauritz. Ultrasonographic findings of the axillary part of the brachial plexus. *Anesth Analg.* 2001;92:1271-1275.
59. De Jong RH. Axillary block of the brachial plexus. *Anesthesiology.* 1961;22(2):215-225.
60. Lavoie J, Martin R, Tétrault JP, Côté DJ, Colas MJ. Axillary plexus block using a peripheral nerve stimulator: single or multiple injections. *Can J Anaesth.* 1992;39(6):583-586.
61. Sites BD, Beach ML, Spence BC, et al. Ultrasound guidance improves the success rate of a perivascular axillary plexus block. *Acta Anaesthesiol Scand.* 2006;50(6):678-684.
62. Spence BC, Sites BD, Beach ML. Ultrasound-guided musculocutaneous block: a description of a novel technique. *Reg Anesth Pain Med.* 2005;30(2):198-201.
63. Moore KL, Agur AMR. *Essential Clinical Anatomy.* Baltimore, MD: Williams & Wilkins; 1995.
64. The New York School of Regional Anesthesia. <http://www.nysora.com>. Accessed on June 15, 2010.
65. Marhofer P, Nasel C, Sitzwohl C, Kapral S. Magnetic resonance imaging of the distribution of local anesthetic during the three-in-one block. *Anesth Analg.* 2000;90:119.

66. Capdevila X, Macaire P, Dadure C, et al. Continuous psoas compartment block for postoperative analgesia after total hip arthroplasty: new landmarks, technical guidelines, and clinical evaluation. *Anesth Analg*. 2002; 94:1606-1613.
67. Labat G. *Regional Anesthesia: Its Technic and Clinical Application*. Philadelphia, PA: WB Saunders; 1924.
68. O'Donnell BD, Mannion S. Ultrasound-guided femoral nerve block, the safest way to proceed? *Reg Anesth Pain Med*. 2006;31:387-388.
69. Vloka JD, Hadzic A, Reiss W, et al. Femoral nerve block: needle insertion at the inguinal crease results in more consistent nerve localization. *Reg Anesth*. 1998;23:53.
70. van der Wal M, Lang SA, Yip RW. Transsartorial approach for saphenous nerve block. *Can J Anaesth*. 1993;40:542-546.
71. Muraskin SI, Conrad B, Zheng N, Morey TE, Enneking FK. Falls associated with lower-extremity-nerve blocks: a pilot investigation of mechanisms. *Reg Anesth Pain Med*. 2007;32:67-72.
72. Gomez D, Sidhu S, Jha K. Femoral and saphenous nerve palsy post-total knee replacement. *Internet J Orthop Surg*. 2006;3(1).
73. Chelly JE, Greger J, Gebhard R, et al. Continuous femoral blocks improve recovery and outcome of patients undergoing total knee arthroplasty. *J Arthroplasty*. 2001;16:436-445.
74. Wardrop PJ, Nishikawa H. Lateral cutaneous nerve of the thigh blockade as primary anaesthesia for harvesting skin grafts. *Br J Plast Surg*. 1995;48:597-600.
75. Shannon J, Lang SA, Yip RW, Gerard M. Lateral femoral cutaneous nerve block revisited. A nerve stimulator technique. *Reg Anesth*. 1995;20(2):100-104.
76. Bouaziz H, Vial F, Jochum D, et al. An evaluation of the cutaneous distribution after obturator nerve block. *Anesth Analg*. 2002;94:445-449.
77. Wassef MR. Interadductor approach to obturator nerve blockade for spastic conditions of adductor thigh muscles. *Reg Anesth*. 1993;18:13.
78. Di Benedetto P, Casati A, Bertini L, Fanelli G. Posterior subgluteal approach to block the sciatic nerve: description of the technique and initial clinical experiences. *Eur J Anaesthesiol*. 2002;19:682-686.
79. Chan VW, Nova H, Abbas S, McCartney CJ, Perlas A, Xu DQ. Ultrasound examination and localization of the sciatic nerve: a volunteer study. *Anesthesiology*. 2006;104:309-314.
80. Chantzi C, Alevizou A, Saranteas T, Zogogiannis J, Iatrou C, Dimitriou V. Usefulness of the two to 5 MHz ultrasound probe in examination and block of the sciatic nerve in orthopedic trauma patients: a preliminary study. *J Clin Anesth*. 2007;19:486-488.
81. Sukhani R, Candido KD, Doty R, Yaghmour E, McCarthy RJ. Infragluteal-parabiceps sciatic nerve block: an evaluation of a novel approach using a single-injection technique. *Anesth Analg*. 2003;96:868-873.
82. Vloka JD, Hadzi A, April E, Thys DM. The division of the sciatic nerve in the popliteal fossa: anatomical implications for popliteal nerve blockade. *Anesth Analg*. 2001;92:215.
83. Ichiyanagi K. Sciatic nerve block: lateral approach with the patient supine. *Anesthesiology*. 1959;20:601.
84. Paqueron X, Bouaziz H, Macalou D, et al. The lateral approach to the sciatic nerve at the popliteal fossa: one or two injections? *Anesth Analg*. 1999;89:1221.
85. Redborg KE, Sites BD, Chinn CD, et al. Ultrasound improves the success rate of a sural nerve block at the ankle. *Reg Anesth Pain Med*. 2009;34:24-28.
86. Choi YK, Liu J. The use of 5% of lidocaine for prolonged analgesia in chronic pain patients: a new technique. *Reg Anesth Pain Med*. 1998;23:96-100.
87. Hadzić A, Vloka JD, Saff GN, Hertz R, Thys DM. The "three-in-one block" for treatment of pain in a patient with acute herpes zoster infection. *Reg Anesth*. 1997;22:575-578.