

Chapter 50

REFERENCES

1. Koller C. The use of cocaine for producing anaesthesia of the eye [translated and reprinted]. *Lancet*. 1884;II:990-992.
2. Atlee JL. *Complications in Anesthesia*. 2nd ed. Philadelphia, PA: Elsevier/Saunders; 2007.
3. Ben-David B. Complications of regional anesthesia: an overview. *Anesthesiol Clin North Am*. 2002;20:665-667, ix.
4. Kaye RD, Heard CMB. Alleged malpractice. In: Atlee JL, ed. *Complications in Anesthesia*. 2nd ed. Philadelphia, PA: Elsevier/Saunders; 2007:933-935.
5. Neal JM, Bernards CM, Hadzic A, et al. ASRA practice advisory on neurologic complications in regional anesthesia and pain medicine. *Reg Anesth Pain Med*. 2008;33:404-415.
6. Brull R, Wijayatilake DS, Perlas A, et al. Practice patterns related to block selection, nerve localization and risk disclosure: a survey of the American Society of Regional Anesthesia and Pain Medicine. *Reg Anesth Pain Med*. 2008;33:395-403.
7. Waisel DB, Truog RD. Informed consent. *Anesthesiology*. 1997;87:968-978.
8. Raj PP. Peripheral nerve stimulators for nerve blocks. In: Raj PP, de Andres J, Grossi P, Banister R, Sala-Blanc X, eds. *Textbook of Regional Anesthesia*. New York, NY: Churchill Livingstone; 2002:251-268.
9. Ford DJ, Pither CE, Raj PP. Electrical characteristics of peripheral nerve stimulators: implication for nerve localization. *Reg Anesth*. 1984;9:73-77.
10. Tsui BCH, Wagner A, Finucane B. Electrophysiologic effect of injectates on peripheral nerve stimulation. *Reg Anesth Pain Med*. 2004;29:189-193.
11. Ercole A. The effect of injectate conductivity on the electric field with the nerve stimulator needle: a computer simulation. *Anesth Analg*. 2008;107:1427-1432.
12. Hadzic A. Peripheral nerve stimulators: cracking the code—one at a time. *Reg Anesth Pain Med*. 2004;29:185-188.
13. Pham Dang C. Stimulating catheter and electrophysiologic effect of the flush injectates: exciting. *Reg Anesth Pain Med*. 2004;29:509-510.
14. Tsui BCH, Kropelin B. The electrophysiological effect of dextrose 5% in water on single-shot peripheral nerve stimulation. *Anesth Analg*. 2005;100:1837-1839.
15. Pham-Dang C, Kick O, Collet T, Gouin F, Pinaud M. Continuous peripheral nerve blocks with stimulating catheters. *Reg Anesth Pain Med*. 2003;28:83-88.
16. Pham Dang C, Lelong A, Guilley J, et al. Effect on neurostimulation of injectates used for perineural space expansion before placement of a stimulating catheter: normal saline versus dextrose 5% in water. *Reg Anesth Pain Med*. 2009;34:398-403.
17. Tsui BCH, Finucane B. Practical recommendations for improving needle-tip visibility under ultrasound guidance? *Reg Anesth Pain Med*. 2005;30:596-597.
18. Habicher M, Ocken M, Birnbaum J, Volk T. [Electrical nerve stimulation for peripheral nerve blocks. Ultrasound-guided needle positioning and effect of 5% glucose injection]. *Anaesthesist*. 2009;58:986-991.
19. Tsui BC, Gupta S, Finucane B. Confirmation of epidural catheter placement using nerve stimulation. *Can J Anaesth*. 1998;45:640-644.
20. Tsui BC, Guenther C, Emery D, Finucane B. Determining epidural catheter location using nerve stimulation with radiological confirmation. *Reg Anesth Pain Med*. 2000;25:306-309.
21. Moore AR, Siddiqui N, Kassel EE, Carvalho JC. Unintentional subdural catheter placement during labor analgesia shows typical radiological pattern but atypical response to the Tsui test. *Int J Obstet Anesth*. 2010;19:111-114.
22. Tsui BC, Gupta S, Finucane B. Detection of subarachnoid and intravascular epidural catheter placement. *Can J Anaesth*. 1999;46:675-678.
23. Tsui BC, Gupta S, Emery D, Finucane B. Detection of subdural placement of epidural catheter using nerve stimulation. *Can J Anaesth*. 2000;47:471-473.
24. Lena P, Martin R. Subdural placement of an epidural catheter detected by nerve stimulation. *Can J Anaesth*. 2005;52:618-621.
25. Hoppenstein R. Percutaneous implantation of chronic spinal cord electrodes for control of intractable pain: preliminary report. *Surg Neurol*. 1975;4:195-198.
26. Krainick JU, Thoden U, Riechert T. Spinal cord stimulation in post-amputation pain. *Surg Neurol*. 1975;4:167-170.
27. North RB, Kidd DH, Zahurak M, James CS, Long DM. Spinal cord stimulation for chronic, intractable pain: experience over two decades. *Neurosurgery*. 1993;32:384-394.
28. Richardson RR, Nunez C, Siqueira EB. Histological reaction to percutaneous epidural neurostimulation: initial and long-term results. *Med Prog Technol*. 1979;6:179-184.

29. Tsui BC, Seal R, Koller J, et al. Thoracic epidural analgesia via the caudal approach in pediatric patients undergoing fundoplication using nerve stimulation guidance. *Anesth Analg*. 2001;93:1152-1155.
30. Tsui BCH, Wagner A, Cave D, Kearney R. Thoracic and lumbar epidural analgesia via the caudal approach using electrical stimulation guidance in pediatric patients: a review of 289 patients. *Anesthesiology*. 2004;100:683-689.
31. Ting PL, Sivagnanratnam V. Ultrasonographic study of the spread of local anesthetic during axillary brachial plexus block. *Br J Anaesth*. 1989;63:326-329.
32. Perlas A, Lobo G, Lo N, et al. Ultrasound-guided supraclavicular block: outcome of 510 consecutive cases. *Reg Anesth Pain Med*. 2009;34:171-176.
33. Klaastad O, Sauter AR, Dodgson MS. Brachial plexus block with or without ultrasound guidance. *Curr Opin Anaesthesiol*. 2009;22:655-660.
34. Sites BD, Chan VW, Neal JM, et al. The American Society of Regional Anesthesia and Pain Medicine and the European Society of Regional Anaesthesia and Pain Therapy joint committee recommendations for education and training in ultrasound-guided regional anesthesia. *Reg Anesth Pain Med*. 2010;35:S74-S80.
35. Koprak S, Krafft P, Eibenberger K, et al. Ultrasound-guided supraclavicular approach for regional anesthesia of the brachial plexus. *Anesth Analg*. 1994;78:503-507.
36. Williams SR, Chouinard P, Arcand G, et al. Ultrasound guidance speeds execution and improves the quality of supraclavicular block. *Anesth Analg*. 2003;97:1518-1523.
37. Griffin J, Nicholls B. Ultrasound in regional anaesthesia. *Anaesthesia* 2010;65(Suppl 1):1-12.
38. Orebaugh SL, Williams BA, Kentor ML. Ultrasound guidance with nerve stimulation reduces the time necessary for resident peripheral nerve blockade. *Reg Anesth Pain Med*. 2007;32:448-454.
39. Lo N, Brull R, Perlas A, et al. Evolution of ultrasound guided axillary brachial plexus blockade: retrospective analysis of 662 blocks. *Can J Anaesth*. 2008;55:408-413.
40. Casati A, Baciarello M, Di Cianni S, et al. Effects of ultrasound guidance on the minimum effective anaesthetic volume required to block the femoral nerve. *Br J Anaesth*. 2007;98:823-827.
41. Willschke H, Bosenberg A, Marhofer P, et al. Ultrasonographic-guided ilioinguinal/iliohypogastric nerve block in pediatric anesthesia: what is the optimal volume? *Anesth Analg*. 2006;102:1680-1684.
42. Warman P, Nicholls B. Ultrasound-guided nerve blocks: efficacy and safety. *Best Pract Res Clin Anaesthesiol*. 2009;23:313-326.
43. Neal JM. Ultrasound-guided regional anesthesia and patient safety: an evidence-based analysis. *Reg Anesth Pain Med*. 2010;35:S59-S67.
44. Chan VWS, Perlas A, Rawson R, Odukoya O. Ultrasound-guided supraclavicular brachial plexus block. *Anesth Analg*. 2003;97:1514-1517.
45. Chan VW. Applying ultrasound imaging to interscalene brachial plexus block. *Reg Anesth Pain Med*. 2003;28:340-343.
46. Tsui B, Suresh S. Ultrasound imaging for regional anesthesia in infants, children, and adolescents: a review of current literature and its application in the practice of extremity and trunk blocks. *Anesthesiology*. 2010;112:473-492.
47. Grau T, Leipold RW, Conradi R, Martin E, Motsch J. Efficacy of ultrasound imaging in obstetric epidural anesthesia. *J Clin Anesth*. 2002;14:169-175.
48. Grau T, Leipold RW, Fatehi S, Martin E, Motsch J. Real-time ultrasonic observation of combined spinal-epidural anaesthesia. *Eur J Anaesthesiol*. 2004;21:25-31.
49. Grau T, Bartussek E, Conradi R, Martin E, Motsch J. Ultrasound imaging improves learning curves in obstetric epidural anesthesia: a preliminary study. *Can J Anaesth*. 2003;50:1047-1050.
50. Grau T, Leipold RW, Horter J, et al. Paramedian access to the epidural space: the optimum window for ultrasound imaging. *J Clin Anesth*. 2001;13:213-217.
51. Grau T, Leipold RW, Conradi R, Martin E. Ultrasound control for presumed difficult epidural puncture. *Acta Anaesthesiol Scand*. 2001;45:766-771.
52. Chawathe MS, Jones RM, Gildersleve CD, et al. Detection of epidural catheters with ultrasound in children. *Paediatr Anaesth*. 2003;13:681-684.
53. Chen CPC, Tang SFT, Hsu T-C, et al. Ultrasound guidance in caudal epidural needle placement. *Anesthesiology*. 2004;101:181-184.
54. Shevde K, Panagopoulos G. A survey of 800 patients' knowledge, attitudes and concerns regarding anesthesia. *Anesth Analg*. 1991;73:190-198.
55. Haugen AS, Eide GE, Olsen MV, et al. Anxiety in the operating theatre: a study of frequency and environmental impact in patients having local, plexus or regional anaesthesia. *J Clin Nurs*. 2009;18:2301-2310.
56. Curley RK, Macfarlane AW, King CM. Contact sensitivity to the amide anesthetics lidocaine, prilocaine, and mepivacaine. Case report and review of the literature. *Arch Dermatol*. 1986;122:924-926.
57. Nazir MS, Holdcroft A. Local anaesthetic drugs: adverse effects as reported through the ADROIT system in the UK. *Pharmacoepidemiol Drug Saf*. 2009;18:1000-1006.
58. Dawkins CJ. An analysis of the complications of extradural and caudal block. *Anaesthesia*. 1969;24:554-563.
59. Auroy Y, Narchi P, Messiah A, et al. Serious complications related to regional anesthesia: results of a prospective survey in France. *Anesthesiology*. 1997;87:479-486.
60. Brown DT, Beamish D, Wildsmith JA. Allergic reaction to an amide local anaesthetic. *Br J Anaesth* 1981;53:435-437.
61. Covino BG, Vassallo HG. Pharmacokinetic aspects of local anesthetic agents. *Local anesthetics: mechanisms of action and clinical use*. New York, NY: Grune & Stratton; 1976: 95-123.
62. Lofstrom JB. Physiologic disposition of local anesthetics. *Reg Anesth*. 1982;7:33-38.
63. Englesson S, Grevsten S. The influence of acid-base changes on central nervous system toxicity of local anaesthetic agents. II. *Acta Anaesthesiol Scand*. 1974;18:88-8103.
64. Mather LE, Cousins MJ. Local anaesthetics and their current clinical use. *Drugs*. 1979;18:185-205.
65. Weinberg GL. Current concepts in resuscitation of patients with local anesthetic cardiac toxicity. *Reg Anesth Pain Med*. 2002;27:568-575.
66. Tanaka M, Kimura T, Goyagi T, et al. Evaluating hemodynamic and T wave criteria of simulated intravascular test doses using bupivacaine or isoproterenol in anesthetized children. *Anesth Analg*. 2000;91:567-572.

67. Tanaka M, Nishikawa T. A comparative study of hemodynamic and T-wave criteria for detecting intravascular injection of the test dose (epinephrine) in sevoflurane-anesthetized adults. *Anesth Analg*. 1999;89:32-36.
68. Tanaka M, Sato M, Kimura T, Nishikawa T. The efficacy of simulated intravascular test dose in sedated patients. *Anesth Analg*. 2001;93:1612-1617.
69. Brull R, Perlas A, Cheng PH, Chan VW. Minimizing the risk of intravascular injection during ultrasound-guided peripheral nerve blockade. *Anesthesiology*. 2008;109:1142; author reply 1144, 1144-1145.
70. Orebaugh SL, Williams BA, Vallejo M, Kentor ML. Adverse outcomes associated with stimulator-based peripheral nerve blocks with versus without ultrasound visualization. *Reg Anesth Pain Med*. 2009;34:251-255.
71. Chin J, Tsui BC. No change in impedance upon intravascular injection of D5W. *Can J Anaesth*. 2010;57:559-564.
72. Faccenda KA, Finucane BT. Complications of regional anaesthesia: Incidence and prevention. *Drug Saf*. 2001;24:413-442.
73. Heavner JE, Dryden CF, Sanghani V, et al. Severe hypoxia enhances central nervous system and cardiovascular toxicity of bupivacaine in lightly anesthetized pigs. *Anesthesiology*. 1992;77:142-147.
74. Rosen MA, Thigpen JW, Shnider SM, et al. Bupivacaine-induced cardiotoxicity in hypoxic and acidotic sheep. *Anesth Analg*. 1985;64:1089-1096.
75. Weinberg G, Ripper R, Feinstein DL, Hoffman W. Lipid emulsion infusion rescues dogs from bupivacaine-induced cardiac toxicity. *Reg Anesth Pain Med*. 2003;28:198-202.
76. Weinberg GL, VadeBoncouer T, Ramaraju GA, Garcia-Amaro MF, Cwik MJ. Pretreatment or resuscitation with a lipid infusion shifts the dose-response to bupivacaine-induced asystole in rats. *Anesthesiology*. 1998;88:1071-1075.
77. Weinberg G. Lipid rescue resuscitation from local anaesthetic cardiac toxicity. *Toxicol Rev*. 2006;25:139-145.
78. Litz RJ, Roessel T, Heller AR, Stehr SN. Reversal of central nervous system and cardiac toxicity after local anesthetic intoxication by lipid emulsion injection. *Anesth Analg*. 2008;106:1575-1577.
79. Picard J, Meek T. Lipid emulsion to treat overdose of local anaesthetic: the gift of the glob. *Anaesthesia*. 2006;61:107-109.
80. Groban L, Butterworth J. Lipid reversal of bupivacaine toxicity: has the silver bullet been identified? *Reg Anesth Pain Med*. 2003;28:167-169.
81. Rosenblatt MA, Abel M, Fischer GW, Itzkovich CJ, Eisenkraft JB. Successful use of a 20% lipid emulsion to resuscitate a patient after a presumed bupivacaine-related cardiac arrest. *Anesthesiology*. 2006;105:217-218.
82. Litz RJ, Popp M, Stehr SN, Koch T. Successful resuscitation of a patient with ropivacaine-induced asystole after axillary plexus block using lipid infusion. *Anaesthesia*. 2006;61:800-801.
83. Leskiw U, Weinberg GL. Lipid resuscitation for local anesthetic toxicity: is it really lifesaving? *Curr Opin Anaesthesiol*. 2009;22:667-671.
84. Foxall G, McCahon R, Lamb J, Hardman JG, Bedforth NM. Levobupivacaine-induced seizures and cardiovascular collapse treated with Intralipid. *Anaesthesia*. 2007;62:516-518.
85. Ludot H, Tharin JY, Belouadah M, Mazoit JX, Malinovsky JM. Successful resuscitation after ropivacaine and lidocaine-induced ventricular arrhythmia following posterior lumbar plexus block in a child. *Anesth Analg*. 2008;106:1572-1574.
86. Marwick PC, Levin AI, Coetzee AR. Recurrence of cardiotoxicity after lipid rescue from bupivacaine-induced cardiac arrest. *Anesth Analg*. 2009;108:1344-1346.
87. Neal JM, Bernards CM, Butterworth JFt, et al. ASRA practice advisory on local anesthetic systemic toxicity. *Reg Anesth Pain Med*. 2010;35:152-161.
88. Heavner JE, Pitkanen MT, Shi B, Rosenberg PH. Resuscitation from bupivacaine-induced asystole in rats: comparison of different cardioactive drugs. *Anesth Analg*. 1995;80:1134-1139.
89. Hiller DB, Gregorio GD, Ripper R, et al. Epinephrine impairs lipid resuscitation from bupivacaine overdose: a threshold effect. *Anesthesiology*. 2009;111:498-505.
90. Krismer AC, Hogan QH, Wenzel V, et al. The efficacy of epinephrine or vasopressin for resuscitation during epidural anesthesia. *Anesth Analg*. 2001;93:734-742.
91. Mayr VD, Raedler C, Wenzel V, Lindner KH, Strohmenger H-U. A comparison of epinephrine and vasopressin in a porcine model of cardiac arrest after rapid intravenous injection of bupivacaine. *Anesth Analg*. 2004;98:1426-1431.
92. Neustein S, Sampson I, Dimich I, Shiang H, Tatu J. Milrinone is superior to epinephrine as treatment of myocardial depression due to ropivacaine in pigs. *Can J Anaesth*. 2000;47:1114-1118.
93. Long WB, Rosenblum S, Grady IP. Successful resuscitation of bupivacaine-induced cardiac arrest using cardiopulmonary bypass. *Anesth Analg*. 1989;69:403-406.
94. Barrington MJ, Watts SA, Gledhill SR, et al. Preliminary results of the Australasian Regional Anaesthesia Collaboration: a prospective audit of more than 7000 peripheral nerve and plexus blocks for neurologic and other complications. *Reg Anesth Pain Med*. 2009;34:534-541.
95. Sorenson EJ. Neurological injuries associated with regional anesthesia. *Reg Anesth Pain Med*. 2008;33:442-448.
96. Renck H. Neurological complications of central nerve blocks. *Acta Anaesthesiol Scand*. 1995;39:859-868.
97. Deschner S, Borgeat A, Hadzic A. Neurologic complications of peripheral nerve blocks: mechanisms and management. In: Hadzic A, ed. *Textbook of Regional Anesthesia and Acute Pain Management*. New York, NY: McGraw-Hill; 2007:969-978.
98. Moore DC, Mulroy MF, Thompson GE. Peripheral nerve damage and regional anaesthesia. *Br J Anaesth*. 1994;73:435-436.
99. Bernards CM, Hadzic A, Suresh S, Neal JM. Regional anesthesia in anesthetized or heavily sedated patients. *Reg Anesth Pain Med*. 2008;33:449-460.
100. Bigeleisen PE. Nerve puncture and apparent intraneural injection during ultrasound-guided axillary block does not invariably result in neurologic injury. *Anesthesiology*. 2006;105:779-783.
101. Robards C, Hadzic A, Somasundaram L, et al. Intraneural injection with low-current stimulation during popliteal sciatic nerve block. *Anesth Analg*. 2009;109:673-677.
102. Sala Blanch X, Lopez AM, Carazo J, et al. Intraneural injection during nerve stimulator-guided sciatic nerve block at the popliteal fossa. *Br J Anaesth*. 2009;102:855-861.
103. Bigeleisen PE, Moayeri N, Groen GJ. Extraneural versus intraneural stimulation thresholds during ultrasound-guided supraclavicular block. *Anesthesiology*. 2009;110:1235-1243.
104. Altermatt FR, Cummings TJ, Auten KM, et al. Ultrasonographic appearance of intraneural injections in the porcine model. *Reg Anesth Pain Med*. 2010;35:203-206.

105. Lupu CM, Kiehl TR, Chan VW, et al. Nerve expansion seen on ultrasound predicts histologic but not functional nerve injury after intraneural injection in pigs. *Reg Anesth Pain Med*. 2010;35:132-139.
106. Selander D, Dhuner KG, Lundborg G. Peripheral nerve injury due to injection needles used for regional anesthesia. An experimental study of the acute effects of needle point trauma. *Acta Anaesthesiol Scand*. 1977;21:182-188.
107. Neal JM, Hebl JR. Complications after peripheral nerve block. In: Benson H, et al, eds. *Essentials of Pain Medicine and Regional Anesthesia*. 2nd ed. New York, NY: Churchill Livingstone; 2004:694-701.
108. Hadzic A, Dilberovic F, Shah S, et al. Combination of intraneural injection and high injection pressure leads to fascicular injury and neurologic deficits in dogs. *Reg Anesth Pain Med*. 2004;29:417-423.
109. Claudio R, Hadzic A, Shih H, et al. Injection pressures by anesthesiologists during simulated peripheral nerve block. *Reg Anesth Pain Med*. 2004;29:201-205.
110. Tsui BC, Li LX, Pillay JJ. Compressed air injection technique to standardize block injection pressures. *Can J Anaesth*. 2006;53:1098-1102.
111. Weaver MA, Tandatnick CA, Hahn MB. Peripheral nerve blockade. In: Raj PP, de Andres J, Grossi P, Banister R, Sala-Blanc X, eds. *Textbook of Regional Anesthesia*. New York, NY: Churchill Livingstone; 2002:857-870.
112. Finucane BT. *Complications of Regional Anesthesia*. 2nd ed. New York, NY: Springer; 2007.
113. Mulroy MF. *Regional Anesthesia: An Illustrated Procedural Guide*. 2nd ed. Boston, MA: Little, Brown; 1996.
114. Munson ES, Paul WL, Embro WJ. Central-nervous-system toxicity of local anesthetic mixtures in monkeys. *Anesthesiology*. 1977;46:179-183.
115. Borgeat A, Ekotodramis G, Kalberer F, Benz C. Acute and nonacute complications associated with interscalene block and shoulder surgery: a prospective study. *Anesthesiology*. 2001;95:875-880.
116. Urban MK, Urquhart B. Evaluation of brachial plexus anesthesia for upper extremity surgery. *Reg Anesth*. 1994;19:175-182.
117. Barutell C, Vidal F, Raich M, Montero A. A neurological complication following interscalene brachial plexus block. *Anaesthesia*. 1980;35:365-367.
118. Passannante AN. Spinal anesthesia and permanent neurologic deficit after interscalene block. *Anesth Analg*. 1996;82:873-874.
119. Winnie AP. *Plexus Anesthesia: I. Perivascular Techniques of Brachial Plexus Block*. Philadelphia, PA: WB Saunders; 1983.
120. Kulenkampff D. Anaesthesia of the brachial plexus [German]. *Zentralb Chir*. 1911;40:849.
121. Brand L, Papper EM. A comparison of supraclavicular and axillary techniques for brachial plexus blocks. *Anesthesiology*. 1961;22:226-229.
122. De Jong RH. *Local Anesthetics*. 2nd ed. Springfield, IL: Thomas; 1977.
123. Winnie AP. Interscalene brachial plexus block. *Anesth Analg*. 1970;49:455-466.
124. Ward ME. The interscalene approach to the brachial plexus. *Anaesthesia*. 1974;29:147-157.
125. Brown DL, Bridenbaugh LD. The upper extremity: somatic block. In: Cousins MJ, Bridenbaugh PO, eds. *Neural Blockade in Clinical Anesthesia and Management of Pain*. 3rd ed. Philadelphia, PA: Lippincott-Raven; 1998:345-371.
126. Tsui BC, Doyle K, Chu K, Pillay J, Dillane D. Case series: ultrasound-guided supraclavicular block using a curvilinear probe in 104 day-case hand surgery patients. *Can J Anaesth*. 2009;56:46-51.
127. Lichtenstein DA. Pneumothorax and introduction to ultrasound signs in the lung. In: Lichtenstein MA, ed. *General Ultrasound in the Critically Ill*. Berlin, Germany: Springer; 2007:105-114.
128. Lambert LA, Lambert DH, Strichartz GR. Irreversible conduction block in isolated nerve by high concentrations of local anesthetics. *Anesthesiology*. 1994;80:1082-1093.
129. Ravindran RS, Bond VK, Tasch MD, Gupta CD, Luerssen TG. Prolonged neural blockade following regional analgesia with 2-chloroprocaine. *Anesth Analg*. 1980;59:447-451.
130. Stevens RA, Urmey WF, Urquhart BL, Kao TC. Back pain after epidural anesthesia with chloroprocaine. *Anesthesiology*. 1993;78:492-497.
131. Fibuch EE, Opper SE. Back pain following epidurally administered Nesacaine-MPF. *Anesth Analg*. 1989;69:113-115.
132. Schneider M, Ettlin T, Kaufmann M, et al. Transient neurologic toxicity after hyperbaric subarachnoid anesthesia with 5% lidocaine. *Anesth Analg*. 1993;76:1154-1157.
133. Benoit PW, Belt WD. Some effects of local anesthetic agents on skeletal muscle. *Exp Neurol* 1972;34:264-278.
134. Benoit PW. Microscarring in skeletal muscle after repeated exposures to lidocaine with epinephrine. *J Oral Surg*. 1978;36:530-533.
135. Benoit PW. Reversible skeletal muscle damage after administration of local anesthetics with and without epinephrine. *J Oral Surg*. 1978;36:198-201.
136. Kytta J, Heinon G, Rosenberg PH. Effects of repeated bupivacaine administration on sciatic nerve and surrounding tissue in rats. *Acta Anaesthesiol Scand*. 1986;30:625-629.
137. Zink W, Bohl JRE, Hacke N, et al. The long term myotoxic effects of bupivacaine and ropivacaine after continuous peripheral nerve blocks. *Anesth Analg*. 2005;101:548-554.
138. Neal JM, Moore JM, Kopacz DJ, et al. Quantitative analysis of respiratory, motor, and sensory function after supraclavicular block. *Anesth Analg*. 1998;86:1239-1244.
139. Hood J, Knoblanche G. Respiratory failure following brachial plexus block. *Anaesth Intensive Care*. 1979;7:285-286.
140. Kayerker UM, Dick MM. Phrenic nerve paralysis following interscalene brachial plexus block. *Anesth Analg*. 1983;62:536-537.
141. Rau RH, Chan YL, Chuang HI, et al. Dyspnea resulting from phrenic nerve paralysis after interscalene brachial plexus block in an obese male—a case report. *Acta Anaesthesiol Sin*. 1997;35:113-118.
142. Bashein G, Robertson HT, Kennedy WF. Persistent phrenic nerve paresis following interscalene brachial plexus block. *Anesthesiology*. 1985;63:102-104.
143. Robaux S, Bouaziz H, Boisseau N, Raucoules-Aime M, Laxenaire MC. Persistent phrenic nerve paralysis following interscalene brachial plexus block. *Anesthesiology*. 2001;95:1519-1521.
144. Urmey WF, Talts KH, Sharrock NE. One hundred percent incidence of hemidiaphragmatic paresis associated with interscalene brachial plexus anesthesia as diagnosed by ultrasonography. *Anesth Analg*. 1991;72:498-503.

145. Farrar MD, Scheybani M, Nolte H. Upper extremity block effectiveness and complications. *Reg Anesth*. 1981;6:133-134.
146. Urmev WF, McDonald M. Hemidiaphragmatic paresis during interscalene brachial plexus block: effects on pulmonary function and chest wall mechanics. *Anesth Analg*. 1992;74:352-357.
147. Renes SH, Rettig HC, Gielen MJ, Wilder-Smith OH, van Geffen GJ. Ultrasound-guided low-dose interscalene brachial plexus block reduces the incidence of hemidiaphragmatic paresis. *Reg Anesth Pain Med*. 2009;34:498-502.
148. Renes SH, Spormans HH, Gielen MJ, Rettig HC, van Geffen GJ. Hemidiaphragmatic paresis can be avoided in ultrasound-guided supraclavicular brachial plexus block. *Reg Anesth Pain Med*. 2009;34:595-599.
149. Al-Khafaji JM, Elias MA. Incidence of Horner syndrome with interscalene brachial plexus block and its importance in the management of head injury. *Anesthesiology*. 1986;64:127-127.
150. Gray AT. Ultrasound-guided regional anesthesia: current state of the art. *Anesthesiology*. 2006;104:368-373, discussion 365A.
151. Neal JM, Brull R, Chan VW, et al. The ASRA evidence-based medicine assessment of ultrasound-guided regional anesthesia and pain medicine: executive summary. *Reg Anesth Pain Med*. 2010;35:S1-9.
152. Wong CA. Neurologic deficits and labor analgesia. *Reg Anesth Pain Med*. 2004;29:341-351.
153. Kao M-C, Tsai S-K, Tsou M-Y, et al. Paraplegia after delayed detection of inadvertent spinal cord injury during thoracic epidural catheterization in an anesthetized elderly patient. *Anesth Analg*. 2004;99:580-583.
154. Rose JB. Spinal cord injury in a child after single-shot epidural anesthesia. *Anesth Analg*. 2003;96:3-6.
155. Bromage PR, Benumof JL. Paraplegia following intracord injection during attempted epidural anesthesia under general anesthesia. *Reg Anesth Pain Med*. 1998;23:104-107.
156. Broadbent CR, Maxwell WB, Ferrie R, et al. Ability of anaesthetists to identify a marked lumbar interspace. *Anaesthesia*. 2000;55:1122-1126.
157. Pysyk CL, Persaud D, Bryson GL, Lui A. Ultrasound assessment of the vertebral level of the palpated intercrystal (Tuffier's) line. *Can J Anaesth*. 2010;57:46-49.
158. Hogan QH. Tuffier's line: the normal distribution of anatomic parameters. *Anesth Analg*. 1994;78:194-195.
159. Kim JT, Bahk JH, Sung J. Influence of age and sex on the position of the conus medullaris and Tuffier's line in adults. *Anesthesiology*. 2003;99:1359-1363.
160. Reynolds F. Damage to the conus medullaris following spinal anaesthesia. *Anaesthesia*. 2001;56:238-247.
161. Margarido CB, Arzola C, Balki M, Carvalho JC. Anesthesiologists' learning curves for ultrasound assessment of the lumbar spine. *Can J Anaesth*. 2010;57:120-126.
162. Hamandi K, Mottershead J, Lewis T, Ormerod IC, Ferguson IT. Irreversible damage to the spinal cord following spinal anesthesia. *Neurology*. 2002;59:624-626.
163. Simon SL, Abrahams JM, Sean Grady M, LeRoux PD, Rushton SA. Intramedullary injection of contrast into the cervical spinal cord during cervical myelography: a case report. *Spine (Phila Pa 1976)*. 2002;27:E274-E277.
164. Kandel ER, Schwartz JH, Jones RM. The perception of pain. In: Kandel ER, Schwartz JH, Jessell TM, eds. *Principles of Neural Science*. 4th ed. New York, NY: McGraw-Hill; 2000: 472-491.
165. Jackson MB, Pounder D, Price C, Matthews AW, Neville E. Percutaneous cervical cordotomy for the control of pain in patients with pleural mesothelioma. *Thorax*. 1999;54: 238-241.
166. Lahuerta J, Bowsher D, Lipton S, Buxton PH. Percutaneous cervical cordotomy: a review of 181 operations on 146 patients with a study on the location of "pain fibers" in the C-2 spinal cord segment of 29 cases. *J Neurosurg*. 1994;80:975-985.
167. Pounder D, Elliott S. An awake patient may not detect spinal cord puncture. *Anaesthesia*. 2000;55:194-194.
168. Tripathi M, Nath SS, Gupta RK. Paraplegia after intracord injection during attempted epidural steroid injection in an awake-patient. *Anesth Analg*. 2005;101:1209-1211.
169. Tsui BCH, Armstrong K. Can direct spinal cord injury occur without paresthesia? A report of delayed spinal cord injury after epidural placement in an awake patient. *Anesth Analg*. 2005;101:1212-1214.
170. Tsui BCH, Emery D, Uwiera RRE, Finucane B. The use of electrical stimulation to monitor epidural needle advancement in a porcine model. *Anesth Analg*. 2005;100:1611-1613.
171. Tsui BCH, Wagner AM, Cunningham K, et al. Threshold current of an insulated needle in the intrathecal space in pediatric patients. *Anesth Analg*. 2005;100:662-665.
172. Usubiaga JE. Neurological complications following epidural anesthesia. *Int Anesthesiol Clin*. 1975;13:1-153.
173. Niemi G. Advantages and disadvantages of adrenaline in regional anaesthesia. *Best Pract Res Clin Anaesthesiol*. 2005; 19:229-245.
174. Kozody R, Palahniuk RJ, Wade JG, Cumming MO, Pucci WR. The effect of subarachnoid epinephrine and phenylephrine on spinal cord blood flow. *Can Anaesth Soc J*. 1984;31: 503-508.
175. Horlocker TT, Wedel DJ, Benzon H, et al. Regional anesthesia in the anticoagulated patient: defining the risks (the second ASRA Consensus Conference on Neuraxial Anesthesia and Anticoagulation). *Reg Anesth Pain Med*. 2003;28:172-197.
176. Vandermeulen EP, Van Aken H, Vermeylen J. Anticoagulants and spinal-epidural anesthesia. *Anesth Analg*. 1994;79: 1165-1177.
177. Perkins WJ, Davis DH, Huntoon MA, Horlocker TT. A retained Racz catheter fragment after epidural neurolysis: implications during magnetic resonance imaging. *Anesth Analg*. 2003;96:1717-1719.
178. Kane RE. Neurologic deficits following epidural or spinal anesthesia. *Anesth Analg*. 1981;60:150-161.
179. Moen V, Dahlgren N, Irestedt L. Severe neurological complications after central neuraxial blockades in Sweden 1990-1999. *Anesthesiology*. 2004;101:950-959.
180. Hughes SC. HIV and anesthesia. *Anesthesiol Clin North America*. 2004;22:379-404, v.
181. Verma A. Epidemiology and clinical features of HIV-1 associated neuropathies. *J Peripher Nerv Syst*. 2001;6:8-13.
182. Kost-Byerly S, Tobin JR, Greenberg RS, et al. Bacterial colonization and infection rate of continuous epidural catheters in children. *Anesth Analg*. 1998;86:712-716.
183. Bubeck J, Boos K, Krause H, Thies K-C. Subcutaneous tunneling of caudal catheters reduces the rate of bacterial colonization to that of lumbar epidural catheters. *Anesth Analg*. 2004; 99:689-693.
184. Baker AS, Ojemann RG, Swartz MN, Richardson EP. Spinal epidural abscess. *N Engl J Med*. 1975;293:463-468.

185. Eng RH, Seligman SJ. Lumbar puncture-induced meningitis. *JAMA*. 1981;245:1456-1459.
186. Feldman JM, Chapin-Robertson K, Turner J. Do agents used for epidural analgesia have antimicrobial properties? *Reg Anesth*. 1994;19:43-47.
187. Tsui BCH, Malherbe S, Koller J, Aronyk K. Reversal of an unintentional spinal anesthetic by cerebrospinal lavage. *Anesth Analg*. 2004;98:434-436.
188. Ben-David B, Rawa R. Complications of neuraxial blockade. *Anesthesiol Clin North Am*. 2002;20:669-693.
189. Ben-David B, Vaida S, Gaitini L. The influence of high spinal anesthesia on sensitivity to midazolam sedation. *Anesth Analg*. 1995;81:525-528.
190. Tverskoy M, Shagal M, Finger J, Kissin I. Spinal anesthesia and midazolam hypnotic requirements. *Anesth Analg*. 1996;83:198-199.
191. Tverskoy M, Shifrin V, Finger J, Fleyshman G, Kissin I. Effect of epidural bupivacaine block on midazolam hypnotic requirements. *Reg Anesth*. 1996;21:209-213.
192. Lubenow T, Keh-Wong E, Kristof K, Ivankovich O, Ivankovich AD. Inadvertent subdural injection: a complication of an epidural block. *Anesth Analg*. 1988;67:175-179.
193. Reynolds F, Speedy HM. The subdural space: the third place to go astray. *Anaesthesia*. 1990;45:120-123.
194. Mollmann M, Holst D, Enk D, et al. [Subdural intra-arachnoid spread of local anesthetics. A complication of spinal anesthesia]. *Anaesthesist*. 1992;41:685-688.
195. Wills JH. Rapid onset of massive subdural anesthesia. *Reg Anesth Pain Med*. 2005;30:299-302.
196. Brown DL, Ransom DM, Hall JA, et al. Regional anesthesia and local anesthetic-induced systemic toxicity: seizure frequency and accompanying cardiovascular changes. *Anesth Analg*. 1995;81:321-328.
197. Albright GA. Cardiac arrest following regional anesthesia with etidocaine or bupivacaine. *Anesthesiology*. 1979;51:285-287.
198. Reynolds F. Bupivacaine and intravenous regional anaesthesia. *Anaesthesia*. 1984;39:105-107.
199. Banwell BR, Morley-Forster P, Krause R. Decreased incidence of complications in parturients with the arrow (FlexTip Plus) epidural catheter. *Can J Anaesth*. 1998;45:370-372.
200. Tsui BC, Gupta S, Finucane B. Determination of epidural catheter placement using nerve stimulation in obstetric patients. *Reg Anesth Pain Med*. 1999;24:17-23.
201. Turnbull DK, Shepherd DB. Post-dural puncture headache: pathogenesis, prevention and treatment. *Br J Anaesth*. 2003;91:718-729.
202. Patin DJ, Eckstein EC, Harum K, Pallares VS. Anatomic and biomechanical properties of human lumbar dura mater. *Anesth Analg*. 1993;76:535-540.
203. Fink BR. Postspinal headache. *Anesth Analg*. 1990;71:208-209.
204. Norris MC, Leighton BL, DeSimone CA. Needle bevel direction and headache after inadvertent dural puncture. *Anesthesiology*. 1989;70:729-731.
205. Lybecker H, Moller JT, May O, Nielsen HK. Incidence and prediction of postdural puncture headache. A prospective study of 1021 spinal anesthetics. *Anesth Analg*. 1990;70:389-394.
206. Ready LB, Cuplin S, Haschke RH, Nessly M. Spinal needle determinants of rate of transdural fluid leak. *Anesth Analg*. 1989;69:457-460.
207. Baraz R, Collis RE. The management of accidental dural puncture during labour epidural analgesia: a survey of UK practice. *Anaesthesia*. 2005;60:673-679.
208. Kuczkowski KM, Benumof JL. Decrease in the incidence of post-dural puncture headache: maintaining CSF volume. *Acta Anaesthesiol Scand*. 2003;47:98-100.
209. Munnur U, Suresh MS. Backache, headache, and neurologic deficit after regional anesthesia. *Anesthesiol Clin North America*. 2003;21:71-86.
210. Halker RB, Demaerschalk BM, Wellik KE, et al. Caffeine for the prevention and treatment of postdural puncture headache: debunking the myth. *Neurologist*. 2007;13:323-327.
211. Camann WR, Murray RS, Mushlin PS, Lambert DH. Effects of oral caffeine on postdural puncture headache. A double-blind, placebo-controlled trial. *Anesth Analg*. 1990;70:181-184.
212. Bolton VE, Leicht CH, Scanlon TS. Postpartum seizure after epidural blood patch and intravenous caffeine sodium benzoate. *Anesthesiology*. 1989;70:146-149.
213. Carp H, Singh PJ, Vadhera R, Jayaram A. Effects of the serotonin-receptor agonist sumatriptan on postdural puncture headache: report of six cases. *Anesth Analg*. 1994;79:180-182.
214. Thew M, Paech MJ. Management of postdural puncture headache in the obstetric patient. *Curr Opin Anaesthesiol*. 2008;21:288-292.
215. Bussone G, Tullo V, d'Onofrio F, et al. Frovatriptan for the prevention of postdural puncture headache. *Cephalalgia*. 2007;27:809-813.
216. Kshatri AM, Foster PA. Adrenocorticotrophic hormone infusion as a novel treatment for postdural puncture headache. *Reg Anesth*. 1997;22:432-434.
217. Gormley JB. Treatment of post spinal headache. *Anesthesiology*. 1960;21:565.
218. DiGiovanni AJ, Dunbar BS. Epidural injections of autologous blood for postlumbar-puncture headache. *Anesth Analg*. 1970;49:268-271.
219. DiGiovanni AJ, Galbert MW, Wahle WM. Epidural injection of autologous blood for postlumbar-puncture headache. II. Additional clinical experiences and laboratory investigation. *Anesth Analg*. 1972;51:226-232.
220. Szeinfeld M, Ihmeidan IH, Moser MM, et al. Epidural blood patch: evaluation of the volume and spread of blood injected into the epidural space. *Anesthesiology*. 1986;64:820-822.
221. Beards SC, Jackson A, Griffiths AG, Horsman EL. Magnetic resonance imaging of extradural blood patches: appearances from 30 min to 18 h. *Br J Anaesth*. 1993;71:182-188.
222. Andrews PJ, Ackerman WE, Juneja M, Cases-Cristobal V, Rigor BM. Transient bradycardia associated with extradural blood patch after inadvertent dural puncture in parturients. *Br J Anaesth*. 1992;69:401-403.
223. Palmer JH, Wilson DW, Brown CM. Lumbovertebral syndrome after repeat extradural blood patch. *Br J Anaesth*. 1997;78:334-336.
224. Perez M, Olmos M, Garrido FJ. Facial nerve paralysis after epidural blood patch. *Reg Anesth*. 1993;18:196-198.
225. Seeberger MD, Urwyler A. Lumbovertebral syndrome after extradural blood patch. *Br J Anaesth*. 1992;69:414-416.
226. Tekkok IH, Carter DA, Brinker R. Spinal subdural haematoma as a complication of immediate epidural blood patch. *Can J Anaesth*. 1996;43:306-309.

227. Kowbel MA, Comfort VK. Caudal epidural blood patch for the treatment of a paediatric subarachnoid-cutaneous fistula. *Can J Anaesth.* 1995;42:625-627.
228. Roy L, Vischoff D, Lavoie J. Epidural blood patch in a seven-year-old child. *Can J Anaesth.* 1995;42:621-624.
229. Cheek TG, Banner R, Sauter J, Gutsche BB. Prophylactic extradural blood patch is effective. A preliminary communication. *Br J Anaesth.* 1988;61:340-342.
230. Palahniuk RJ, Cumming M. Prophylactic blood patch does not prevent post lumbar puncture headache. *Can Anaesth Soc J.* 1979;26:132-133.
231. Aldrete JA, Brown TL. Intrathecal hematoma and arachnoiditis after prophylactic blood patch through a catheter. *Anesth Analg.* 1997;84:233-234.
232. Boonmak P, Boonmak S. Epidural blood patching for preventing and treating post-dural puncture headache. *Cochrane Database Syst Rev.* 2010:CD001791.
233. Baysinger CL, Menk EJ, Harte E, Middaugh R. The successful treatment of dural puncture headache after failed epidural blood patch. *Anesth Analg.* 1986;65:1242-1244.
234. Stevens RA, Jorgensen N. Successful treatment of dural puncture headache with epidural saline infusion after failure of epidural blood patch. Case report. *Acta Anaesthesiol Scand.* 1988;32:429-431.
235. Gerritse BM, van Dongen RT, Crul BJ. Epidural fibrin glue injection stops persistent cerebrospinal fluid leak during long-term intrathecal catheterization. *Anesth Analg.* 1997;84:1140-1141.
236. Crul BJ, Gerritse BM, van Dongen RT, Schoonderwaldt HC. Epidural fibrin glue injection stops persistent postdural puncture headache. *Anesthesiology.* 1999;91:576-577.
237. Reynvoet ME, Cosaert PA, Desmet MF, Plasschaert SM. Epidural dextran 40 patch for postdural puncture headache. *Anaesthesia.* 1997;52:886-888.
238. McCormack JG, Malherbe S. Applications of ultrasound in paediatric anaesthesia. *Curr Anaesth Crit Care.* 2008;19:302-308.
239. Sharrock NE. Recordings of, and an anatomical explanation for, false positive loss of resistance during lumbar extradural analgesia. *Br J Anaesth.* 1979;51:253-258.
240. Gallart L, Blanco D, Samsó E, Vidal F. Clinical and radiologic evidence of the epidural plica mediana dorsalis. *Anesth Analg.* 1990;71:698-701.
241. Portnoy D, Vadhera RB. Mechanisms and management of an incomplete epidural block for cesarean section. *Anesthesiol Clin North America.* 2003;21:39-57.
242. Alahuhta S, Kangas-Saarela T, Hollmen AI, Edstrom HH. Visceral pain during caesarean section under spinal and epidural anaesthesia with bupivacaine. *Acta Anaesthesiol Scand.* 1990;34:95-98.
243. Bourne TM, deMelo AE, Bastianpillai BA, May AE. A survey of how British obstetric anaesthetists test regional anaesthesia before caesarean section. *Anaesthesia.* 1997;52:901-903.
244. Russell IF. Levels of anaesthesia and intraoperative pain at caesarean section under regional block. *Int J Obstet Anesth.* 1995;4:71-77.
245. Russell IF. The futility of using sharp pinprick (or cold) to assess spinal or epidural anesthesia for cesarean delivery. *Reg Anesth Pain Med.* 2001;26:385-387.
246. Wong CA, Norris MC. Acute situations: obstetrics. In: Raj PP, de Andres J, Grossi P, Banister R, Sala-Blanc X, eds. *Textbook of Regional Anesthesia.* New York, NY: Churchill Livingstone; 2002:471-504.
247. Tsui BC, Kulkarni P. Migration of a thoracic epidural catheter into the intercostal space via the intervertebral foramen. *Anesth Analg* 2009;108:371-373.
248. Borghi B, Casati A, Iuorio S, et al. Frequency of hypotension and bradycardia during general anesthesia, epidural anesthesia, or integrated epidural-general anesthesia for total hip replacement. *J Clin Anesth.* 2002;14:102-106.
249. Meissner A, Rolf N, Van Aken H. Thoracic epidural anesthesia and the patient with heart disease: benefits, risks, and controversies. *Anesth Analg.* 1997;85:517-528.
250. Schmidt C, Hinder F, Van Aken H, et al. The effect of high thoracic epidural anesthesia on systolic and diastolic left ventricular function in patients with coronary artery disease. *Anesth Analg.* 2005;100:1561-1569.
251. Alvarez J, Hernandez B, Atanassoff PG. High thoracic epidural anesthesia and coronary artery disease in surgical and non-surgical patients. *Curr Opin Anaesthesiol.* 2005;18:501-506.
252. Wright PM, Fee JP. Cardiovascular support during combined extradural and general anaesthesia. *Br J Anaesth.* 1992;68:585-589.
253. Hahn RG. Increased haemodilution in hypotension induced by epidural anaesthesia. *Acta Anaesthesiol Scand.* 1993;37:357-360.
254. Hahn RG. Haemoglobin dilution from epidural-induced hypotension with and without fluid loading. *Acta Anaesthesiol Scand.* 1992;36:241-244.
255. Holte K, Foss NB, Svensen C, et al. Epidural anesthesia, hypotension, and changes in intravascular volume. *Anesthesiology.* 2004;100:281-286.
256. Dalens B, Hasnaoui A. Caudal anesthesia in pediatric surgery: success rate and adverse effects in 750 consecutive patients. *Anesth Analg.* 1989;68:83-89.
257. Tsuji MH, Horigome H, Yamashita M. Left ventricular functions are not impaired after lumbar epidural anaesthesia in young children. *Paediatr Anaesth.* 1996;6:405-409.
258. McCarthy GS. The effect of thoracic extradural analgesia on pulmonary gas distribution, functional residual capacity and airway closure. *Br J Anaesth.* 1976;48:243-248.
259. Sundberg A, Wattwil M, Arvill A. Respiratory effects of high thoracic epidural anaesthesia. *Acta Anaesthesiol Scand.* 1986;30:215-217.
260. Kochi T, Sako S, Nishino T, Mizuguchi T. Effect of high thoracic extradural anaesthesia on ventilatory response to hypercapnia in normal volunteers. *Br J Anaesth.* 1989;62:362-367.
261. Sakura S, Saito Y, Kosaka Y. Effect of extradural anaesthesia on the ventilatory response to hypoxaemia. *Anaesthesia.* 1993;48:205-209.
262. Gruber EM, Tschernko EM, Kritzing M, et al. The effects of thoracic epidural analgesia with bupivacaine 0.25% on ventilatory mechanics in patients with severe chronic obstructive pulmonary disease. *Anesth Analg.* 2001;92:1015-1019.
263. Scammell SJ. Case report: inadvertent epidural anaesthesia as a complication of interscalene brachial plexus block. *Anaesth Intensive Care.* 1979;7:56-57.
264. Bonnet F, Derosier JP, Pluskwa F, Abhay K, Gaillard A. Cervical epidural anaesthesia for carotid artery surgery. *Can J Anaesth.* 1990;37:353-358.

265. Michalek P, David I, Adamec M, Janousek L. Cervical epidural anesthesia for combined neck and upper extremity procedure: a pilot study. *Anesth Analg*. 2004;99:1833-1836.
266. Nystrom UM, Nystrom NA. Continuous cervical epidural anesthesia in reconstructive hand surgery. *J Hand Surg Am*. 1997;22:906-912.
267. Tzeng JJ, Wang JJ, Ho ST, et al. Dexamethasone for prophylaxis of nausea and vomiting after epidural morphine for post-Caesarean section analgesia: comparison of droperidol and saline. *Br J Anaesth*. 2000;85:865-868.
268. Wang JJ, Ho ST, Liu YH, et al. Dexamethasone decreases epidural morphine-related nausea and vomiting. *Anesth Analg*. 1999;89:117-120.
269. Jeon Y, Hwang J, Kang J, et al. Effects of epidural naloxone on pruritus induced by epidural morphine: a randomized controlled trial. *Int J Obstet Anesth*. 2005;14:22-25.
270. Szarvas S, Harmon D, Murphy D. Neuraxial opioid-induced pruritus: a review. *J Clin Anesth*. 2003;15:234-239.
271. Ganesh A, Maxwell LG. Pathophysiology and management of opioid-induced pruritus. *Drugs*. 2007;67:2323-2333.
272. Emanuelsson BM, Persson J, Alm C, Heller A, Gustafsson LL. Systemic absorption and block after epidural injection of ropivacaine in healthy volunteers. *Anesthesiology*. 1997;87:1309-1317.
273. Borgeat A, Ekatomdramis G, Schenker CA. Postoperative nausea and vomiting in regional anesthesia: a review. *Anesthesiology*. 2003;98:530-547.
274. Sjostrom S, Tamsen A, Persson MP, Hartvig P. Pharmacokinetics of intrathecal morphine and meperidine in humans. *Anesthesiology*. 1987;67:889-895.
275. Lanz E, Kehrberger E, Theiss D. Epidural morphine: a clinical double-blind study of dosage. *Anesth Analg*. 1985;64:786-791.
276. Gedney JA, Liu EH. Side-effects of epidural infusions of opioid bupivacaine mixtures. *Anaesthesia*. 1998;53:1148-1155.
277. Tzeng JJ, Hsing CH, Chu CC, Chen YH, Wang JJ. Low-dose dexamethasone reduces nausea and vomiting after epidural morphine: a comparison of metoclopramide with saline. *J Clin Anesth*. 2002;14:19-23.
278. Wang J-J, Tzeng J-I, Ho S-T, et al. The prophylactic effect of tropisetron on epidural morphine-related nausea and vomiting: a comparison of dexamethasone with saline. *Anesth Analg*. 2002;94:749-753.
279. Allen TK, Habib AS. P6 stimulation for the prevention of nausea and vomiting associated with cesarean delivery under neuraxial anesthesia: a systematic review of randomized controlled trials. *Anesth Analg*. 2008;107:1308-1312.
280. Wein AJ. Neuromuscular dysfunction of the lower urinary tract and its management. In: Campbell MF, Walsh PC, Retik AB, eds. *Campbell's Urology*. 8th ed. Philadelphia, PA: Saunders; 2002:954.
281. Darrah DM, Griebing TL, Silverstein JH. Postoperative urinary retention. *Anesthesiol Clin*. 2009;27:465-484.
282. Givens CD, Wenzel RP. Catheter-associated urinary tract infections in surgical patients: a controlled study on the excess morbidity and costs. *J Urol*. 1980;124:646-648.
283. Platt R, Polk BF, Murdock B, Rosner B. Mortality associated with nosocomial urinary-tract infection. *N Engl J Med*. 1982;307:637-642.
284. Stenseth R, Sellevold O, Breivik H. Epidural morphine for postoperative pain: experience with 1085 patients. *Acta Anaesthesiol Scand*. 1985;29:148-156.
285. Rawal N, Mollefors K, Axelsson K, Lingardh G, Widman B. An experimental study of urodynamic effects of epidural morphine and of naloxone reversal. *Anesth Analg*. 1983;62:641-647.
286. Lytle SA, Goldsmith DM, Neuendorf TL, Lowry ME. Postoperative analgesia with epidural fentanyl. *J Am Osteopath Assoc*. 1991;91:547-550.
287. Lamonerie L, Marret E, Deleuze A, et al. Prevalence of postoperative bladder distension and urinary retention detected by ultrasound measurement. *Br J Anaesth*. 2004;92:544-546.
288. Keita H, Diouf E, Tubach F, et al. Predictive factors of early postoperative urinary retention in the postanesthesia care unit. *Anesth Analg*. 2005;101:592-596.
289. Baldini G, Bagry H, Aprikian A, Carli F. Postoperative urinary retention: anesthetic and perioperative considerations. *Anesthesiology*. 2009;110:1139-1157.
290. Rawal N, Mollefors K, Axelsson K, Lingardh G, Widman B. Naloxone reversal of urinary retention after epidural morphine. *Lancet*. 1981;2:1411-1411.
291. Malinovsky J-M, Lepage J-Y, Karam G, Pinaud M. Nalbuphine reverses urinary effects of epidural morphine: a case report. *J Clin Anesth*. 2002;14:535-538.
292. Basse L, Werner M, Kehlet H. Is urinary drainage necessary during continuous epidural analgesia after colonic resection? *Reg Anesth Pain Med*. 2000;25:498-501.
293. Brown EM, Elman DS. Postoperative backache. *Anesth Analg*. 1961;40:683-685.
294. MacArthur C, Lewis M, Knox EG, Crawford JS. Epidural anaesthesia and long term backache after childbirth. *BMJ*. 1990;301:9-12.
295. Russell R, Groves P, Taub N, O'Dowd J, Reynolds F. Assessing long term backache after childbirth. *BMJ*. 1993;306:1299-1303.
296. Breen TW, Ransil BJ, Groves PA, Oriol NE. Factors associated with back pain after childbirth. *Anesthesiology*. 1994;81:29-34.
297. Russell R, Dundas R, Reynolds F. Long term backache after childbirth: prospective search for causative factors. *BMJ*. 1996;312:1384-1388.
298. Levy L, Randel GI, Pandit SK. Does chloroprocaine (Nesacaine MPF) for epidural anesthesia increase the incidence of backache? *Anesthesiology*. 1989;71:476-476.
299. Drolet P, Veillette Y. Back pain following epidural anesthesia with 2-chloroprocaine (EDTA-free) or lidocaine. *Reg Anesth*. 1997;22:303-307.
300. Bier A. Uben einen neueun weg lokalanasthesie an den gliedmassen zu erzeugen. *Arch Clin Chir*. 1908;86:1007-1016.
301. Holmes CM. Intravenous regional analgesia. A useful method of producing analgesia of the limbs. *Lancet*. 1963;1:245-247.
302. Mabee JR, Bostwick TL, Burke MK. Iatrogenic compartment syndrome from hypertonic saline injection in Bier block. *J Emerg Med*. 1994;12:473-476.
303. Quigley JT, Popich GA, Lanz UB. Compartment syndrome of the forearm and hand: a case report. *Clin Orthop Relat Res*. 1981;247-251.
304. Luce EA, Mangubat E. Loss of hand and forearm following Bier block: a case report. *J Hand Surg Am*. 1983;8:280-283.
305. Carney AL, Anderson EM. Tourniquet subclavian steal: brainstem ischemia and cortical blindness—clinical significance and testing. *Adv Neurol*. 1981;30:283-290.

306. Pitkanen MT, Suzuki N, Rosenberg PH. Intravenous regional anaesthesia with 0.5% prilocaine or 0.5% chlorprocaine. A double-blind comparison in volunteers. *Anaesthesia*. 1992;47:618-619.
307. Asik I, Kocum AI, Goktug A, Turhan KS, Alkis N. Comparison of ropivacaine 0.2% and 0.25% with lidocaine 0.5% for intravenous regional anesthesia. *J Clin Anesth*. 2009;21:401-407.
308. Kumar C, Dowd T. Ophthalmic regional anaesthesia. *Curr Opin Anaesthesiol*. 2008;21:632-637.
309. Anonymous. Anaesthesia for cataract surgery. *Eur J Implant*. 1993;5:1-3.
310. Lichter PR. Avoiding complications from local anesthesia. *Ophthalmology*. 1988;95:565-566.
311. Smith R. Cataract extraction without retrobulbar anaesthetic injection. *Br J Ophthalmol*. 1990;74:205-207.
312. Shuler JD, Paschal JF, Holland GN. Antiplatelet therapy and cataract surgery. *J Cataract Refract Surg*. 1992;18:567-571.
313. Hamilton RC. Brain-stem anesthesia as a complication of regional anesthesia for ophthalmic surgery. *Can J Ophthalmol*. 1992;27:323-325.
314. Katsev DA, Drews RC, Rose BT. An anatomic study of retrobulbar needle path length. *Ophthalmology*. 1989;96:1221-1224.
315. Edge KR, Nicoll JM. Retrobulbar hemorrhage after 12,500 retrobulbar blocks. *Anesth Analg*. 1993;76:1019-1022.
316. Hamilton RC, Gimbel HV, Strunin L. Regional anaesthesia for 12,000 cataract extraction and intraocular lens implantation procedures. *Can J Anaesth*. 1988;35:615-623.
317. Manners TD, Burton RL. Randomised trial of topical versus sub-Tenon's local anaesthesia for small-incision cataract surgery. *Eye (Lond)*. 1996;10 (Pt 3):367-370.
318. Jeganathan VS, Jeganathan VP. Sub-Tenon's anaesthesia: a well tolerated and effective procedure for ophthalmic surgery. *Curr Opin Ophthalmol*. 2009;20:205-209.
319. Davis DB, Mandel MR. Posterior peribulbar anesthesia: an alternative to retrobulbar anesthesia. *J Cataract Refract Surg*. 1986;12:182-184.
320. Berglin L, Stenkula S, Algvere PV. Ocular perforation during retrobulbar and peribulbar injections. *Ophthalmic Surg Lasers*. 1995;26:429-434.
321. Hunter DG, Lam GC, Guyton DL. Inferior oblique muscle injury from local anesthesia for cataract surgery. *Ophthalmology*. 1995;102:501-509.
322. Peutrell JM, Lonnqvist P-A. Neuraxial blocks for anaesthesia and analgesia in children. *Curr Opin Anaesthesiol*. 2003;16:461-470.