

Chapter 51

REFERENCES

1. Jannetta PJ. Developments in neurosurgery: "the 4 factors." *Neurosurgery*. 2009;65(4 Suppl):A9-A10.
2. Ahmed A. Anaesthesia for interventional neuroradiology. *J Ayub Med Coll Abbottabad*. 2007;19(3):80-84.
3. Leithner C, Royl G, Offenhauser N, et al. Pharmacological uncoupling of activation induced increases in CBF and CMRO₂. *J Cereb Blood Flow Metab*. 2010;30(2):311-322.
4. Duncan R. Epilepsy, cerebral blood flow, and cerebral metabolic rate. *Cerebrovasc Brain Metab Rev*. 1992;4(2):105-121.
5. Mori K, Maeda M, Miyazaki M, Iwase H. Effects of mild (33°C) and moderate (29°C) hypothermia on cerebral blood flow and metabolism, lactate, and extracellular glutamate in experimental head injury. *Neurol Res*. 1998;20(8):719-726.
6. Royl G, Fuchtemeier M, Leithner C, et al. Hypothermia effects on neurovascular coupling and cerebral metabolic rate of oxygen. *Neuroimage*. 2008;40(4):1523-1532.
7. Sloan TB. Anesthetics and the brain. *Anesthesiol Clin North America*. 2002;20:265-292.
8. Reinstrup P, Ryding E, Algotsson L, Berntman L, Uski T. Effects of nitrous oxide on human regional cerebral blood flow and isolated pial arteries. *Anesthesiology*. 1994;81(2):396-402.
9. Schlunzen L, Cold GE, Rasmussen M, Vafaee MS. Effects of dose-dependent levels of isoflurane on cerebral blood flow in healthy subjects studied using positron emission tomography. *Acta Anaesthesiol Scand*. 2006;50(3):306-312.
10. Langsjo JW, Maksimow A, Salmi E, et al. S-ketamine anesthesia increases cerebral blood flow in excess of the metabolic needs in humans. *Anesthesiology*. 2005;103(2):258-268.
11. Ito H, Kanno I, Ibaraki M, Hatazawa J, Miura S. Changes in human cerebral blood flow and cerebral blood volume during hypercapnia and hypocapnia measured by positron emission tomography. *J Cereb Blood Flow Metab*. 2003;23(6):665-670.
12. Schlunzen L, Vafaee MS, Juul N, Cold GE. Regional cerebral blood flow responses to hyperventilation during sevoflurane anaesthesia studied with PET. *Acta Anaesthesiol Scand*. 2010;54(5):610-615.
13. Henriksen L, Paulson OB. The effects of sodium nitroprusside on cerebral blood flow and cerebral venous blood gases. II. Observations in awake man during successive blood pressure reduction. *Eur J Clin Invest*. 1982;12(5):389-393.
14. Schumann-Bard P, Touzani O, Young AR, et al. Cerebrovascular effects of sodium nitroprusside in the anaesthetized baboon: a positron emission tomographic study. *J Cereb Blood Flow Metab*. 2005;25(4):535-544.
15. Oshima T, Karasawa F, Satoh T. Effects of propofol on cerebral blood flow and the metabolic rate of oxygen in humans. *Acta Anaesthesiol Scand*. 2002;46(7):831-835.
16. Petersen KD, Landsfeldt U, Cold GE, et al. ICP is lower during propofol anaesthesia compared to isoflurane and sevoflurane. *Acta Neurochir Suppl*. 2002;81:89-91.
17. Voss LJ, Sleigh JW, Barnard JP, Kirsch HE. The howling cortex: seizures and general anesthetic drugs. *Anesth Analg*. 2008;107(5):1689-1703.
18. Krauss JK, Akeyson EW, Giam P, Jankovic J. Propofol-induced dyskinesias in Parkinson's disease. *Anesth Analg*. 1996;83(2):420-422.
19. Ramesh VJ, Umamaheswara Rao GS. Quantification of burst suppression and bispectral index with 2 different bolus doses of thiopentone sodium. *J Neurosurg Anesthesiol*. 2007;19(3):179-182.
20. Randomized clinical study of thiopental loading in comatose survivors of cardiac arrest. Brain Resuscitation Clinical Trial I Study Group. *N Engl J Med*. 1986;314(7):397-403.
21. Monsalve F, Rucabado L, Ruano M, et al. The neurologic effects of thiopental therapy after cardiac arrest. *Intensive Care Med*. 1987;13(4):244-248.
22. Ostwald P, Doenicke AW. Etomidate revisited. *Curr Opin Anaesthesiol*. 1998;11(4):391-398.
23. Hoffman WE, Charbel FT, Ausman JI. Cerebral blood flow and metabolic response to etomidate and ischemia. *Neurol Res*. 2007;19(1):41-44.
24. Pastor J, Wix R, Meilan ML, et al. Etomidate accurately localizes the epileptic area in patients with temporal lobe epilepsy. *Epilepsia*. 2010;51(4):602-609.
25. Mayberg TS, Lam AM, Matta BF, Domino KB, Winn HR. Ketamine does not increase cerebral blood flow velocity or intracranial pressure during isoflurane/nitrous oxide anesthesia in patients undergoing craniotomy. *Anesth Analg*. 1995;81(1):84-89.
26. Albanese J, Viviani X, Potie F, et al. Sufentanil, fentanyl, and alfentanil in head trauma patients: a study on cerebral hemodynamics. *Crit Care Med*. 1999;27(2):407-411.
27. Nakada J, Nishira M, Hosoda R, et al. Priming with rocuronium or vecuronium prevents remifentanil-mediated muscle rigidity and difficult ventilation. *J Anesth*. 2009;23(3):323-328.
28. Fodale V, Schifilliti D, Pratico C, Santamaria LB. Remifentanil and the brain. *Acta Anaesthesiol Scand*. 2008;52(3):319-326.

29. Angst MS, Koppert W, Pahl I, Clark DJ, Schmelz M. Short-term infusion of the μ -opioid agonist remifentanyl in humans causes hyperalgesia during withdrawal. *Pain*. 2003;106(1-2):49-57.
30. Hoffman WE, Miletich DJ, Albrecht RF. The effects of midazolam on cerebral blood flow and oxygen consumption and its interaction with nitrous oxide. *Anesth Analg*. 1986;65(7):729-733.
31. Farag E. Dexmedetomidine in the neurointensive care unit. *Discov Med*. 2010;9(44):42-45.
32. Rozet I. Anesthesia for functional neurosurgery: the role of dexmedetomidine. *Curr Opin Anaesthesiol*. 2008;21(5):537-543.
33. Yeh JS, Dhir JS, Green AL, Bodiwala D, Brydon HL. Changes in plasma phenytoin level following craniotomy. *Br J Neurosurg*. 2006;20(6):403-406.
34. Platt PR, Thackray NM. Phenytoin-induced resistance to vecuronium. *Anaesth Intensive Care*. 1993;21(2):185-191.
35. David HN, Leveille F, Chazalviel L, et al. Reduction of ischemic brain damage by nitrous oxide and xenon. *J Cereb Blood Flow Metab*. 2003;23(10):1168-1173.
36. Stirt JA, Grosslight KR, Bedford RF, Vollmer D. "Defasciculation" with metocurine prevents succinylcholine-induced increases in intracranial pressure. *Anesthesiology*. 1987;67(1):50-53.
37. Ducey JP, Deppe SA, Foley KT. A comparison of the effects of suxamethonium, atracurium and vecuronium on intracranial haemodynamics in swine. *Anaesth Intensive Care*. 1989;17(4):448-455.
38. Matta BF, Heath KJ, Tipping K, Summors AC. Direct cerebral vasodilatory effects of sevoflurane and isoflurane. *Anesthesiology*. 1999;91(3):677-680.
39. Matta BF, Lam AM. Nitrous oxide increases cerebral blood flow velocity during pharmacologically induced EEG silence in humans. *J Neurosurg Anesthesiol*. 1995;7(2):89-93.
40. Van Hemelrijck J, Fitch W, Mattheussen M, et al. Effect of propofol on cerebral circulation and autoregulation in the baboon. *Anesth Analg*. 1990;71(1):49-54.
41. Jevtovic-Todorovic V, Beals J, Benshoff N, Olney JW. Prolonged exposure to inhalational anesthetic nitrous oxide kills neurons in adult rat brain. *Neuroscience*. 2003;122(3):609-616.
42. Xiong L, Zheng Y, Wu M, et al. Preconditioning with isoflurane produces dose-dependent neuroprotection via activation of adenosine triphosphate-regulated potassium channels after focal cerebral ischemia in rats. *Anesth Analg*. 2003;96(1):233-237.
43. Eberspacher E, Eckel B, Engelhard K, et al. Effects of sevoflurane on cognitive deficit, motor function, and histopathology after cerebral ischemia in rats. *Acta Anaesthesiol Scand*. 2009;53(6):774-782.
44. McAuliffe JJ, Loepke AW, Miles L, et al. Desflurane, isoflurane, and sevoflurane provide limited neuroprotection against neonatal hypoxia-ischemia in a delayed preconditioning paradigm. *Anesthesiology*. 2009;111(3):533-546.
45. Codaccioni JL, Velly LJ, Moubarik C, et al. Sevoflurane preconditioning against focal cerebral ischemia: inhibition of apoptosis in the face of transient improvement of neurological outcome. *Anesthesiology*. 2009;110(6):1271-1278.
46. Baumgartner WA. Neurologic injury after cardiopulmonary bypass surgery. *J Neurosurg Anesthesiol*. 2004;16(1):102-104.
47. Ikonomidou C, Turski L. Why did NMDA receptor antagonists fail clinical trials for stroke and traumatic brain injury? *Lancet Neurol*. 2002;1(6):383-386.
48. Hardingham GE, Bading H. The Yin and Yang of NMDA receptor signaling. *Trends Neurosci*. 2003;26(2):81-89.
49. Allan SM, Rothwell NJ. Cytokines and acute neurodegeneration. *Nat Rev Neurosci*. 2001;2(10):734-744.
50. Mantz J, Degos V, Laigle C. Recent advances in pharmacologic neuroprotection. *Eur J Anaesthesiol*. 2010;27(1):6-10.
51. Palace J. Neuroprotection and repair. *J Neurol Sci*. 2008;265(1-2):21-25.
52. Yatsu FM, Diamond I, Graziano C, Lindquist P. Experimental brain ischemia: protection from irreversible damage with a rapid-acting barbiturate (methohexital). *Stroke*. 1972;3(6):726-732.
53. Todd MM, Chadwick HS, Shapiro HM, et al. The neurologic effects of thiopental therapy following experimental cardiac arrest in cats. *Anesthesiology*. 1982;57(2):76-86.
54. Koerner IP, Brambrink AM. Brain protection by anesthetic agents. *Curr Opin Anaesthesiol*. 2006;19(5):481-486.
55. Warner DS, Takaoka S, Wu B, et al. Electroencephalographic burst suppression is not required to elicit maximal neuroprotection from pentobarbital in a rat model of focal cerebral ischemia. *Anesthesiology*. 1996;84(6):1475-1484.
56. Head BP, Patel P. Anesthetics and brain protection. *Curr Opin Anaesthesiol*. 2007;20(5):395-399.
57. Kochs E, Hoffman WE, Werner C, et al. The effects of propofol on brain electrical activity, neurologic outcome, and neuronal damage following incomplete ischemia in rats. *Anesthesiology*. 1992;76(2):245-252.
58. Yamaguchi S, Midorikawa Y, Okuda Y, Kitajima T. Propofol prevents delayed neuronal death following transient forebrain ischemia in gerbils. *Can J Anaesth*. 1999;46(6):593-598.
59. Kanbak M, Saricaoglu F, Avci A, et al. Propofol offers no advantage over isoflurane anesthesia for cerebral protection during cardiopulmonary bypass: a preliminary study of S-100beta protein levels. *Can J Anaesth*. 2004;51(7):712-717.
60. Hoffman WE, Kochs E, Werner C, Thomas C, Albrecht RF. Dexmedetomidine improves neurologic outcome from incomplete ischemia in the rat. Reversal by the α_2 -adrenergic antagonist atipamezole. *Anesthesiology*. 1991;75(2):328-332.
61. Sanders RD, Ma D, Maze M. Anaesthesia induced neuroprotection. *Best Pract Res Clin Anaesthesiol*. 2005;19(3):461-474.
62. Himmelseher S, Durieux ME. Revising a dogma: ketamine for patients with neurological injury? *Anesth Analg*. 2005;101(2):524-534.
63. Nagels W, Demeyere R, Van Hemelrijck J, et al. Evaluation of the neuroprotective effects of S(+)-ketamine during open-heart surgery. *Anesth Analg*. 2004;98(6):1595-1603.
64. Drummond JC, McKay LD, Cole DJ, Patel PM. The role of nitric oxide synthase inhibition in the adverse effects of etomidate in the setting of focal cerebral ischemia in rats. *Anesth Analg*. 2005;100(3):841-846.
65. Hoffman WE, Charbel FT, Edelman G, Misra M, Ausman JJ. Comparison of the effect of etomidate and desflurane on brain tissue gases and pH during prolonged middle cerebral artery occlusion. *Anesthesiology*. 1998;88(5):1188-1194.
66. Lei B, Popp S, Capuano-Waters C, Cottrell JE, Kass IS. Lidocaine attenuates apoptosis in the ischemic penumbra and reduces infarct size after transient focal cerebral ischemia in rats. *Neuroscience*. 2004;125(3):691-701.
67. Niyama S, Tanaka E, Tsuji S, et al. Neuroprotective mechanisms of lidocaine against in vitro ischemic insult of the rat hippocampal CA1 pyramidal neurons. *Neurosci Res*. 2005;53(3):271-278.

68. Patel PMM, Drummond JCM, Cole DJM, Goskowitz RLM. Isoflurane reduces ischemia-induced glutamate release in rats subjected to forebrain ischemia. *Anesthesiology*. 1995;82(4):996-1003.
69. Bickler PE, Fahlman CS. The inhaled anesthetic, isoflurane, enhances Ca²⁺-dependent survival signaling in cortical neurons and modulates MAP kinases, apoptosis proteins and transcription factors during hypoxia. *Anesth Analg*. 2006;103(2):419-429.
70. Kitano H, Young JM, Cheng J, et al. Gender-specific response to isoflurane preconditioning in focal cerebral ischemia. *J Cereb Blood Flow Metab*. 2007;27(7):1377-1386.
71. Franks NP, Honore E. The TREK K₂P channels and their role in general anaesthesia and neuroprotection. *Trends Pharmacol Sci*. 2004;25(11):601-608.
72. Heurteaux C, Guy N, Laigle C, et al. TREK-1, a K⁺ channel involved in neuroprotection and general anesthesia. *EMBO J*. 2004;23(13):2684-2695.
73. Jevtovic-Todorovic V, Hartman RE, Izumi Y, et al. Early exposure to common anesthetic agents causes widespread neurodegeneration in the developing rat brain and persistent learning deficits. *J Neurosci*. 2003;23(3):876-882.
74. Dingley J, Tooley J, Porter H, Thoresen M. Xenon provides short-term neuroprotection in neonatal rats when administered after hypoxia-ischemia. *Stroke*. 2006;37(2):501-506.
75. Ma D, Hossain M, Pettet GK, et al. Xenon preconditioning reduces brain damage from neonatal asphyxia in rats. *J Cereb Blood Flow Metab*. 2006;26(2):199-208.
76. Fornai F, Longone P, Cafaro L, et al. Lithium delays progression of amyotrophic lateral sclerosis. *Proc Natl Acad Sci U S A*. 2008;105(6):2052-2057.
77. Verdonck O, Lahrech H, Francony G, et al. Erythropoietin protects from post-traumatic edema in the rat brain. *J Cereb Blood Flow Metab*. 2007;27(7):1369-1376.
78. Leist M, Ghezzi P, Grasso G, et al. Derivatives of erythropoietin that are tissue protective but not erythropoietic. *Science*. 2004;305(5681):239-242.
79. Kreuter M, Langer C, Kerkhoff C, et al. Stroke, myocardial infarction, acute and chronic inflammatory diseases: caspases and other apoptotic molecules as targets for drug development. *Arch Immunol Ther Exp (Warsz)*. 2004;52(3):141-155.
80. Irazuzta J, Pretzlaff RK, DeCourten-Myers G, Zemlan F, Zingarelli B. Dexamethasone decreases neurological sequelae and caspase activity. *Intensive Care Med*. 2005;31(1):146-150.
81. Hypothermia after Cardiac Arrest Study Group. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med*. 2002;346(8):549-556.
82. Todd MM, Hindman BJ, Clarke WR, Torner JC; Intraoperative Hypothermia for Aneurysm Surgery Trial (IHAST) Investigators. Mild intraoperative hypothermia during surgery for intracranial aneurysms. *N Engl J Med*. 2005;352(2):135-145.
83. Singhal AB, Lo EH, Dalkara T, Moskowitz MA. Advances in stroke neuroprotection: hyperoxia and beyond. *Neuroimaging Clin N Am*. 2005;15(3):697-720.
84. Hancock SM, Eastwood JR, Mahajan RP. Effects of inhaled nitrous oxide 50% on estimated cerebral perfusion pressure and zero flow pressure in healthy volunteers. *Anaesthesia*. 2005;60(2):129-132.
85. Randell T, Niskanen M. Management of physiological variables in neuroanaesthesia: maintaining homeostasis during intracranial surgery. *Curr Opin Anaesthesiol*. 2006;19(5):492-497.
86. Rasmussen M, Bundgaard H, Cold GE. Craniotomy for supratentorial brain tumors: risk factors for brain swelling after opening the dura mater. *J Neurosurg*. 2004;101(4):621-626.
87. Sharma D, Bithal PK, Dash HH, et al. Cerebral autoregulation and CO₂ reactivity before and after elective supratentorial tumor resection. *J Neurosurg Anesthesiol*. 2010;22(2):132-137.
88. Bundgaard H, Jensen K, Cold GE, et al. Effects of perioperative indomethacin on intracranial pressure, cerebral blood flow, and cerebral metabolism in patients subjected to craniotomy for cerebral tumors. *J Neurosurg Anesthesiol*. 1996;8(4):273-279.
89. Rasmussen M, Ostergaard L, Juul N, et al. Do indomethacin and propofol cause cerebral ischemic damage? Diffusion-weighted magnetic resonance imaging in patients undergoing craniotomy for brain tumors. *Anesthesiology*. 2004;101(4):872-878.
90. Ringel F, Chang RC, Staub F, Baethmann A, Plesnila N. Contribution of anion transporters to the acidosis-induced swelling and intracellular acidification of glial cells. *J Neurochem*. 2000;75(1):125-132.
91. Holmstrom A, Akeson J. Desflurane increases intracranial pressure more and sevoflurane less than isoflurane in pigs subjected to intracranial hypertension. *J Neurosurg Anesthesiol*. 2004;16(2):136-143.
92. Kaye A, Kucera IJ, Heavner J, et al. The comparative effects of desflurane and isoflurane on lumbar cerebrospinal fluid pressure in patients undergoing craniotomy for supratentorial tumors. *Anesth Analg*. 2004;98(4):1127-1132.
93. Magni G, Rosa IL, Melillo G, Savio A, Rosa G. A comparison between sevoflurane and desflurane anesthesia in patients undergoing craniotomy for supratentorial intracranial surgery. *Anesth Analg*. 2009;109(2):567-571.
94. Holmstrom A, Akeson J. Sevoflurane induces less cerebral vasodilation than isoflurane at the same A-line autoregressive index level. *Acta Anaesthesiol Scand*. 2005;49(1):16-22.
95. Cole CD, Gottfried ON, Gupta DK, Couldwell WT. Total intravenous anesthesia: advantages for intracranial surgery. *Neurosurgery*. 2007;61(5 Suppl 2):369-377.
96. Engelhard K, Werner C. Inhalational or intravenous anesthetics for craniotomies? Pro inhalational. *Curr Opin Anaesthesiol*. 2006;19(5):504-508.
97. Kaisti KK, Langsjö JW, Aalto S, et al. Effects of sevoflurane, propofol, and adjunct nitrous oxide on regional cerebral blood flow, oxygen consumption, and blood volume in humans. *Anesthesiology*. 2003;99(3):603-613.
98. Kawano Y, Kawaguchi M, Inoue S, et al. Jugular bulb oxygen saturation under propofol or sevoflurane/nitrous oxide anesthesia during deliberate mild hypothermia in neurosurgical patients. *J Neurosurg Anesthesiol*. 2004;16(1):6-10.
99. Liao R, Li J, Liu J. Volatile induction/maintenance of anaesthesia with sevoflurane increases jugular venous oxygen saturation and lumbar cerebrospinal fluid pressure in patients undergoing craniotomy. *Eur J Anaesthesiol*. 2010;27(4):369-376.
100. Bein B, Renner J, Caliebe D, et al. Sevoflurane but not propofol preserves myocardial function during minimally invasive direct coronary artery bypass surgery. *Anesth Analg*. 2005;100(3):610-616.
101. Filipovic M, Wang J, Michaux I, et al. Effects of halothane, sevoflurane and propofol on left ventricular diastolic function in humans during spontaneous and mechanical ventilation. *Br J Anaesth*. 2005;94(2):186-192.

102. Marval PD, Perrin ME, Hancock SM, Mahajan RP. The effects of propofol or sevoflurane on the estimated cerebral perfusion pressure and zero flow pressure. *Anesth Analg*. 2005;100(3):835-840.
103. Lauta E, Abbinante C, Del Gaudio A, et al. Emergence times are similar with sevoflurane and total intravenous anesthesia: results of a multicenter RCT of patients scheduled for elective supratentorial craniotomy. *J Neurosurg Anesthesiol*. 2010;22(2):110-118.
104. Liolios A, Guerit JM, Scholtes JL, Raftopoulos C, Hantson P. Propofol infusion syndrome associated with short-term large-dose infusion during surgical anesthesia in an adult. *Anesth Analg*. 2005;100(6):1804-1806.
105. Cremer OL, Moons KG, Bouman EA, et al. Long-term propofol infusion and cardiac failure in adult head-injured patients. *Lancet*. 2001;357(9250):117-118.
106. Girling KJ, Cavill G, Mahajan RP. The effects of nitrous oxide and oxygen on transient hyperemic response in human volunteers. *Anesth Analg*. 1999;89(1):175-180.
107. Tanskanen PE, Kytta JV, Randell TT, Aantaa RE. Dexmedetomidine as an anaesthetic adjuvant in patients undergoing intracranial tumour surgery: a double-blind, randomized and placebo-controlled study. *Br J Anaesth*. 2006;97(5):658-665.
108. Muizelaar JP, van der Poel HG, Li ZC, Kontos HA, Levasseur JE. Pial arteriolar vessel diameter and CO₂ reactivity during prolonged hyperventilation in the rabbit. *J Neurosurg*. 1988;69(6):923-927.
109. Wilkinson ID, Jellineck DA, Levy D, et al. Dexamethasone and enhancing solitary cerebral mass lesions: alterations in perfusion and blood-tumor barrier kinetics shown by magnetic resonance imaging. *Neurosurgery*. 2006;58(4):640-646.
110. Bell BA, Smith MA, Kean DM, et al. Brain water measured by magnetic resonance imaging. Correlation with direct estimation and changes after mannitol and dexamethasone. *Lancet*. 1987;1(8524):66-69.
111. Lukins MB, Manninen PH. Hyperglycemia in patients administered dexamethasone for craniotomy. *Anesth Analg*. 2005;100(4):1129-1133.
112. Schwarz G, Fuchs G, Weihs W, et al. Sitting position for neurosurgery: experience with preoperative contrast echocardiography in 301 patients. *J Neurosurg Anesthesiol*. 1994;6(2):83-88.
113. Mammoto T, Hayashi Y, Ohnishi Y, Kuro M. Incidence of venous and paradoxical air embolism in neurosurgical patients in the sitting position: detection by transesophageal echocardiography. *Acta Anaesthesiol Scand*. 1998;42(6):643-647.
114. Black S, Cucchiara RF, Nishimura RA, Michenfelder JD. Parameters affecting occurrence of paradoxical air embolism. *Anesthesiology*. 1989;71(2):235-241.
115. Black S, Ockert DB, Oliver WC Jr, Cucchiara RF. Outcome following posterior fossa craniectomy in patients in the sitting or horizontal positions. *Anesthesiology*. 1988;69(1):49-56.
116. Reasoner DK, Todd MM, Scamman FL, Warner DS. The incidence of pneumocephalus after supratentorial craniotomy. Observations on the disappearance of intracranial air. *Anesthesiology*. 1994;80(5):1008-1012.
117. Bundgaard H, von Oettingen G, Larsen KM, et al. Effects of sevoflurane on intracranial pressure, cerebral blood flow and cerebral metabolism. A dose-response study in patients subjected to craniotomy for cerebral tumours. *Acta Anaesthesiol Scand*. 1998;42(6):621-627.
118. Rolighed Larsen JK, Haure P, Cold GE. Reverse Trendelenburg position reduces intracranial pressure during craniotomy. *J Neurosurg Anesthesiol*. 2002;14(1):16-21.
119. Schramko A, Suojaranta-Ylinen R, Kuitunen A, et al. Hydroxyethylstarch and gelatin solutions impair blood coagulation after cardiac surgery: a prospective randomized trial. *Br J Anaesth*. 2010;104:691-697.
120. Niemi TT, Kuitunen AH. Hydroxyethyl starch impairs in vitro coagulation. *Acta Anaesthesiol Scand*. 1998;42(9):1104-1109.
121. Schramko AA, Suojaranta-Ylinen RT, Kuitunen AH, Kukkonen SI, Niemi TT. Rapidly degradable hydroxyethyl starch solutions impair blood coagulation after cardiac surgery: a prospective randomized trial. *Anesth Analg*. 2009;108(1):30-36.
122. Kawaguchi M, Sakamoto T, Furuya H, Ohnishi H, Karasawa J. Pseudoankylosis of the mandible after supratentorial craniotomy. *Anesth Analg*. 1996;83(4):731-734.
123. Marquardt G, Setzer M, Schick U, Seifert V. Cerebellar hemorrhage after supratentorial craniotomy. *Surg Neurol*. 2002;57(4):241-251.
124. Verchere E, Grenier B, Mesli A, et al. Postoperative pain management after supratentorial craniotomy. *J Neurosurg Anesthesiol*. 2002;14(2):96-101.
125. Rapana A, Lamaida E, Pizza V. Multiple postoperative intracerebral haematomas remote from the site of craniotomy. *Br J Neurosurg*. 1998;12(4):364-368.
126. Colao A. Pituitary tumours: the prolactinoma. *Best Pract Res Clin Endocrinol Metab*. 2009;23(5):575-596.
127. Karavitaki N, Wass JA. Non-adenomatous pituitary tumours. *Best Pract Res Clin Endocrinol Metab*. 2009;23(5):651-665.
128. Bonert V. Diagnostic challenges in acromegaly: a case-based review. *Best Pract Res Clin Endocrinol Metab*. 2009;23(Suppl 1):S23-S30.
129. Bertagna X, Guignat L, Groussin L, Bertherat J. Cushing's disease. *Best Pract Res Clin Endocrinol Metab*. 2009;23(5):607-623.
130. Beck-Peccoz P, Persani L, Mannavola D, Campi I. Pituitary tumours: TSH-secreting adenomas. *Best Pract Res Clin Endocrinol Metab*. 2009;23(5):597-606.
131. Pereira O, Bevan JS. Preoperative assessment for pituitary surgery. *Pituitary*. 2008;11(4):347-351.
132. Kim SH, Lee KC, Kim SH. Cranial nerve palsies accompanying pituitary tumour. *J Clin Neurosci*. 2007;14(12):1158-1162.
133. Schlechte JA. Long-term management of prolactinomas. *J Clin Endocrinol Metab*. 2007;92(8):2861-2865.
134. Nemergut EC, Dumont AS, Barry UT, Laws ER. Perioperative management of patients undergoing transsphenoidal pituitary surgery. *Anesth Analg*. 2005;101(4):1170-1181.
135. Scacchi M, Cavagnini F. Acromegaly. *Pituitary*. 2006;9(4):297-303.
136. Gross BA, Mindea SA, Pick AJ, Chandler JP, Batjer HH. Diagnostic approach to Cushing disease. *Neurosurg Focus*. 2007;23(3):E1.
137. Vance ML. Perioperative management of patients undergoing pituitary surgery. *Endocrinol Metab Clin North Am*. 2003;32(2):355-365.
138. Schteingart DE. Drugs in the medical treatment of Cushing's syndrome. *Expert Opin Emerg Drugs*. 2009;14(4):661-671.
139. Marquardt G, Yahya H, Hermann E, Seifert V. Direct transnasal approach for pituitary surgery. *Neurosurg Rev*. 2004;27(2):83-88.

140. Neal JG, Patel SJ, Kulbersh JS, Osguthorpe JD, Schlosser RJ. Comparison of techniques for transsphenoidal pituitary surgery. *Am J Rhinol.* 2007;21(2):203-206.
141. Maurtua MA, Deogaonkar A, Bakri MH, et al. Dosing of remifentanyl to prevent movement during craniotomy in the absence of neuromuscular blockade. *J Neurosurg Anesthesiol.* 2008;20(4):221-225.
142. Avitsian R, Schubert A. Anesthetic considerations for intraoperative management of cerebrovascular disease in neurovascular surgical procedures. *Anesthesiol Clin.* 2007;25(3):441-463.
143. Zhang J, Clatterbuck RE. Molecular genetics of human intracranial aneurysms. *Int J Stroke.* 2008;3(4):272-287.
144. Rosen DS, Macdonald RL. Subarachnoid hemorrhage grading scales: a systematic review. *Neurocrit Care.* 2005;2(2):110-118.
145. Dupont SA, Wijidicks EF, Manno EM, et al. Timing of computed tomography and prediction of vasospasm after aneurysmal subarachnoid hemorrhage. *Neurocrit Care.* 2009;11(1):71-75.
146. Frontera JA, Claassen J, Schmidt JM, et al. Prediction of symptomatic vasospasm after subarachnoid hemorrhage: the modified fisher scale. *Neurosurgery.* 2006;59(1):21-27.
147. McLaughlin N, Bojanowski MW. Aneurysmal surgery in the presence of angiographic vasospasm: an outcome assessment. *Can J Neurol Sci.* 2006;33(2):181-188.
148. Kassell NF, Torner JC, Jane JA, Haley EC Jr, Adams HP. The International Cooperative Study on the Timing of Aneurysm Surgery. Part 2: Surgical results. *J Neurosurg.* 1990;73(1):37-47.
149. Whitfield PC, Kirkpatrick PJ. Timing of surgery for aneurysmal subarachnoid haemorrhage. *Cochrane Database Syst Rev.* 2001;(2):CD001697.
150. Muroi C, Keller M, Pangalu A, et al. Neurogenic pulmonary edema in patients with subarachnoid hemorrhage. *J Neurosurg Anesthesiol.* 2008;20(3):188-192.
151. Baumann A, Audibert G, McDonnell J, Mertes PM. Neurogenic pulmonary edema. *Acta Anaesthesiol Scand.* 2007;51(4):447-455.
152. McLaughlin N, Bojanowski MW, Girard F, Denault A. Pulmonary edema and cardiac dysfunction following subarachnoid hemorrhage. *Can J Neurol Sci.* 2005;32(2):178-185.
153. Lee VH, Oh JK, Mulvagh SL, Wijidicks EF. Mechanisms in neurogenic stress cardiomyopathy after aneurysmal subarachnoid hemorrhage. *Neurocrit Care.* 2006;5(3):243-249.
154. Bulsara KR, McGirt MJ, Liao L, et al. Use of the peak troponin value to differentiate myocardial infarction from reversible neurogenic left ventricular dysfunction associated with aneurysmal subarachnoid hemorrhage. *J Neurosurg.* 2003;98(3):524-528.
155. Kao L, Al Lawati Z, Vavao J, Steinberg GK, Katznelson L. Prevalence and clinical demographics of cerebral salt wasting in patients with aneurysmal subarachnoid hemorrhage. *Pituitary.* 2009;12(4):347-351.
156. Brilstra EH, Rinkel GJ, Algra A, van Gijn J. Rebleeding, secondary ischemia, and timing of operation in patients with subarachnoid hemorrhage. *Neurology.* 2000;55(11):1656-1660.
157. Diringner MN. Management of aneurysmal subarachnoid hemorrhage. *Crit Care Med.* 2009;37(2):432-440.
158. Keyrouz SG, Diringner MN. Clinical review: prevention and therapy of vasospasm in subarachnoid hemorrhage. *Crit Care.* 2007;11(4):220.
159. Tseng MY, Czosnyka M, Richards H, Pickard JD, Kirkpatrick PJ. Effects of acute treatment with pravastatin on cerebral vasospasm, autoregulation, and delayed ischemic deficits after aneurysmal subarachnoid hemorrhage: a phase II randomized placebo-controlled trial. *Stroke.* 2005;36(8):1627-1632.
160. Kramer AH, Gurka MJ, Nathan B, et al. Statin use was not associated with less vasospasm or improved outcome after subarachnoid hemorrhage. *Neurosurgery.* 2008;62(2):422-427.
161. Badjatia N, Topcuoglu MA, Pryor JC, et al. Preliminary experience with intra-arterial nicardipine as a treatment for cerebral vasospasm. *Am J Neuroradiol.* 2004;25(5):819-826.
162. Hosoda K, Fujita S, Kawaguchi T, et al. Effect of clot removal and surgical manipulation on regional cerebral blood flow and delayed vasospasm in early aneurysm surgery for subarachnoid hemorrhage. *Surg Neurol.* 1999;51(1):81-88.
163. Riina HA, Lemole GM Jr, Spetzler RF. Anterior communicating artery aneurysms. *Neurosurgery.* 2002;51(4):993-996.
164. Hindman BJ, Bayman EO, Pfisterer WK, et al. No association between intraoperative hypothermia or supplemental protective drug and neurologic outcomes in patients undergoing temporary clipping during cerebral aneurysm surgery: findings from the Intraoperative Hypothermia for Aneurysm Surgery Trial. *Anesthesiology.* 2010;112(1):86-101.
165. Imizu S, Kato Y, Sangli A, Oguri D, Sano H. Assessment of incomplete clipping of aneurysms intraoperatively by a near-infrared indocyanine green-video angiography (Niicg-Va) integrated microscope. *Minim Invasive Neurosurg.* 2008;51(4):199-203.
166. Grzyska U, Fiehler J. Pathophysiology and treatment of brain AVMs. *Klin Neuroradiol.* 2009;19(1):82-90.
167. Spetzler RF, Martin NA. A proposed grading system for arteriovenous malformations. *J Neurosurg.* 1986;65(4):476-483.
168. Starke RM, Komotar RJ, Hwang BY, et al. Treatment guidelines for cerebral arteriovenous malformation microsurgery. *Br J Neurosurg.* 2009;23(4):376-386.
169. Spetzler RF, Wilson CB, Weinstein P, et al. Normal perfusion pressure breakthrough theory. *Clin Neurosurg.* 1978;25:651-672.
170. al Rodhan NR, Sundt TM Jr, Piepgras DG, et al. Occlusive hyperemia: a theory for the hemodynamic complications following resection of intracerebral arteriovenous malformations. *J Neurosurg.* 1993;78(2):167-175.
171. Hashimoto N, Nozaki K, Takagi Y, Kikuta K, Mikuni N. Surgery of cerebral arteriovenous malformations. *Neurosurgery.* 2007;61(1 Suppl):375-387.
172. Frost EA, Booij LH. Anesthesia in the patient for awake craniotomy. *Curr Opin Anaesthesiol.* 2007;20(4):331-335.
173. Bustillo MA, Lazar RM, Finck AD, et al. Dexmedetomidine may impair cognitive testing during endovascular embolization of cerebral arteriovenous malformations: a retrospective case report series. *J Neurosurg Anesthesiol.* 2002;14(3):209-212.
174. Wanebo JE, Zabramski JM, Spetzler RF. Superficial temporal artery-to-middle cerebral artery bypass grafting for cerebral revascularization. *Neurosurgery.* 2004;55(2):395-398.
175. Bilotta F, Rosa G. 'Anesthesia' for awake neurosurgery. *Curr Opin Anaesthesiol.* 2009;22(5):560-565.
176. Costello TG, Cormack JR, Hoy C, et al. Plasma ropivacaine levels following scalp block for awake craniotomy. *J Neurosurg Anesthesiol.* 2004;16(2):147-150.
177. Audu PB, Wilkerson C, Bartkowski R, et al. Plasma ropivacaine levels during awake intracranial surgery. *J Neurosurg Anesthesiol.* 2005;17(3):153-155.

178. Farag E, Deboer G, Cohen BH, Niezgodka J. Metabolic acidosis due to propofol infusion. *Anesthesiology*. 2005;102(3):697-698.
179. Bekker A, Sturaitis MK. Dexmedetomidine for neurological surgery. *Neurosurgery*. 2005;57(1 Suppl):1-10.
180. Tanaka K, Oda Y, Funao T, et al. Dexmedetomidine decreases the convulsive potency of bupivacaine and levobupivacaine in rats: involvement of α_2 -adrenoceptor for controlling convulsions. *Anesth Analg*. 2005;100(3):687-696.
181. Szelenyi A, Joksimovic B, Seifert V. Intraoperative risk of seizures associated with transient direct cortical stimulation in patients with symptomatic epilepsy. *Clin Neurophysiol*. 2007;24(1):39-43.
182. Conte V, Baratta P, Tomaselli P, et al. Awake neurosurgery: an update. *Minerva Anestesiologica*. 2008;74(6):289-292.
183. Benabid AL, Pollak P, Gervason C, et al. Long-term suppression of tremor by chronic stimulation of the ventral intermediate thalamic nucleus. *Lancet*. 1991;337(8738):403-406.
184. Deiner S, Hagen J. Parkinson's disease and deep brain stimulator placement. *Anesthesiol Clin*. 2009;27(3):391-415.
185. Frost EA, Osborn I. Deep brain stimulation—surgery for movement disorders and Parkinson's disease. *Int Anesthesiol Clin*. 2009;47(2):57-68.
186. Fickers BF, Zandstra DF. Primary laryngospasm in a patient with Parkinson's disease: treatment with CPAP via mini-tracheostomy following intubation. *Intensive Care Med*. 1995;21(10):863-864.
187. Khatib R, Ebrahim Z, Rezai A, et al. Perioperative events during deep brain stimulation: the experience at Cleveland Clinic. *J Neurosurg Anesthesiol*. 2008;20(1):36-40.
188. *Recommendation of the ICRP*. Elmsford, NY: Pergamon Press; 1991:1-25.
189. Varma MK, Price K, Jayakrishnan V, Manickam B, Kessell G. Anaesthetic considerations for interventional neuroradiology. *Br J Anaesth*. 2007;99(1):75-85.
190. Rinkel GJ, Djibuti M, Algra A, van Gijn J. Prevalence and risk of rupture of intracranial aneurysms: a systematic review. *Stroke*. 1998;29(1):251-256.
191. Vinuela F, Duckwiler G, Mawad M. Guglielmi detachable coil embolization of acute intracranial aneurysm: perioperative anatomical and clinical outcome in 403 patients. *J Neurosurg*. 1997;86(3):475-482.
192. Malisch TW, Guglielmi G, Vinuela F, et al. Intracranial aneurysms treated with the Guglielmi detachable coil: midterm clinical results in a consecutive series of 100 patients. *J Neurosurg*. 1997;87(2):176-183.
193. Molyneux AJ, Kerr RS, Yu LM, et al. International subarachnoid aneurysm trial (ISAT) of neurosurgical clipping versus endovascular coiling in 2143 patients with ruptured intracranial aneurysms: a randomised comparison of effects on survival, dependency, seizures, rebleeding, subgroups, and aneurysm occlusion. *Lancet*. 2005;366(9488):809-817.
194. Brisman JL, Song JK, Newell DW. Cerebral aneurysms. *N Engl J Med*. 2006;355(9):928-939.
195. Crawford PM, West CR, Chadwick DW, Shaw MD. Arteriovenous malformations of the brain: natural history in unoperated patients. *J Neurol Neurosurg Psychiatry*. 1986;49(1):1-10.
196. Spetzler RF, Martin NA, Carter LP, et al. Surgical management of large AVMs by staged embolization and operative excision. *J Neurosurg*. 1987;67(1):17-28.
197. Lakhani S, Guha A, Nahser HC. Anaesthesia for endovascular management of cerebral aneurysms. *Eur J Anaesthesiol*. 2006;23(11):902-913.
198. Davies KR, Gelb AW, Manninen PH, Boughner DR, Bisnaire D. Cardiac function in aneurysmal subarachnoid haemorrhage: a study of electrocardiographic and echocardiographic abnormalities. *Br J Anaesth*. 1991;67(1):58-63.
199. Naredi S, Lambert G, Eden E, et al. Increased sympathetic nervous activity in patients with nontraumatic subarachnoid hemorrhage. *Stroke*. 2000;31(4):901-906.
200. Report of World Federation of Neurological Surgeons Committee on a universal subarachnoid hemorrhage grading scale. *J Neurosurg*. 1988;68(6):985-986.
201. Claassen J, Vu A, Kreiter KT, et al. Effect of acute physiologic derangements on outcome after subarachnoid hemorrhage. *Crit Care Med*. 2004;32(3):832-838.
202. Maroon JC, Nelson PB. Hypovolemia in patients with subarachnoid hemorrhage: therapeutic implications. *Neurosurgery*. 1979;4(3):223-226.
203. Kasuya H, Onda H, Yoneyama T, Sasaki T, Hori T. Bedside monitoring of circulating blood volume after subarachnoid hemorrhage. *Stroke*. 2003;34(4):956-960.
204. van den Bergh WM, Algra A, van der Sprenkel JW, Tulleken CA, Rinkel GJ. Hypomagnesemia after aneurysmal subarachnoid hemorrhage. *Neurosurgery*. 2003;52(2):276-281.
205. Castagnini HE, van Eijs F, Salevsky FC, Nathanson MH. Sevoflurane for interventional neuroradiology procedures is associated with more rapid early recovery than propofol. *Can J Anaesth*. 2004;51(5):486-491.
206. Johnson JO. Anesthesia for minimally invasive neurosurgery. *Anesthesiol Clin North America*. 2002;20(2):361-375.
207. Alvarez JA, Cohen AR. Neonatal applications of neuroendoscopy. *Neurosurg Clin N Am*. 1998;9(2):405-413.
208. Fabregas N, Craen RA. Anaesthesia for minimally invasive neurosurgery. *Best Pract Res Clin Anaesthesiol*. 2002;16(1):81-93.
209. Gaab MR, Schroeder HW. Neuroendoscopic approach to intraventricular lesions. *J Neurosurg*. 1998;88(3):496-505.
210. Teo C, Rahman S, Boop FA, Cherny B. Complications of endoscopic neurosurgery. *Childs Nerv Syst*. 1996;12(5):248-253.
211. Kato Y, Sano H, Nagahisa S, et al. Endoscope-assisted microsurgery for cerebral aneurysms. *Minim Invasive Neurosurg*. 2000;43(2):91-97.
212. Wadsworth R, Anderton JM, Vohra A. The effect of four different surgical prone positions on cardiovascular parameters in healthy volunteers. *Anaesthesia*. 1996;51(9):819-822.
213. Toyota S, Amaki Y. Hemodynamic evaluation of the prone position by transesophageal echocardiography. *J Clin Anesth*. 1998;10(1):32-35.
214. Kim KA, Wang MY. Anesthetic considerations in the treatment of cervical myelopathy. *Spine J*. 2006;6(6 Suppl):207S-211S.
215. Takizawa D, Hiraoka H, Nakamura K, Yamamoto K, Horiuchi R. Influence of the prone position on propofol pharmacokinetics. *Anaesthesia*. 2004;59(12):1250-1251.
216. Sudheer PS, Logan SW, Ateleanu B, Hall JE. Haemodynamic effects of the prone position: a comparison of propofol total intravenous and inhalation anaesthesia. *Anaesthesia*. 2006;61(2):138-141.
217. Pearce DJ. The role of posture in laminectomy. *Proc R Soc Med*. 1957;50(2):109-112.

218. Bhardwaj A, Long DM, Ducker TB, Toung TJ. Neurologic deficits after cervical laminectomy in the prone position. *J Neurosurg Anesthesiol.* 2001;13(4):314-319.
219. Nyren S, Mure M, Jacobsson H, Larsson SA, Lindahl SG. Pulmonary perfusion is more uniform in the prone than in the supine position: scintigraphy in healthy humans. *J Appl Physiol.* 1999;86(4):1135-1141.
220. Tobin A, Kelly W. Prone ventilation—it's time. *Anaesth Intensive Care.* 1999;27(2):194-201.
221. Pelosi P, Croci M, Calappi E, et al. The prone positioning during general anesthesia minimally affects respiratory mechanics while improving functional residual capacity and increasing oxygen tension. *Anesth Analg.* 1995;80(5):955-960.
222. Sawin PD, Todd MM, Traynelis VC, et al. Cervical spine motion with direct laryngoscopy and orotracheal intubation. An in vivo cinefluoroscopic study of subjects without cervical abnormality. *Anesthesiology.* 1996;85(1):26-36.
223. Watts AD, Gelb AW, Bach DB, Pelz DM. Comparison of the Bullard and Macintosh laryngoscopes for endotracheal intubation of patients with a potential cervical spine injury. *Anesthesiology.* 1997;87(6):1335-1342.
224. Sahin A, Salman MA, Erden IA, Aypar U. Upper cervical vertebrae movement during intubating laryngeal mask, fiberoptic and direct laryngoscopy: a video-fluoroscopic study. *Eur J Anaesthesiol.* 2004;21(10):819-823.
225. LeGrand SA, Hindman BJ, Dexter F, Weeks JB, Todd MM. Craniocervical motion during direct laryngoscopy and orotracheal intubation with the Macintosh and Miller blades: an in vivo cinefluoroscopic study. *Anesthesiology.* 2007;107(6):884-891.
226. Robitaille A, Williams SR, Tremblay MH, et al. Cervical spine motion during tracheal intubation with manual in-line stabilization: direct laryngoscopy versus GlideScope videolaryngoscopy. *Anesth Analg.* 2008;106(3):935-941.
227. Keller C, Brimacombe J, Keller K. Pressures exerted against the cervical vertebrae by the standard and intubating laryngeal mask airways: a randomized, controlled, cross-over study in fresh cadavers. *Anesth Analg.* 1999;89(5):1296-1300.
228. Lennarson PJ, Smith D, Todd MM, et al. Segmental cervical spine motion during orotracheal intubation of the intact and injured spine with and without external stabilization. *J Neurosurg.* 2000;92(2 Suppl):201-206.
229. Donaldson WF, III, Heil BV, Donaldson VP, Silvaggio VJ. The effect of airway maneuvers on the unstable C1-C2 segment. A cadaver study. *Spine (Phila Pa 1976).* 1997;22(11):1215-1218.
230. Breig A, Turnbull I, Hassler O. Effects of mechanical stresses on the spinal cord in cervical spondylosis. A study on fresh cadaver material. *J Neurosurg.* 1966;25(1):45-56.
231. McLeod AD, Calder I. Spinal cord injury and direct laryngoscopy—the legend lives on. *Br J Anaesth.* 2000;84(6):705-709.
232. Rau CS, Liang CL, Lui CC, Lee TC, Lu K. Quadriplegia in a patient who underwent posterior fossa surgery in the prone position. Case report. *J Neurosurg.* 2002;96(1 Suppl):101-103.
233. Crosby ET. Airway management in adults after cervical spine trauma. *Anesthesiology.* 2006;104(6):1293-1318.
234. Sagi HC, Beutler W, Carroll E, Connolly PJ. Airway complications associated with surgery on the anterior cervical spine. *Spine (Phila Pa 1976).* 2002;27(9):949-953.
235. Santoni BG, Hindman BJ, Puttlitz CM, et al. Manual in-line stabilization increases pressures applied by the laryngoscope blade during direct laryngoscopy and orotracheal intubation. *Anesthesiology.* 2009;110(1):24-31.
236. Sinha A, Agarwal A, Gaur A, Pandey CK. Oropharyngeal swelling and macroglossia after cervical spine surgery in the prone position. *J Neurosurg Anesthesiol.* 2001;13(3):237-239.
237. Wattenmaker I, Concepcion M, Hibberd P, Lipson S. Upper-airway obstruction and perioperative management of the airway in patients managed with posterior operations on the cervical spine for rheumatoid arthritis. *J Bone Joint Surg Am.* 1994;76(3):360-365.
238. Cheng MA, Sigurdson W, Tempelhoff R, Laurysen C. Visual loss after spine surgery: a survey. *Neurosurgery.* 2000;46(3):625-630.
239. Hayreh SS. Anterior ischemic optic neuropathy. *Clin Neurosci.* 1997;4(5):251-263.
240. Brown RH, Schauble JF, Miller NR. Anemia and hypotension as contributors to perioperative loss of vision. *Anesthesiology.* 1994;80(1):222-226.
241. Chang SH, Miller NR. The incidence of vision loss due to perioperative ischemic optic neuropathy associated with spine surgery: the Johns Hopkins Hospital Experience. *Spine (Phila Pa 1976).* 2005;30(11):1299-1302.
242. Dunker S, Hsu HY, Sebag J, Sadun AA. Perioperative risk factors for posterior ischemic optic neuropathy. *J Am Coll Surg.* 2002;194(6):705-710.
243. Cheng MA, Todorov A, Tempelhoff R, et al. The effect of prone positioning on intraocular pressure in anesthetized patients. *Anesthesiology.* 2001;95(6):1351-1355.
244. Hunt K, Bajekal R, Calder I, et al. Changes in intraocular pressure in anesthetized prone patients. *J Neurosurg Anesthesiol.* 2004;16(4):287-290.
245. Lam AK, Douthwaite WA. Does the change of anterior chamber depth or/and episcleral venous pressure cause intraocular pressure change in postural variation? *Optom Vis Sci.* 1997;74(8):664-667.
246. Brucculeri M, Hammel T, Harris A, Malinovsky V, Martin B. Regulation of intraocular pressure after water drinking. *J Glaucoma.* 1999;8(2):111-116.
247. Martin B, Harris A, Hammel T, Malinovsky V. Mechanism of exercise-induced ocular hypotension. *Invest Ophthalmol Vis Sci.* 1999;40(5):1011-1015.
248. Roth S. The effects of isovolumic hemodilution on ocular blood flow. *Exp Eye Res.* 1992;55(1):59-63.
249. James CB, Smith SE. The effect of posture on the intraocular pressure and pulsatile ocular blood flow in patients with non-arteritic anterior ischaemic optic neuropathy. *Eye.* 1991;5(Pt 3):309-314.
250. Yamamoto T, Kitazawa Y. Vascular pathogenesis of normal-tension glaucoma: a possible pathogenetic factor, other than intraocular pressure, of glaucomatous optic neuropathy. *Prog Retin Eye Res.* 1998;17(1):127-143.
251. Buono LM, Foroozan R. Perioperative posterior ischemic optic neuropathy: review of the literature. *Surv Ophthalmol.* 2005;50(1):15-26.
252. Schobel GA, Schmidbauer M, Millesi W, Undt G. Posterior ischemic optic neuropathy following bilateral radical neck dissection. *Int J Oral Maxillofac Surg.* 1995;24(4):283-287.
253. Aldrich MS, Alessi AG, Beck RW, Gilman S. Cortical blindness: etiology, diagnosis, and prognosis. *Ann Neurol.* 1987;21(2):149-158.
254. Kollarits CR, Lubow M, Hissong SL. Retinal strokes. I. Incidence of carotid atheromata. *JAMA.* 1972;222(10):1273-1275.
255. Huber JF, Grob D. Bilateral cortical blindness after lumbar spine surgery. A case report. *Spine (Phila Pa 1976).* 1998;23(16):1807-1809.

256. Myers MA, Hamilton SR, Bogosian AJ, Smith CH, Wagner TA. Visual loss as a complication of spine surgery. A review of 37 cases. *Spine (Phila Pa 1976)*. 1997;22(12):1325-1329.
257. Grossman W, Ward WT. Central retinal artery occlusion after scoliosis surgery with a horseshoe headrest. Case report and literature review. *Spine (Phila Pa 1976)*. 1993;18(9):1226-1228.
258. Hollenhorst R, Svien H, Benoit CF. Unilateral blindness occurring during anesthesia for neurosurgical operations. *AMA Arch Ophthalmol*. 1954;52(6):819-830.
259. Robinson MK, Halpern JI. Retinal vein occlusion. *Am Fam Physician*. 1992;45(6):2661-2666.
260. Stevens WR, Glazer PA, Kelley SD, Lietman TM, Bradford DS. Ophthalmic complications after spinal surgery. *Spine (Phila Pa 1976)*. 1997;22(12):1319-1324.
261. Lee LA, Deem S, Glenn RW, et al. Effects of anemia and hypotension on porcine optic nerve blood flow and oxygen delivery. *Anesthesiology*. 2008;108(5):864-872.
262. Abraham M, Sakhuja N, Sinha S, Rastogi S. Unilateral visual loss after cervical spine surgery. *J Neurosurg Anesthesiol*. 2003;15(4):319-322.
263. Wilson JF, Freeman SB, Breene DP. Anterior ischemic optic neuropathy causing blindness in the head and neck surgery patient. *Arch Otolaryngol Head Neck Surg*. 1991;117(11):1304-1306.
264. Farag E, Doyle DJ. Vision loss after spine surgery: a new hypothesis. *Can J Anaesth*. 2006;53(4):420.
265. Abbott MA, McLaren AD, Algie T. Intra-ocular pressure during cardiopulmonary bypass—a comparison of crystalloid and colloid priming solutions. *Anaesthesia*. 1994;49(4):343-346.
266. American Society of Anesthesiologists Task Force on Perioperative Blindness: Practice advisory for perioperative visual loss associated with spine surgery: a report by the American Society of Anesthesiologists Task Force on Perioperative Blindness. *Anesthesiology*. 2006;104(6):1319-1328.
267. Kamming D, Clarke S. Postoperative visual loss following prone spinal surgery. *Br J Anaesth*. 2005;95(2):257-260.
268. Jeon YT, Park YO, won HJ, et al. Effect of head position on postoperative chemosis after prone spinal surgery. *J Neurosurg Anesthesiol*. 2007;19(1):1-4.
269. Parkin B, Turner A, Moore E, Cook S. Bacterial keratitis in the critically ill. *Br J Ophthalmol*. 1997;81(12):1060-1063.
270. Yu YH, Chen WJ, Chen LH, Chen WC. Ischemic orbital compartment syndrome after posterior spinal surgery. *Spine (Phila Pa 1976)*. 2008;33(16):E569-E572.
271. Delattre O, Thoreux P, Liverneaux P, et al. Spinal surgery and ophthalmic complications: a French survey with review of 17 cases. *J Spinal Disord Tech*. 2007;20(4):302-307.
272. Foroglou N, Zamani A, Black P. Intra-operative MRI (iop-MR) for brain tumour surgery. *Br J Neurosurg*. 2009;23(1):14-22.
273. Mutchnick IS, Moriarty TM. Neurosurgical uses for intraprocedural magnetic resonance imaging. *Top Magn Reson Imaging*. 2005;16(5):383-395.
274. Fahlbusch R, Ganslandt O, Buchfelder M, Schott W, Nimsky C. Intraoperative magnetic resonance imaging during transsphenoidal surgery. *J Neurosurg*. 2001;95(3):381-390.
275. Brown PD, Maurer MJ, Rummans TA, et al. A prospective study of quality of life in adults with newly diagnosed high-grade gliomas: the impact of the extent of resection on quality of life and survival. *Neurosurgery*. 2005;57(3):495-504.
276. Sutherland GR, Kaibara T, Wallace C, Tomanek B, Richter M. Intraoperative assessment of aneurysm clipping using magnetic resonance angiography and diffusion-weighted imaging: technical case report. *Neurosurgery*. 2002;50(4):893-897.
277. Martin AJ, Larson PS, Ostrem JL, et al. Placement of deep brain stimulator electrodes using real-time high-field interventional magnetic resonance imaging. *Magn Reson Med*. 2005;54(5):1107-1114.
278. Jolesz FA. Future perspectives for intraoperative MRI. *Neurosurg Clin N Am*. 2005;16(1):201-213.
279. Schmitz B, Nimsky C, Wendel G, et al. Anesthesia during high-field intraoperative magnetic resonance imaging experience with 80 consecutive cases. *J Neurosurg Anesthesiol*. 2003;15(3):255-262.
280. Kanal E. Clinical utility of the American College of Radiology MR Safe Practice Guidelines. *J Magn Reson Imaging*. 2004;19(1):2-5.
281. Colletti PM. Size “H” oxygen cylinder: accidental MR projectile at 1.5 Tesla. *J Magn Reson Imaging*. 2004;19(1):141-143.
282. Tan TK, Goh J. The anaesthetist's role in the setting up of an intraoperative MR imaging facility. *Singapore Med J*. 2009;50(1):4-10.
283. Roberts GC. Post-craniotomy analgesia: current practices in British neurosurgical centres—a survey of post-craniotomy analgesic practices. *Eur J Anaesthesiol*. 2005;22(5):328-332.
284. Fabling JM, Gan TJ, Guy J, et al. Postoperative nausea and vomiting. A retrospective analysis in patients undergoing elective craniotomy. *J Neurosurg Anesthesiol*. 1997;9(4):308-312.
285. Fabling JM, Gan TJ, el Moalem HE, Warner DS, Borel CO. A randomized, double-blinded comparison of ondansetron, droperidol, and placebo for prevention of postoperative nausea and vomiting after supratentorial craniotomy. *Anesth Analg*. 2000;91(2):358-361.
286. Halloran K, Barash PG. Inside the black box: current policies and concerns with the United States Food and Drug Administration's highest drug safety warning system. *Curr Opin Anaesthesiol*. 2010;23:423-427.