

Chapter 62

REFERENCES

1. Tsen LC. Neuraxial techniques for labor analgesia should be placed in the lateral position. *Int J Obstet Anesth.* 2008;17:146-149.
2. Cook TM. Combined spinal-epidural techniques. *Anaesthesia.* 2000;55:42-64.
3. Cappiello E, O'Rourke N, Segal S, Tsen LC. A randomized trial of dural puncture epidural technique compared with the standard epidural technique for labor analgesia. *Anesth Analg.* 2008;107:1646-1651.
4. Arkoosh VA. Continuous spinal analgesia and anesthesia in obstetrics. *Reg Anesth.* 1993;18:402-405.
5. Huckaby T, Skerman JH, Hurley RJ, et al. Sensory analgesia for vaginal deliveries: a preliminary report of continuous spinal anesthesia with a 32-gauge catheter. *Reg Anesth.* 1991;16:150-153.
6. Lambert DH, Hurley RJ. Cauda equina syndrome and continuous spinal anesthesia. *Anesth Analg.* 1991;72:817-819.
7. Ginossar Y, Riley ET, Angst MS. The site of action of epidural fentanyl in humans: the difference between infusion and bolus administration. *Anesth Analg.* 2003;97:1428-1438.
8. Palot M, Chale JJ, Colladon B, et al. Anesthesia and analgesia practice patterns in French obstetrical patients. *Ann Fr Anesth Reanim.* 1998;17:210-219.
9. Bucklin BA, Hawkins JL, Anderson JR, et al. Obstetric anesthesia workforce survey: twenty-year update. *Anesthesiology.* 2005;103:645-653.
10. Shepard TH. *Catalog of Teratogenic Agents.* Baltimore, MD: Johns Hopkins University Press; 2001.
11. Briggs GG, Freeman RK, Sumner JY. *Drugs in Pregnancy and Lactation.* Philadelphia, PA: Lippincott Williams & Wilkins; 2005.
12. Tsen LC. Darwin to desflurane: anesthesia for assisted reproductive technologies. *Anesth Analg.* 2002;94(Suppl):109-114.
13. Fujinaga M, Mazze RI, Jackson EC, et al. Reproductive and teratogenic effects of sufentanil and alfentanil in Sprague-Dawley rats. *Anesth Analg.* 1988;67:166-169.
14. Fujinaga M, Mazze RI, Baden JM, et al. Rat whole embryo culture: an in vitro model for testing nitrous oxide teratogenicity. *Anesthesiology.* 1988;69:401-404.
15. Fujinaga M, Baden JM, Mazze RI. Developmental toxicity of nondepolarizing muscle relaxants in cultured rat embryos. *Anesthesiology.* 1992;76:999-1003.
16. Fujinaga M, Mazze RI. Reproductive and teratogenic effects of lidocaine in Sprague-Dawley rats. *Anesthesiology.* 1986;65:626-632.
17. Mazze RI, Fujinaga M, Rice SA, et al. Reproductive and teratogenic effects of nitrous oxide, halothane, isoflurane, and enflurane in Sprague-Dawley rats. *Anesthesiology.* 1986;64:339-344.
18. Mazze RI, Fujinaga M, Baden JM. Halothane prevents nitrous oxide teratogenicity in Sprague-Dawley rats; folic acid does not. *Teratology.* 1988;38:121-127.
19. Hawkins JL, Chang J, Palmer S, Gibbs CP, Callaghan WM. Anesthesia-related maternal mortality in the United States: 1979-2002. *Obstet Gynecol* 2011;117(1):69-74.
20. Cooper GM, McClure JH. Anaesthesia chapter from Saving mothers' lives; reviewing maternal deaths to make pregnancy safer. *Br J Anaesth.* 2008;100:17-22.
21. Kodali BS, Chandrasekhar S, Bulich LN, Topulos GP, Datta S. Airway changes during labor and delivery. *Anesthesiology.* 2008;108:357-362.
22. Endler GC, Mariona FG, Sokol RJ, et al. Anesthesia-related maternal mortality in Michigan, 1972 to 1984. *Am J Obstet Gynecol.* 1988;159:187-193.
23. Tsen LC, Pitner R, Camann WR. General anesthesia for cesarean section at a tertiary care hospital 1990-1995: indications and implications. *Int J Obstet Anesth.* 1998;7:147-152.
24. Mhyre JM, Riesner MN, Polley LS, Naughton NN. A series of anesthesia-related maternal deaths in Michigan, 1985-2003. *Anesthesiology.* 2007;106:1096-1104.
25. Marx GF. Aortocaval compression syndrome: its 50 year history. *Int J Obstet Anesth.* 1992;1:60-64.
26. Ramanathan J, Bennett K. Pre-eclampsia: fluids, drugs, and anesthetic management. *Anesthesiol Clin North America.* 2003;21:145-163.
27. Tsen LC, Schultz R, Martin R, et al. Intrathecal low-dose bupivacaine versus lidocaine for in vitro fertilization procedures. *Reg Anesth. Pain Med.* 2001;26:52-56.
28. Martin R, Tsen LC, Tzeng G, Hornstein MD, Datta S. Anesthesia for in vitro fertilization: the addition of fentanyl to 1.5% lidocaine. *Anesth Analg.* 1999;88:523-526.
29. Kauppila A, Leinonen P, Vihko R, et al. Metoclopramide-induced hyperprolactinemia impairs ovarian follicle maturation and corpus luteum function in women. *J Clin Endocrinol Metab.* 1982;54:955-960.
30. Practice Committee of American Society for Reproductive Medicine. Ovarian hyperstimulation syndrome. *Fertil Steril.* 2008;90(Suppl):S188-S193.

31. ACOG practice bulletin. Cervical insufficiency. *Int J Gynaecol Obstet.* 2004;85:81-89.
32. Lavoie A, Guay J. Anesthetic dose neuraxial blockade increases the success rate of external fetal version: a meta-analysis. *Can J Anaesth.* 2010;57:408-414.
33. Cherayil G, Feinberg B, Robinson J, et al. Central neuraxial blockade promotes external cephalic version success after a failed attempt. *Anesth Analg.* 2002;94:1589-1592.
34. Deprest JA, Devlieger R, Srisupundit K, et al. Fetal surgery is a clinical reality. *Semin Fetal Neonatal Med.* 2010;15:58-67.
35. Johnson MD, Birnbach DJ, Burchman C, et al. Fetal surgery and general anesthesia: a case report and review. *J Clin Anesth.* 1989;1:363-367.
36. McCullagh P. Fetal sentience and fetal surgery. In: Adams AP, Cashman JN, eds. *Recent Advances in Anaesthesia and Analgesia.* Vol. 20. Edinburgh, UK: Churchill Livingstone; 1998:107-121.
37. Lee SJ, Ralston HJ, Drey EA, et al. Fetal pain: a systematic multidisciplinary review of the evidence. *JAMA.* 2005;294:947-954.
38. Strumper D, Durieux ME, Gogarten W, et al. Fetal plasma concentrations after intraamniotic sufentanil in chronically instrumented pregnant sheep. *Anesthesiology.* 2003;98:1400-1406.
39. Zhang J, Troendle JF, Yancey MK. Reassessing the labor curve in nulliparous women. *Am J Obstet Gynecol.* 2002;187:824-828.
40. Eltzschig HK, Lieberman ES, Camann WR. Regional anesthesia and analgesia for labor and delivery. *N Engl J Med.* 2003;348:319-332.
41. Tsen LC, Thomas J, Segal S, et al. Transcutaneous electrical nerve stimulation does not augment combined spinal epidural labour analgesia. *Can J Anaesth.* 2000;47:38-42.
42. Tsen LC, Thomas J, Segal S, et al. Transcutaneous electrical nerve stimulation does not augment epidural labor analgesia. *J Clin Anesth.* 2001;13:571-575.
43. Practice guidelines for obstetric anesthesia: an updated report by the American Society of Anesthesiologists Task Force on Obstetric Anesthesia. *Anesthesiology.* 2007;106:843-863.
44. Santos AC, DeArmas PI. Systemic toxicity of levobupivacaine, bupivacaine, and ropivacaine during continuous intravenous infusion to nonpregnant and pregnant ewes. *Anesthesiology.* 2001;95:1256-1264.
45. Ohmura S, Kawada M, Ohta T, et al. Systemic toxicity and resuscitation in bupivacaine-, levobupivacaine-, or ropivacaine-infused rats. *Anesth Analg.* 2001;93:743-748.
46. Camann WR, Denney RA, Holby ED, et al. A comparison of intrathecal, epidural, and intravenous sufentanil for labor analgesia. *Anesthesiology.* 1992;77:884-887.
47. Celleno D, Capogna G. Epidural fentanyl plus bupivacaine 0.125 per cent for labour: analgesic effects. *Can J Anaesth.* 1988;35:375-378.
48. van der Vyver M, Halpern S, Joseph G. Patient-controlled epidural analgesia versus continuous infusion for labour analgesia: a meta-analysis. *Br J Anaesth.* 2002;89:459-465.
49. Wong CA, Scavone BM, Peaceman AM, et al. The risk of cesarean delivery with neuraxial analgesia given early versus late in labor. *N Engl J Med.* 2005;352:655-665.
50. Tsen LC, Thue B, Datta S, et al. Is combined spinal-epidural analgesia associated with more rapid cervical dilation in nulliparous patients when compared with conventional epidural analgesia? *Anesthesiology.* 1999;91:920-925.
51. Thomas JA, Pan PH, Harris LC, Owen MD, D'Angelo R. Dural puncture with a 27-gauge Whitacre needle as part of a combined spinal-epidural technique does not improve labor epidural catheter function. *Anesthesiology.* 2005;103:1046-1051.
52. Halpern SH, Leighton BL, Ohlsson A, et al. Effect of epidural vs parenteral opioid analgesia on the progress of labor: a meta-analysis. *JAMA.* 1998;280:2105-2110.
53. American College of Obstetricians and Gynecologists Task Force on Cesarean Delivery Rates. *Evaluation of Cesarean Delivery.* Washington, DC: American College of Obstetricians and Gynecologists; 2000.
54. Segal S, Su M, Gilbert P. The effect of a rapid change in availability of epidural analgesia on the cesarean delivery rate: a meta-analysis. *Am J Obstet Gynecol.* 2000;183:974-978.
55. Tsen LC, Boosalis P, Segal S, et al. Hemodynamic effects of simultaneous administration of intravenous ephedrine and spinal anesthesia for cesarean delivery. *J Clin Anesth.* 2000;12:378-382.
56. Corke BC, Datta S, Ostheimer GW, et al. Spinal anaesthesia for cesarean section. The influence of hypotension on neonatal outcome. *Anaesthesia.* 1982;37:658-662.
57. Datta S, Alper MH, Ostheimer GW, et al. Method of ephedrine administration and nausea and hypotension during spinal anesthesia for cesarean section. *Anesthesiology.* 1982;56:68-70.
58. Turnbull DK, Shepherd DB. Post-dural puncture headache: pathogenesis, prevention and treatment. *Br J Anaesth.* 2003;91:718-729.
59. Jones RJ. The role of recumbency in the prevention and treatment of postspinal headache. *Anesth Analg.* 1974;53:788-796.
60. Abouleish E, Vega S, Blendinger I, et al. Long-term follow-up of epidural blood patch. *Anesth Analg.* 1975;54:459-463.
61. Holmstrom B, Rawal N, Axelsson K, et al. Risk of catheter migration during combined spinal epidural block: percutaneous epiduroscopy study. *Anesth Analg.* 1995;80:747-753.
62. Albright GA, Forster RM. The safety and efficacy of combined spinal and epidural analgesia/anesthesia (6,002 blocks) in a community hospital. *Reg Anesth Pain Med.* 1999;24:117-125.
63. Hamilton BE, Martin JA, Ventura SJ. *National Vital Statistics Reports. Births: Preliminary Data for 2007.* Washington, DC: US Department of Health and Human Services, National Center for Health Statistics, Center for Disease Control and Prevention; 2009;57:1-23.
64. Betran AP, Merialdi M, Lauer JA, et al. Rates of caesarean section: analysis of global, regional and national estimates. *Paediatr Perinat Epidemiol.* 2007;21:98-113.
65. Balki M, Ronayne M, Davies S, et al. Minimum oxytocin dose requirement after cesarean delivery for labor arrest. *Obstet Gynecol.* 2006;107:45-50.
66. Carvalho JC, Balki M, Kingdom J, Windrim R. Oxytocin requirements at elective cesarean delivery: a dose-finding study. *Obstet Gynecol.* 2004;104:1005-1010.
67. Tsen LC, Balki M. Oxytocin protocols during cesarean delivery: time to acknowledge the risk/benefit ratio? *Int J Obstet Anesth.* 2010;19(3):243-245.
68. Secher NJ, Arnsbo P, Wallin L. Haemodynamic effects of oxytocin (syntocinon) and methyl ergometrine (methergin) on the systemic and pulmonary circulations of pregnant anaesthetized women. *Acta Obstet Gynecol Scand.* 1978;57:97-103.
69. Heytens L, Camu F. Pulmonary edema during cesarean section related to the use of oxytocic drugs. *Acta Anaesthesiol Belg.* 1984;35:155-164.
70. Park EH, Sachs blood pressure. Postpartum hemorrhage and other problems of the third stage. In: Gonik B, ed. *High Risk*

Pregnancy: Management Options. London, UK: WB Saunders; 1999:1231-1246.

71. Palmer CM, Emerson S, Volgoropolous D, et al. Dose-response relationship of intrathecal morphine for post-cesarean analgesia. *Anesthesiology*. 1999;90:437-444.
72. Abouleish EI. Epinephrine improves the quality of spinal hyperbaric bupivacaine for cesarean section. *Anesth Analg*. 1987;66:395-400.
73. Datta S, Lambert DH, Gregus J, et al. Differential sensitivities of mammalian nerve fibers during pregnancy. *Anesth Analg*. 1983;62:1070-1072.
74. Bromage PR. Continuous lumbar epidural analgesia for obstetrics. *Can Med Assoc J*. 1961;85:1136-1140.
75. Philipson EH, Kuhnert BR, Syracuse CD. Fetal acidosis, 2-chloroprocaine, and epidural anesthesia for cesarean section. *Am J Obstet Gynecol*. 1985;151:322-324.
76. Camann WR, Hartigan PM, Gilbertson LI, et al. Chloroprocaine antagonism of epidural opioid analgesia: a receptor-specific phenomenon? *Anesthesiology*. 1990;73:860-863.
77. Stevens RA, Urmey WF, Urquhart BL, et al. Back pain after epidural anesthesia with chloroprocaine. *Anesthesiology*. 1993;78:492-497.
78. Stevens RA, Chester WL, Schubert A, et al. pH-adjustment of 2-chloroprocaine quickens the onset of epidural anaesthesia. *Can J Anaesth*. 1989;36:515-518.
79. Peterfreund RA, Datta S, Ostheimer GW. pH adjustment of local anesthetic solutions with sodium bicarbonate: laboratory evaluation of alkalization and precipitation. *Reg Anesth*. 1989;14:265-270.
80. Portnoy D, Vadhera RB. Mechanisms and management of an incomplete epidural block for cesarean section. *Anesthesiol Clin North America*. 2003;21:39-57.
81. Tsen LC, Pitner R, Camann WR. General anesthesia at a tertiary care hospital 1990-1995: indications and implications. *Int J Obstet Anesth*. 1998;7:147-152.
82. Aya AG, Mangin R, Vialles N, et al. Patients with severe preeclampsia experience less hypotension during spinal anesthesia for elective cesarean delivery than healthy parturients: a prospective cohort comparison. *Anesth Analg*. 2003;97:867-872.
83. Santos AC, Birnbach DJ. Spinal anesthesia in the parturient with severe preeclampsia: time for reconsideration. *Anesth Analg*. 2003;97:621-622.
84. Chestnut DH, Dewan DM, Redick LF, et al. Anesthetic management for obstetric hysterectomy: a multi-institutional study. *Anesthesiology*. 1989;70:607-610.
85. Park GE, Hauch MA, Curlin F, et al. The effects of varying volumes of crystalloid administration before cesarean delivery on maternal hemodynamics and colloid osmotic pressure. *Anesth Analg*. 1996;83:299-303.
86. Lee A, Ngan Kee WD, Gin T. A quantitative, systematic review of randomized controlled trials of ephedrine versus phenylephrine for the management of hypotension during spinal anesthesia for cesarean delivery. *Anesth Analg*. 2002;94:920-926.
87. Ueyama H, He YL, Tanigami H, et al. Effects of crystalloid and colloid preload on blood volume in the parturient undergoing spinal anesthesia for elective cesarean section. *Anesthesiology*. 1999;91:1571-1576.
88. Loughrey JP, Yao N, Datta S, et al. Hemodynamic effects of spinal anesthesia and simultaneous intravenous bolus of combined phenylephrine and ephedrine versus ephedrine for cesarean delivery. *Int J Obstet Anesth*. 2005;14:43-47.
89. Mercier FJ, Riley ET, Frederickson WL, et al. Phenylephrine added to prophylactic ephedrine infusion during spinal anesthesia for elective cesarean section. *Anesthesiology*. 2001;95:668-674.
90. Ngan Kee WD, Lee A, Khaw KS, Ng FF, Karmakar MK, Gin T. A randomized double-blinded comparison of phenylephrine and ephedrine infusion combinations to maintain blood pressure during spinal anesthesia for cesarean delivery: the effects on fetal acid-base status and hemodynamic control. *Anesth Analg*. 2008;107:1295-1302.
91. Balki M, Carvalho JC. Intraoperative nausea and vomiting during cesarean section under regional anesthesia. *Int J Obstet Anesth*. 2005;14:230-241.
92. Szarvas S, Harmon D, Murphy D. Neuraxial opioid-induced pruritus: a review. *J Clin Anesth*. 2003;15:234-239.
93. Kranke P, Eberhart LH, Roewer N, et al. Pharmacological treatment of postoperative shivering: a quantitative systematic review of randomized controlled trials. *Anesth Analg*. 2002;94:453-460.
94. Ross BK. ASA closed claims in obstetrics: lessons learned. *Anesthesiol Clin North America*. 2003;21:183-197.
95. Royal College of Obstetricians and Gynaecologists. *Why Mothers Die 1997-1999: The Confidential Enquiries into Maternal Deaths in the United Kingdom*. London, UK: Royal College of Obstetricians and Gynaecologists; 2001.
96. Malan TP Jr, Johnson MD. The difficult airway in obstetric anesthesia: techniques for airway management and the role of regional anesthesia. *J Clin Anesth*. 1988;1:104-111.
97. Practice guidelines for management of the difficult airway: an updated report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. *Anesthesiology*. 2003;98:1269-1277.
98. Han TH, Brimacombe J, Lee EJ, et al. The laryngeal mask airway is effective (and probably safe) in selected healthy parturients for elective cesarean section: a prospective study of 1067 cases. *Can J Anaesth*. 2001;48:1117-1121.
99. James CF, Modell JH, Gibbs CP, et al. Pulmonary aspiration—effects of volume and pH in the rat. *Anesth Analg*. 1984;63:665-668.
100. Johnston JR, Moore J, McCaughey W, et al. Use of cimetidine as an oral antacid in obstetric anesthesia. *Anesth Analg*. 1983;62:720-726.
101. Murphy DF, Nally B, Gardiner J, et al. Effect of metoclopramide on gastric emptying before elective and emergency cesarean section. *Br J Anaesth*. 1984;56:1113-1116.
102. Norris MC, Dewan DM. Preoxygenation for cesarean section: a comparison of two techniques. *Anesthesiology*. 1985;62:827-829.
103. Khaw KS, Ngan Kee WD. Fetal effects of maternal supplementary oxygen during cesarean section. *Curr Opin Anaesthesiol*. 2004;309-313.
104. Dogru K, Yildiz K, Dalgic H, et al. Inhibitory effects of desflurane and sevoflurane on contractions of isolated gravid rat myometrium under oxytocin stimulation. *Acta Anaesthesiol Scand*. 2003;47:472-474.
105. Mattingly JE, D'Alessio J, Ramanathan J. Effects of obstetric analgesics and anesthetics on the neonate: a review. *Paediatr Drugs*. 2003;5:615-627.
106. Datta S, Ostheimer GW, Weiss JB, et al. Neonatal effect of prolonged anesthetic induction for cesarean section. *Obstet Gynecol*. 1981;58:331-335.

107. Palmer CM, Nogami WM, Van Maren G, et al. Postcesarean epidural morphine: a dose-response study. *Anesth Analg*. 2000;90:887-891.
108. Carvalho B, Riley E, Cohen SE, et al. Single-dose, sustained-release epidural morphine in the management of postoperative pain after elective cesarean delivery: results of a multicenter randomized controlled study. *Anesth Analg*. 2005;100:1150-1158.
109. Kotelko DM, Dailey PA, Shnider SM, et al. Epidural morphine analgesia after cesarean delivery. *Obstet Gynecol*. 1984;63:409-413.
110. Wang BC, Hillman DE, Spielholz NI, et al. Chronic neurologic deficits and Nesacaine-CE--an effect of the anesthetic, 2-chloroprocaine, or the antioxidant, sodium bisulfite? *Anesth Analg*. 1984;63:445-447.
111. Fischer RL, Lubenow TR, Liceaga A, et al. Comparison of continuous epidural infusion of fentanyl-bupivacaine and morphine-bupivacaine in management of postoperative pain. *Anesth Analg*. 1988;67:559-563.
112. Lavand'homme P. Chronic pain after vaginal and cesarean delivery: a reality questioning our daily practice of obstetric anesthesia. *Int J Obstet Anesth*. 2010;19:1-2.
113. Gonik B. Intensive care monitoring of the critically ill pregnant patient. In: Creasy RK, Resnik R, eds. *Maternal-Fetal Medicine*. Philadelphia, PA: WB Saunders; 1994:865-890.
114. *Why Mothers Die 1997-1999: Confidential Enquiries into Maternal Deaths in the United Kingdom*. London, UK: Department of Health, HMSO; 2001:36-37.
115. Fuller AJ, Carvalho B, Brummel C, et al. Epidural anesthesia for elective cesarean delivery with intraoperative arterial occlusion balloon catheter placement. *Anesth Analg*. 2006;102:585-587.
116. Georgiou C. Balloon tamponade in the management of postpartum haemorrhage: a review. *BJOG*. 2009;116:748-757.
117. Practice guidelines for blood component therapy: a report by the American Society of Anesthesiologists Task Force on Blood Component Therapy. *Anesthesiology*. 1996;84:732-747.
118. Sibai BM. Diagnosis and management of gestational hypertension and preeclampsia. *Obstet Gynecol*. 2003;102:181-192.
119. Jouppila P, Jouppila R, Hollmen A, et al. Lumbar epidural analgesia to improve intervillous blood flow during labor in severe preeclampsia. *Obstet Gynecol*. 1982;59:158-161.
120. Dyer RA, Piercy JL, Reed AR, et al. Hemodynamic changes associated with spinal anesthesia for cesarean delivery in severe preeclampsia. *Anesthesiology*. 2008;108:802-811.
121. VanWijk MJ, Boer K, Berckmans RJ, et al. Enhanced coagulation activation in preeclampsia: the role of APC resistance, microparticles and other plasma constituents. *Thromb Haemost*. 2002;88:415-420.
122. Wallace DH, Leveno KJ, Cunningham FG, et al. Randomized comparison of general and regional anesthesia for cesarean delivery in pregnancies complicated by severe preeclampsia. *Obstet Gynecol*. 1995;86:193-199.
123. Izi B, Riha RL, Martin SE, et al. The upper airway in pregnancy and pre-eclampsia. *Am J Respir Crit Care Med*. 2003;167:137-140.
124. Clinical competence in hemodynamic monitoring. A statement for physicians from the ACP/ACC/AHA Task Force on Clinical Privileges in Cardiology. *Circulation*. 1990;81:2036-2040.
125. Sibbald WJ, Keenan SP. Show me the evidence: a critical appraisal of the Pulmonary Artery Catheter Consensus Conference and other musings on how critical care practitioners need to improve the way we conduct business. *Crit Care Med*. 1997;25:2060-2063.
126. Connors AF Jr, Speroff T, Dawson NV, et al. The effectiveness of right heart catheterization in the initial care of critically ill patients. SUPPORT Investigators. *JAMA*. 1996;276:889-897.
127. Newsome LR, Bramwell RS, Curling PE. Severe preeclampsia: hemodynamic effects of lumbar epidural anesthesia. *Anesth Analg*. 1986;65:31-36.
128. Sise MJ, Hollingsworth P, Brimm JE, et al. Complications of the flow-directed pulmonary artery catheter: a prospective analysis in 219 patients. *Crit Care Med*. 1981;9:315-318.
129. Madden BP. Pulmonary hypertension and pregnancy. *Int J Obstet Anesth*. 2009;18:156-164.
130. Benedetti TJ, Kates R, Williams V. Hemodynamic observations in severe preeclampsia complicated by pulmonary edema. *Am J Obstet Gynecol*. 1985;152:330-334.
131. Clark SL, Cotton DB. Clinical indications for pulmonary artery catheterization in the patient with severe preeclampsia. *Am J Obstet Gynecol*. 1988;158:453-458.
132. Belfort MA, Mares A, Saade G, et al. Two-dimensional echocardiography and Doppler ultrasound in managing obstetric patients. *Obstet Gynecol*. 1997;90:326-330.
133. Fischbach F. *A Manual of Laboratory and Diagnostic Tests*. 5th ed. Philadelphia, PA: JB Lippincott; 1996:362-363.
134. Gao S, Escalante C. Venous thromboembolism and malignancy. *Exp Rev Anticancer Ther*. 2004;4:303-320.
135. Schafer AI, Levine MN, Konkle BA, et al. Thrombotic disorders: diagnosis and treatment. *Hematology Am Soc Hematol Educ Program*. 2003:520-539.
136. Fillingim RB, King CD, Ribeiro-Dasilva MC, et al. Sex, gender, and pain: a review of recent clinical and experimental findings. *J Pain*. 2009;10:447-485.
137. Dawson-Basoa M, Gintzler AR. Nociceptin (Orphanin FQ) abolishes gestational and ovarian sex steroid-induced antinociception and induces hyperalgesia. *Brain Res*. 1997;750:48-52.
138. Ratka A, Simpkins JW. Effects of estradiol and progesterone on the sensitivity to pain and on morphine-induced antinociception in female rats. *Horm Behav*. 1991;25:217-228.
139. Weiland NG, Wise PM. Estrogen and progesterone regulate opiate receptor densities in multiple brain regions. *Endocrinology*. 1990;126:804-808.
140. Tsen LC, Natale M, Datta S, et al. Can estrogen influence the response to noxious stimuli? *J Clin Anesth*. 2001;13:118-121.
141. Tong C, Ma W, Shin SW, et al. Uterine cervical distension induces cFos expression in deep dorsal horn neurons of the rat spinal cord. *Anesthesiology*. 2003;99:205-211.
142. Pleym H, Spigset O, Kharasch ED, et al. Gender differences in drug effects: implications for anesthesiologists. *Acta Anaesthesiol Scand*. 2003;47:241-259.
143. Wilson K, Oram M, Horth CE, et al. The influence of the menstrual cycle on the metabolism and clearance of methaqualone. *Br J Clin Pharmacol*. 1982;14:333-339.
144. Kirkwood C, Moore A, Hayes P, et al. Influence of menstrual cycle and gender on alprazolam pharmacokinetics. *Clin Pharmacol Ther*. 1991;50:404-409.
145. Kashuba AD, Bertino JS Jr, Rocci ML Jr, et al. Quantification of 3-month intraindividual variability and the influence of sex and menstrual cycle phase on CYP3A activity as measured by phenotyping with intravenous midazolam. *Clin Pharmacol Ther*. 1998;64:269-277.
146. Gustavson LE, Benet LZ. Menopause: pharmacodynamics and pharmacokinetics. *Exp Gerontol*. 1994;29:437-444.

147. Hoymork SC, Raeder J. Why do women wake up faster than men from propofol anaesthesia? *Br J Anaesth.* 2005;95:627-633.
148. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA.* 2010;303:235-241.
149. King JC, Sherline DM. Paracervical and pudendal block. *Clin Obstet Gynecol.* 1981;24:587-595.
150. Alp E, Bijl D, Bleichrodt RP, et al. Surgical smoke and infection control. *J Hosp Infect.* 2006;62:1-5.
151. Ray BK, Vallejo MC, Creinin MD, et al. Amniotic fluid embolism with second trimester pregnancy termination: a case report. *Can J Anaesth.* 2004;51:139-144.
152. Maslow AD, Breen TW, Sarna MC, et al. Prevalence of coagulation abnormalities associated with intrauterine fetal death. *Can J Anaesth.* 1996;43:1237-1243.
153. Kleiner GJ, Greston WM. Current concepts of defibrination in the pregnant woman. *J Reprod Med.* 1976;17:309-317.
154. Sarwar N, Newlands ES, Seckl MJ. Gestational trophoblastic neoplasia: the management of relapsing patients and other recent advances. *Curr Oncol Rep.* 2004;6:476-482.
155. Yildiz K, Dogru K, Dalgic H, et al. Inhibitory effects of desflurane and sevoflurane on oxytocin-induced contractions of isolated pregnant human myometrium. *Acta Anaesthesiol Scand.* 2005;49:1355-1359.
156. Hirvonen EA, Nuutinen LS, Kauko M. Hemodynamic changes due to Trendelenburg positioning and pneumoperitoneum during laparoscopic hysterectomy. *Acta Anaesthesiol Scand.* 1995;39:949-955.
157. Irvin W, Andersen W, Taylor P, et al. Minimizing the risk of neurologic injury in gynecologic surgery. *Obstet Gynecol.* 2004;103:374-382.
158. Litwiller JP, Wells RE Jr, Halliwill JR, et al. Effect of lithotomy positions on strain of the obturator and lateral femoral cutaneous nerves. *Clin Anat.* 2004;17:45-49.
159. Practice advisory for the prevention of perioperative peripheral neuropathies: a report by the American Society of Anesthesiologists Task Force on Prevention of Perioperative Peripheral Neuropathies. *Anesthesiology.* 2000;92:1168-1182.
160. Bradley LD. Complications in hysteroscopy: prevention, treatment and legal risk. *Curr Opin Obstet Gynecol.* 2002;14:409-415.
161. Verbalis JG. Adaptation to acute and chronic hyponatremia: implications for symptomatology, diagnosis, and therapy. *Semin Nephrol* 1998;18:3-19.
162. Stoff-Khalili MA, Dall P, Curiel DT. From gene therapy to virotherapy for ovarian cancer. *Minerva Ginecol.* 2004;56:503-514.
163. Marana R, Catalano GF, Muzii L. Salpingoscopy. *Curr Opin Obstet Gynecol.* 2003;15:333-336.
164. Ramirez NC, Lawrence WD, Ginsburg KA. Ectopic pregnancy. A recent five-year study and review of the last 50 years' literature. *J Reprod Med.* 1996;41:733-740.
165. Rehm M, Orth V, Kreimeier U, et al. Changes in intravascular volume during acute normovolemic hemodilution and intraoperative retransfusion in patients with radical hysterectomy. *Anesthesiology.* 2000;92:657-664.
166. Esler MD, Douglas MJ. Planning for hemorrhage. Steps an anesthesiologist can take to limit and treat hemorrhage in the obstetric patient. *Anesthesiol Clin North America.* 2003;21:127-144, vii.
167. Hansen E, Bechmann V, Altmeyen J. Intraoperative blood salvage in cancer surgery: safe and effective? *Transfus Apher Sci.* 2002;27:153-157.
168. Moiniche S, Mikkelsen S, Wetterslev J, et al. A qualitative systematic review of incisional local anaesthesia for postoperative pain relief after abdominal operations. *Br J Anaesth.* 1998;81:377-383.
169. Medeiros L, Fachel J, Garry R, et al. Laparoscopy versus laparotomy for benign ovarian tumours. *Cochrane Database Syst Rev.* 2005;CD004751.
170. Johnson N, Barlow D, Lethaby A, et al. Methods of hysterectomy: systematic review and meta-analysis of randomised controlled trials. *BMJ.* 2005;330:1478.
171. Zapico A, Fuentes P, Grassa A, et al. Laparoscopic-assisted vaginal hysterectomy versus abdominal hysterectomy in stages I and II endometrial cancer. Operating data, follow up and survival. *Gynecol Oncol.* 2005;98:222-227.
172. Krivak TC, Elkas JC, Rose GS, et al. The utility of hand-assisted laparoscopy in ovarian cancer. *Gynecol Oncol.* 2005;96:72-76.
173. Holzheimer RG. Laparoscopic procedures as a risk factor of deep venous thrombosis, superficial ascending thrombophlebitis and pulmonary embolism—case report and review of the literature. *Eur J Med Res.* 2004;9:417-422.