

Chapter 64

REFERENCES

1. Tiret L, Nivoche Y, Hatton F, et al. Complications related to anaesthesia in infants and children. *Br J Anaesth.* 1988;61:263.
2. Murat I, Constant I, Maud'huy H. Perioperative anaesthetic morbidity in children: a database of 24,165 anaesthetics over a 30-month period. *Paediatr Anaesth.* 2004;14:158.
3. Tait AR, Malviya S. Anesthesia for the child with an upper respiratory tract infection: still a dilemma? *Anesth Analg.* 2005;100:59.
4. Warner MA, Warner ME, Warner DO, et al. Perioperative pulmonary aspiration in infants and children. *Anesthesiology.* 1999;90:66.
5. Keller C, Brimacombe J, Bittersohl J, et al. Aspiration and the laryngeal mask airway: three cases and a review of the literature. *Br J Anaesth.* 2004;93:579.
6. Cote CJ, Zaslavsky A, Downes JJ, et al. Postoperative apnea in former preterm infants after inguinal herniorrhaphy. *Anesthesiology.* 1995;82:809.
7. Welborn LG, Hannallah RS, Fink R, et al. High-dose caffeine suppresses postoperative apnea in former preterm infants. *Anesthesiology.* 1989;71:347.
8. Welborn LG, Rice LJ, Hannallah RS, et al. Postoperative apnea in former preterm infants: prospective comparison of spinal and general anesthesia. *Anesthesiology.* 1990;72:838.
9. William JM, Stoddart PA, Williams SA, et al. Post-operative recovery after inguinal herniotomy in ex-premature infants: comparison between sevoflurane and spinal anaesthesia. *Br J Anaesth.* 2001;86:366.
10. Murray JP, Geiduschek JM, Caplan RA, et al. A comparison of pediatric and adult anesthesia closed malpractice claims. *Anesthesiology.* 1993;78:461.
11. Murray JP, Geiduschek JM, Ramamoorthy C, et al. Anesthesia-related cardiac arrest in children: initial findings of the Pediatric Perioperative Cardiac Arrest (POCA) Registry. *Anesthesiology.* 2000;93:6.
12. Bhananker SM, Ramamoorthy C, Geiduschek JM, et al. Anesthesia-related cardiac arrests in children: update from Pediatric Perioperative Cardiac Arrest Registry. *Anesth Analg.* 2007;105: 344-350.
13. Ramamoorthy C, Haberkern CM, Bhananker SM, et al. Anesthesia-related cardiac arrest in children with heart disease: data from the Pediatric Perioperative Cardiac Arrest (POCA) registry. *Anesth Analg.* 2010;110:1376-1382.
14. Larach MG, Rosenberg H, Gronert GA, et al. Hyperkalemic cardiac arrest during anesthesia in infants and children with occult myopathies. *Clin Pediatr.* 1997;36:9.
15. Istaphanous GK, Loepke AW. General anesthetics and the developing brain. *Curr Opin Anaesthesiol.* 2009;22:368-373.
16. Bartels M, Althoff RR, Boomsma DI. Anesthesia and cognitive performance in children: no evidence for a causal relationship. *Twin Res Hum Genet.* 2009;12:246-253.
17. Davidson AJ, McCann ME, Morton NS, et al. Anesthesia and outcome after neonatal surgery: the role for randomized trials. *Anesthesiology.* 2008;109:941-944.
18. Keenan RL, Shapiro JH, Kane FR, et al. Bradycardia during anesthesia in infants. An epidemiologic study. *Anesthesiology.* 1994;80:976.
19. Mamie C, Habre W, Delhumeau C, et al. Incidence and risk factors of perioperative respiratory adverse events in children undergoing elective surgery. *Paediatr Anaesth.* 2004;14:218.
20. Hackel A, Badgwell JM, Binding RR, et al. Guidelines for the pediatric perioperative anesthesia environment. *Pediatrics.* 1999;103:512.
21. Stevenson GW, Tobin MG, Horn BJ, et al. The effect of circuit compliance on delivered ventilation with use of an adult circle system for time cycled volume controlled ventilation using an infant lung model. *Paediatr Anaesth.* 1998;8:139.
22. Kain ZN, Mayes LC, Wang SM, et al. Parental presence and a sedative premedicant for children undergoing surgery: a hierarchical study. *Anesthesiology.* 2000;92:939.
23. Chorney JM, Torrey C, Blount R, et al. Healthcare provider and parent behavior and children's coping and distress at anesthesia induction. *Anesthesiology.* 2009;111:1290-1296.
24. Al-alami AA, Zestos MM, Baraka AS. Pediatric laryngospasm: prevention and treatment. *Curr Opin Anaesthesiol.* 2009;22: 388-395.
25. Rapp HJ, Altenmueller CA, Waschke C. Neuromuscular recovery following rocuronium bromide single dose in infants. *Paediatr Anaesth.* 2004;14:329.
26. Flick RP, Wilder RT, Pieper SF, et al. Risk factors for laryngospasm in children during general anesthesia. *Paediatr Anaesth.* 2008;18:289-296.
27. von Ungern-Sternberg BS, Boda K, Schwab C, et al. Laryngeal mask airway is associated with an increased incidence of adverse respiratory events in children with recent upper respiratory tract infections. *Anesthesiology.* 2007;107: 714-719.
28. Khine HH, Corrdry DH, Kettrick RG, et al. Comparison of cuffed and uncuffed endotracheal tubes in young children during general anesthesia. *Anesthesiology.* 1997;86:627.

29. Vergheze ST, Hannallah RS, Slack MC, et al. Auscultation of bilateral breath sounds does not rule out endobronchial intubation in children. *Anesth Analg*. 2004;99:56.
30. Bray RJ. Propofol infusion syndrome in children. *Paediatr Anaesth*. 1998;8:491.
31. Leelanukrom R, Cunliffe M. Intraoperative fluid and glucose management in children. *Paediatr Anaesth*. 2000;10:353.
32. Bosenberg AT, Raw R, Boezaart AP. Surface mapping of peripheral nerves in children with a nerve stimulator. *Paediatr Anaesth*. 2002;12:398.
33. Flandin-Blety C, Barrier G. Accidents following extradural analgesia in children. The results of a retrospective study. *Paediatr Anaesth*. 1995;5:41.
34. Krane E, Dalens B, Murat I, et al. Epidural catheters in anesthetized patients. *Reg Anesth Pain Med*. 1998;23:433.
35. Berde C. Convulsions associated with pediatric regional anesthesia [editorial]. *Anesth Analg*. 1992;75:164.
36. Tanaka M, Nishikawa T. The efficacy of a simulated intravascular test dose in sevoflurane-anesthetized children: a dose-response study. *Anesth Analg*. 1999;89:632.
37. Giaufre E, Dalens B, Gombert A. Epidemiology and morbidity of regional anesthesia in children: a one-year prospective survey of the French-Language Society of Pediatric Anesthesiologists. *Anesth Analg*. 1996;83:904.
38. Vergheze ST, Mostello LA, Patel RI, et al. Testing anal sphincter tone predicts the effectiveness of caudal analgesia in children. *Anesth Analg*. 2002;94:1161.
39. Vergheze ST, Hannallah RS, Rice LJ, et al. Caudal anesthesia in children: effect of volume versus concentration of bupivacaine on blocking spermatic cord traction response during orchidopexy. *Anesth Analg*. 2002;95:1219.
40. Groban L, Deal D, Vernon J, et al. Cardiac resuscitation after incremental overdosage with lidocaine, bupivacaine, levobupivacaine, and ropivacaine in anesthetized dogs. *Anesth Analg*. 2001;92:37.
41. Ivani G, DeNegri P, Conio A, et al. Comparison of racemic bupivacaine, ropivacaine, and levo-bupivacaine for pediatric caudal anesthesia: effects on postoperative analgesia and motor block. *Reg Anesth Pain Med*. 2002;27:157.
42. Ivani G, De Negri P, Conio A, et al. Ropivacaine-clonidine combination for caudal blockade in children. *Acta Anaesthesiol Scand*. 2000;44:446.
43. Wheeler M, Patel A, Suresh S, et al. The addition of clonidine 2 microg kg⁻¹ does not enhance the postoperative analgesia of a caudal block using 0.125% bupivacaine and epinephrine 1:200,000 in children: a prospective, double-blind, randomized study. *Paediatr Anaesth*. 2005;15:476.
44. Hansen TG, Henneberg SW, Walther-Larsen S, et al. Caudal bupivacaine supplemented with caudal or intravenous clonidine in children undergoing hypospadias repair: a double-blind study. *Br J Anaesth*. 2004;92:223.
45. Kumar P, Rudra A, Pan AK, et al. Caudal additives in pediatrics: a comparison among midazolam, ketamine, and neostigmine coadministered with bupivacaine. *Anesth Analg*. 2005;101:69-73.
46. van Schoor AN, Boon JM, Bosenberg AT, et al. Anatomical considerations of the pediatric ilioinguinal/iliohypogastric nerve block. *Paediatr Anaesth*. 2005;15:371.
47. Willschke H, Marhofer P, Bosenberg A, et al. Ultrasonography for ilioinguinal/iliohypogastric nerve blocks in children. *Br J Anaesth*. 2005;95:226.
48. Choi WY, Irwin MG, Hui TW, et al. EMLA cream versus dorsal penile nerve block for postcircumcision analgesia in children. *Anesth Analg*. 2003;96:396.
49. Antok E, Bordet F, Duflo F, et al. Patient-controlled epidural analgesia versus continuous epidural infusion with ropivacaine for postoperative analgesia in children. *Anesth Analg*. 2003;97:1608.
50. Lejus C. What does analgesia mask? *Paediatr Anaesth*. 2004;14:622.
51. Bosenberg AT, Bland BA, Schulte-Steinberg O, et al. Thoracic epidural anesthesia via caudal route in infants. *Anesthesiology*. 1988;69:265.
52. Valairucha S, Seefelder C, Houck CS. Thoracic epidural catheters placed by the caudal route in infants: the importance of radiographic confirmation. *Paediatr Anaesth*. 2002;12:424.
53. Tsui BC, Wagner A, Cave D, et al. Thoracic and lumbar epidural analgesia via the caudal approach using electrical stimulation guidance in pediatric patients: a review of 289 patients. *Anesthesiology*. 2004;100:683.
54. Benumof JL. Permanent loss of cervical spinal cord function associated with interscalene block performed under general anesthesia. *Anesthesiology*. 2000;93:1541.
55. Schuepfer G, Johr M. Psoas compartment block in children: part I—description of the technique. *Paediatr Anaesth*. 2005;15:461.
56. Schuepfer G, Johr M. Psoas compartment block in children: part II—generation of an institutional learning curve with a new technique. *Paediatr Anaesth*. 2005;15:465.
57. Birmingham PK, Tobin MJ, Fisher DM, et al. Initial and subsequent dosing of rectal acetaminophen in children: a 24-hour pharmacokinetic study of new dose recommendations. *Anesthesiology*. 2001;94:385.
58. Bolton P, Bridge H, Montgomery C, et al. The analgesic efficacy of preoperative high dose (40 mg kg⁻¹) oral acetaminophen after bilateral myringotomy and tube insertion in children. *Paediatr Anaesth*. 2002;12:29.
59. Williams DG, Patel A, Howard RF. Pharmacogenetics of codeine metabolism in an urban population of children and its implications for analgesic reliability. *Br J Anaesth*. 2002;89:839.
60. Galinkin JL, Fazi LM, Cuy RM, et al. Use of intranasal fentanyl in children undergoing myringotomy and tube placement during halothane and sevoflurane anesthesia. *Anesthesiology*. 2000;93:1378.
61. Gan TJ, Meyer T, Apfel CC, et al. Consensus guidelines for managing postoperative nausea and vomiting. *Anesth Analg*. 2003;97:62.
62. Eberhart LH, Morin AM, Guber D, et al. Applicability of risk scores for postoperative nausea and vomiting in adults to paediatric patients. *Br J Anaesth*. 2004;93:386.
63. Eberhart LH, Geldner G, Kranke P, et al. The development and validation of a risk score to predict the probability of postoperative vomiting in pediatric patients. *Anesth Analg*. 2004;99:1630.
64. Sarti A, Busoni P, Dell'Oste C, et al. Incidence of vomiting in susceptible children under regional analgesia with two different anaesthetic techniques. *Paediatr Anaesth*. 2004;14:251.
65. Bortone L, Picetti E, Mergoni M. Anaesthesia with sevoflurane in children: nitrous oxide does not increase postoperative vomiting. *Paediatr Anaesth*. 2002;12:775.

66. Olutoye O, Jantzen EC, Alexis R, et al. A comparison of the costs and efficacy of ondansetron and dolasetron in the prophylaxis of postoperative vomiting in pediatric patients undergoing ambulatory surgery. *Anesth Analg*. 2003;97:390.
67. Kim MS, Coté CJ, Cristoloveanu C, et al. There is no dose-escalation response to dexamethasone (0.0625-1.0 mg/kg) in pediatric tonsillectomy or adenotonsillectomy patients for preventing vomiting, reducing pain, shortening time to first liquid intake, or the incidence of voice change. *Anesth Analg*. 2007;104:1052-1058.
68. Mathew PJ, Madan R, Subramaniam R, et al. Efficacy of low-dose dexamethasone for preventing postoperative nausea and vomiting following strabismus repair in children. *Anaesth Intensive Care*. 2004;32:372.
69. Splinter WM, Rhine EJ. Low-dose ondansetron with dexamethasone more effectively decreases vomiting after strabismus surgery in children than does high-dose ondansetron. *Anesthesiology*. 1998;88:72.
70. Domino KB, Anderson EA, Polissar NL, Posner KL. Comparative efficacy and safety of ondansetron, droperidol, and metoclopramide for preventing postoperative nausea and vomiting: a meta-analysis. *Anesth Analg*. 1999;88:1370-1379.
71. Scuderi PE, Weaver RG, James RL, et al. A randomized, double-blind, placebo controlled comparison of droperidol, ondansetron, and metoclopramide for the prevention of vomiting following outpatient strabismus surgery in children. *J Clin Anesth*. 1997;9:551.
72. Wang SM, Kain ZN. P6 acupoint injections are as effective as droperidol in controlling early postoperative nausea and vomiting in children. *Anesthesiology*. 2002;97:359.
73. Voepel-Lewis T, Malviya S, Tait AR. A prospective cohort study of emergence agitation in the pediatric postanesthesia care unit. *Anesth Analg*. 2003;96:1625.
74. Cravero J, Surgenor S, Whalen K. Emergence agitation in pediatric patients after sevoflurane anaesthesia and no surgery: a comparison with halothane. *Paediatr Anaesth*. 2000;10:419.
75. Cole JW, Murray DJ, McAllister JD, et al. Emergence behaviour in children: defining the incidence of excitement and agitation following anaesthesia. *Paediatr Anaesth*. 2002;12:442.
76. Weldon BC, Bell M, Craddock T. The effect of caudal analgesia on emergence agitation in children after sevoflurane versus halothane anesthesia. *Anesth Analg*. 2004;98:321.
77. Cohen IT, Finkel JC, Hannallah RS, et al. Rapid emergence does not explain agitation following sevoflurane anaesthesia in infants and children: a comparison with propofol. *Paediatr Anaesth*. 2003;13:63.
78. Oh AY, Seo KS, Kim SD, et al. Delayed emergence process does not result in a lower incidence of emergence agitation after sevoflurane anesthesia in children. *Acta Anaesthesiol Scand*. 2005;49:297.
79. Shibata S, Shigeomi S, Sato W, et al. Nitrous oxide administration during washout of sevoflurane improves postanesthetic agitation in children. *J Anesth*. 2005;19:160.
80. Kain ZN, Caldwell-Andrews AA, Weinberg ME, et al. Sevoflurane versus halothane: postoperative maladaptive behavioral changes: a randomized, controlled trial. *Anesthesiology*. 2005;102:720.
81. Keaney A, Diviney D, Harte S, et al. Postoperative behavioral changes following anesthesia with sevoflurane. *Paediatr Anaesth*. 2004;14:866.
82. Kain ZN, Caldwell-Andrews AA, Maranets I, et al. Preoperative anxiety, emergence delirium and postoperative maladaptive behaviors: are they related? *Anesth Analg*. 2004;99:1648.
83. Park JM, Houck CS, Sethna NF, et al. Ketorolac suppresses postoperative bladder spasms after pediatric ureteral reimplantation. *Anesth Analg*. 2000;91:11-15.
84. Halachmi S, El-Ghoneimi A, Bissonnette B, et al. Hemodynamic and respiratory effect of pediatric urological laparoscopic surgery: a retrospective study. *J Urol*. 2003;170:1651.
85. Weld KJ, Ames CD, Landman J, et al. Evaluation of intra-abdominal pressures and gas embolism during laparoscopic partial nephrectomy in a porcine model. *J Urol*. 2005;174:1457.
86. Sakka SG, Huettemann E, Petrat G, et al. Transoesophageal echocardiographic assessment of haemodynamic changes during laparoscopic herniorrhaphy in small children. *Br J Anaesth*. 2000;84:330.
87. Gomez Dammeyer BH, Karanik E, Gluer S, et al. Anuria during pneumoperitoneum in infants and children: a prospective study. *J Pediatr Surg*. 2005;40:1454.
88. Hammer GB. Single-lung ventilation in infants and children. *Paediatr Anaesth*. 2004;14:98.
89. Kalfa N, Allal H, Montes-Tapia F, et al. Ideal timing of thoracoscopic decortication and drainage for empyema in children. *Surg Endosc*. 2004;18:472-477.
90. Hammer GB, Fitzmaurice BG, Brodsky JB. Methods for single-lung ventilation in pediatric patients. *Anesth Analg*. 1999;89:1426.
91. Lawson ML, Mellins RB, Tabangin M, et al. Impact of pectus excavatum on pulmonary function before and after repair with the Nuss procedure. *J Pediatr Surg*. 2005;40:174.
92. Nuss D. Recent experiences with minimally invasive pectus excavatum repair "Nuss procedure." *Jpn J Thorac Cardiovasc Surg*. 2005;53:338.
93. Hammer GB. Anaesthetic management for the child with a mediastinal mass. *Paediatr Anaesth*. 2004;14:95.
94. Sponheim S, Skraastad O, Helseth E, et al. Effects of 0.5 and 1.0 MAC isoflurane, sevoflurane and desflurane on intracranial and cerebral perfusion pressures in children. *Acta Anaesthesiol Scand*. 2003;47:932-938.
95. Hayward R, Gonzalez S. How low can you go? Intracranial pressure, cerebral perfusion pressure, and respiratory obstruction in children with complex craniosynostosis. *J Neurosurg*. 2005;102:16.
96. Ririe DG, David LR, Glazier SS, et al. Surgical advancement influences perioperative care: a comparison of two surgical techniques for sagittal craniosynostosis repair. *Anesth Analg*. 2003;97:699.
97. Williams GD, Ellenbogen RG, Gruss JS. Abnormal coagulation during pediatric craniofacial surgery. *Pediatr Neurosurg*. 2001;35:5.
98. Faberowski LW, Black S, Mickle JP. Incidence of venous air embolism during craniectomy for craniosynostosis repair. *Anesthesiology*. 2000;92:20.