

Chapter 67

REFERENCES

- Bolton P, Bridge HS, Montgomery CJ, et al. The analgesic efficacy of preoperative high dose (40mg/kg) oral acetaminophen after bilateral myringotomy and tube insertion in children. *Pediatr Anesth.* 2002;12:29-35.
- Galinkin JL, Fazi LM, Cuy RM, et al. Use of intranasal fentanyl in children undergoing myringotomy and tube placement during halothane and sevoflurane anesthesia. *Anesthesiology.* 2000;93:1378-1383.
- Voronov P, Tobin MJ, Billings K, et al. Postoperative pain relief in infants undergoing myringotomy and tube placement: a comparison of a novel regional anesthetic block to intranasal fentanyl—a pilot analysis. *Pediatr Anesth.* 2008;18:1196-1201.
- Kawasaki T, Sata T, Kawasaki C, et al. Airway management of a child with temporomandibular joint ankylosis following otitis media. *Anesthesia.* 2002;57:294-295.
- Ho LC, Crosby G, Sundaram P, et al. Ulnar train-of-four stimulation in predicting face movement during intracranial facial nerve stimulation. *Anesth Analg.* 1989;69:242-244.
- Marchal JM, Gomez-Luque A, Martos-Crespo F, et al. Clonidine decreases intraoperative bleeding in middle ear microsurgery. *Acta Anaesth Scand.* 2001;45:627-633.
- Ryu JH, Sohn IS, Do SH. Controlled hypotension for middle ear surgery: a comparison between remifentanil and magnesium sulphate. *Br J Anesth.* 2009;103:490-495.
- Gombar S, Kaur J, Gombar KK, et al. Superior anti-emetic efficacy of granisetron-dexamethasone combination in children undergoing middle ear surgery. *Acta Anesth Scand.* 2007;51:621-624.
- Mukherjee K, Seavell C, Rawlings E, et al. A comparison of total intravenous with balanced anesthesia for middle ear surgery: effects on postoperative nausea and vomiting, pain, and conditions of surgery. *Anesthesia.* 2003;58:176-182.
- Davis I, More JRM, Lahiri SK. Nitrous oxide and the middle ear. *Anesthesia.* 1979;34:147-151.
- Gan TJ, Meyer T, Apfel CC, et al. Consensus guidelines for managing postoperative nausea and vomiting. *Anesth Analg.* 2003;97:62-71.
- Rowlands RG, Harris R, Hern JD, et al. Day-case pediatric mastoid surgery. *Int J Pediatric Otorhinolaryngol.* 2003;67:771-775.
- Uezono S, Holzman RS, Goto T, et al. Prediction of difficult airway in school-aged patients with microtia. *Pediatric Anesth.* 2001;11:409-411.
- Granier M, Dadure C, Bringuier S, et al. Intranasal lidocaine plus naphazoline nitrate improves surgical conditions and perioperative analgesia in septorhinoplasty surgery. *Can J Anesth.* 2009;56:102-108.
- Erbay H, Kara CO, Kara IG, et al. Breathing or ventilation via a mouth mask for rhinoplasty operations in the early postanesthetic period. *Eur J Anaesth.* 2003;20:753-759.
- Gujrathi CS, Daniel SJ, James AL, et al. Management of bilateral choanal atresia in the neonate: an institutional review. *Int J Pediatric Otorhinolaryngol.* 2004;68:399-407.
- Lim EH, Tan S, Lim SL. Anesthesia for a neonate with persistent buccopharyngeal membrane and unilateral choanal atresia. *Pediatr Anesth.* 2005;15:509-511.
- Aqil M, Ulhaq A, Arif A, et al. Venous air embolism during use of a Nd YAG laser. *Anesthesia.* 2008;63:1006-1009.
- Yang JJ, Li WY, Wang ZY, et al. Local anesthesia for functional endoscopic sinus surgery employing small volumes of epinephrine-containing solutions of lidocaine produces profound hypotension. *Acta Anaesth Scand.* 2005;49:1471-1476.
- Durmus M, But AK, Dogan Z, et al. Effect of dexmedetomidine on bleeding during tympanoplasty or septorhinoplasty. *Eur J Anaesth.* 2007;24:447-453.
- Goksu S, Arik H, Demiryurek S, et al. Effects of dexmedetomidine infusion in patients undergoing functional endoscopic sinus surgery under local anaesthesia. *Eur J Anaesth.* 2008;25:22-28.
- Turan A, Memis D, Karamanlioglu B, et al. The analgesic effects of gabapentin in monitored anesthesia care for ear-nose-throat surgery. *Anesth Analg.* 2004;99:375-378.
- Yellon RF. Prevention and management of complications of airway surgery in children. *Pediatr Anesth.* 2004;14:107-111.
- Kain ZN, Caldwell-Andrews AA, Maranets I, et al. Predicting which child-parent pair will benefit from parental presence during induction of anesthesia: a decision making approach. *Anesth Analg.* 2006;102:81-84.
- Arai YCP, Kandatsu N, Ito H, et al. Induction and emergence behavior of children undergoing general anesthesia correlates with maternal salivary amylase activity before anesthesia. *Acta Anaesth Scand.* 2007;52:285-288.
- Homer JR, Elwood T, Peterson D, Rampersad S. Risk factors for adverse events in children with colds emerging from anesthesia: a logistic regression. *Pediatr Anesth.* 2006;17:154-161.
- Tait AR, Malviya S. Anesthesia for the child with an upper respiratory tract infection: still a dilemma? *Anesth Analg.* 2005;100:59-65.

28. Mamie C, Habre W, Delhumeau C, et al. Incidence and risk factors of perioperative respiratory adverse events in children undergoing elective surgery. *Pediatr Anesth.* 2004;14:218-224.
29. Nafiu OO, Green, GE, Walton S, et al. Obesity and risk of perioperative complications in children presenting for adenotonsillectomy. *Int J Pediatr Otorhinolaryngol.* 2009;73:89-95.
30. Granzotto, HG, Aquino FV, Flores JA, Neto JFL. Tonsil size as a predictor of cardiac complications in children with sleep-disordered breathing. *Laryngoscope.* 2010;120:1246-1251.
31. Yellon RF. Is polysomnography required prior to tonsillectomy and adenoidectomy for diagnosis of obstructive sleep apnea versus mild sleep disordered breathing in children? *Laryngoscope.* 2010;120:868-869.
32. Kim JA, Lee JJ. Preoperative predictors of difficult intubation in patients with obstructive sleep apnea syndrome. *Can J Anesth.* 2006;53:393-397.
33. Isono S. Obstructive sleep apnea of obese adults. *Anesthesiology.* 2009;110:908-921.
34. American Society of Anesthesiologists. Practice Guidelines for the Perioperative Management of Patients With Obstructive Sleep Apnea Syndrome. *Anesthesiology.* 2006;104:1081-1093.
35. Reber A, Paganoni R, Frei FJ. Effects of common airway maneuvers on upper airway dimensions and clinical signs in anaesthetized, spontaneously breathing children. *Br J Anaesth.* 2001;86:217-222.
36. Young-Chang PA, Nakayama M, Kato N, et al. The effects of jaw thrust and the lateral position on heart rate variability in anesthetized children with obstructive sleep apnea syndrome. *Anesth Analg.* 2007;104:1352-1355.
37. Khan FA, Memon GA. Comparison of spontaneous with controlled mode of ventilation in tonsillectomy. *Pediatr Anesth.* 2001;11:185-190.
38. Woods AW, Allam S. Tracheal intubation without the use of neuromuscular blocking agents. *Br J Anaesth.* 2005;94:150-158.
39. Gravningsbraten R, Nicklasson B, Raeder J. Safety of laryngeal mask airway and short-stay practice in office-based adenotonsillectomy. *Acta Anaesth Scand.* 2008;53:218-222.
40. Tsui BCH, Wagner A, Cave D, et al. The incidence of laryngospasm with a “no-touch” extubation technique after tonsillectomy and adenoidectomy. *Anesth Analg.* 2004;98:327-329.
41. Isaacson G. Avoiding airway obstruction after pediatric adenotonsillectomy. *Int J Pediatr Otorhinolaryngol.* 2009;73:803-806.
42. Raghavendran S, Bagry H, Dethieux G, et al. Anesthetic management protocol to decrease respiratory complications after adenotonsillectomy in children with severe sleep apnea. *Anesth Analg.* 2010;110:1093-1101.
43. Didem D, Celebi N, Elvan EG, et al. The efficacy of intravenous or peritonsillar ketamine for postoperative pain relief in children following adenotonsillectomy. *Pediatr Anesth.* 2006;17:263-269.
44. Honarmand A, Safavi MR, Jamshidi M. The preventative analgesic effects of preincisional peritonsillar infiltration of two low doses of ketamine for postoperative pain relief in children following adenotonsillectomy. A randomized, double-blind, placebo-controlled study. *Pediatr Anesth.* 2008;18:508-514.
45. Akkaya T, Bedirli N, Ceylan T, et al. Comparison of intravenous and peritonsillar infiltration of tramadol for postoperative pain relief in children following adenotonsillectomy. *Eur J Anaesth.* 2009;26:333-337.
46. Alhashemi JA, Daghastani MF. Effects of intraoperative i.v. acetaminophen vs. i.m. meperidine on post-tonsillectomy pain in children. *Br J Anaesth.* 2006;96:790-795.
47. White MC, Nolan JA. An evaluation of pain and postoperative nausea and vomiting following the introduction of guidelines for tonsillectomy. *Pediatr Anesth.* 2005;15:683-688.
48. Arcioni R, Rocca MD, Romano S, et al. Ondansetron inhibits the analgesic effects of tramadol: a possible 5-HT₃ spinal receptor involvement in acute pain in humans. *Anesth Analg.* 2002;94:1553-1557.
49. Moiniche S, Romsing J, Dahl JB, et al. Nonsteroidal anti-inflammatory drugs and the risk of operative site bleeding after tonsillectomy: a quantitative systematic review. *Anesth Analg.* 2003;96:68-77.
50. Marret E, Flahault A, Samama CM, et al. Effects of postoperative, nonsteroidal, anti-inflammatory drugs on bleeding risk after tonsillectomy. *Anesthesiology.* 2003;98:1497-1502.
51. Cardwell M, Siviter G, Smith A. Non-steroidal anti-inflammatory drugs and perioperative bleeding in pediatric tonsillectomy. *Anesth Analg.* 2005;101:1558.
52. Aouad MT, Siddik SS, Rizak LB, et al. The effect of dexamethasone on postoperative vomiting after tonsillectomy. *Anesth Analg.* 2001;92:636-640.
53. Elhakim M, Ali NM, Rashed I, et al. Dexamethasone reduces postoperative vomiting and pain after pediatric tonsillectomy. *Can J Anaesth.* 2003;50:392-397.
54. Sukhani R, Pappas AL, Lurie J, et al. Ondansetron and dolasetron provide equivalent postoperative vomiting control after ambulatory tonsillectomy in dexamethasone pretreated children. *Anesth Analg.* 2002;95:1230-1235.
55. Karaman M, Ilhan AE, Dereci G, Tek A. Determination of optimum dosage of intraoperative single dose dexamethasone in pediatric tonsillectomy and adenotonsillectomy. *Int J Pediatr Otorhinolaryngol.* 2009;73:1513-1515.
56. Bolton CM, Myles PS, Nolan T, Sterne JA. Prophylaxis of postoperative vomiting in children undergoing tonsillectomy: a systematic review and meta-analysis. *Br J Anaesth.* 2006;97:593-604.
57. Brown KA, Morin I, Hickey C, et al. Urgent adenotonsillectomy. *Anesthesiology.* 2003;99:586-595.
58. Windfuhr JP. Lethal post-tonsillectomy hemorrhage. *Auris Nasus Larynx.* 2003;30:391-396.
59. Windfuhr JP, Schloendorff G, Baburi D, Kremer B. Serious post-tonsillectomy hemorrhage with and without lethal outcome in children and adolescents. *Int J Pediatr Otorhinolaryngol.* 2008;72:1029-1040.
60. Windfuhr JP, Schloendorff G, Baburi D, Kremer B. Life-threatening posttonsillectomy hemorrhage. *Laryngoscope.* 2008;118:1389-1394.
61. Ovassapian A, Tuncbilek M, Weitzel EK, et al. Airway management in adult patients with deep neck infections: a case series and review in the literature. *Anesth Analg.* 2005;100:585-589.
62. Chen L, Zhang X, Li S, et al. The risk factors for hypoxemia in children younger than 5 years old undergoing rigid bronchoscopy for foreign body removal. *Anesth Analg.* 2009;109:1079-1084.
63. Schmitt HJ, Mang H, Schmitt J. Fiberoptic intubation in patients after radiotherapy for carcinoma of the head and neck: difficulty and predictability. *Eur J Anaesth.* 2004;21:914-927.
64. Buczkowski PW, Fombon FN, Lin ES, et al. Air entrainment during high-frequency jet ventilation in a model of upper tracheal stenosis. *Br J Anaesth.* 2007;99:891-897.

65. Davies JM, Hillel AD, Maronian NC, Posner KL. The Hunsaker Mon-Jet tube with jet ventilation is effective for microlaryngeal surgery. *Can J Anesth.* 2009;56:284-290.
66. Cook TM, Alexander R. Major complications during anaesthesia for elective laryngeal surgery in the U.K.: a national survey of the use of high-pressure source ventilation. *Br J Anaesth.* 2008;101:266-272.
67. Unzueta MC, Casas I, Merten A, et al. Endobronchial high frequency jet ventilation for endobronchial laser surgery: an alternative approach. *Anesth Analg.* 2003;96:298-300.
68. Rezaie-Majd A, Bigenzahn W, Denk DM, et al. Superimposed high-frequency jet ventilation (SHFJV) for endoscopic laryngotracheal surgery in more than 1,500 patients. *Br J Anaesth.* 2006;96:650-659.
69. Hautmann H, Gamarra F, Henke M, et al. High frequency jet ventilation interventional fiberoptic bronchoscopy. *Anesth Analg.* 2000;90:1436-1440.
70. Sawchuk WS, Weber PJ, Lowry DR, et al. Infectious papillomavirus in the vapor of warts treated with CO₂ laser or electrocoagulation: detection and protection. *J Am Acad Dermatol.* 1989;21:41-49.
71. American National Standards Institute. *Safe Use of Lasers in Health Care Facilities.* ANSI Z136.3 2005. Washington, DC: ANSI. 2005.
72. Farrell PT. Prenatal diagnosis and intrapartum management of neck masses causing airway obstruction. *Pediatr Anesth.* 2004;14:48-52.
73. Nargozian C. The airway in patients with craniofacial abnormalities. *Pediatr Anesth.* 2004;14:48-52.
74. Hammer GB, Holzki J, Morton NS, eds. The pediatric airway. *Pediatr Anesth.* 2009;19(Suppl):1-197.
75. Mesnil M, Dadure C, Captier G, et al. A new approach for peri-operative analgesia of cleft palate repair in infants: the bilateral suprazygomatic maxillary nerve block. *Pediatr Anesth.* 2010;20:343-349.
76. Austin J, Ali T. Tracheomalacia and bronchomalacia in children: pathophysiology, assessment, treatment, and anesthesia management. *Pediatr Anesth.* 2003;13:3-11.
77. Walner DL. Advances in laryngotracheal reconstruction in children. *Curr Opin Otolaryngol Head Neck Surg.* 1999;7:308-312.
78. Triglia JM, Nicollas R, Roman S. Primary cricotracheal resection in children: indications, technique and outcome. *Int J Pediatr Otorhinolaryngol.* 2001;58:17-25.
79. Wrightson F, Soma M, Smith JH. Anesthetic experience of 100 pediatric tracheostomies. *Pediatr Anesth.* 2009;19:659-666.
80. Diaz-Reganón G, Minambres E, Ruiz A, et al. Safety and complications of percutaneous tracheostomy in a cohort of 800 mixed ICU patients. *Anesthesia.* 2008;63:1198-1203.
81. Kannan S, Chesnut N, McBride G. Intubating LMA guided awake fiberoptic intubation in severe maxillofacial injury. *Can J Anaesth.* 2000;47:989-991.
82. Anwer HMF, Zeitoun IM, Shehata EAA. Submandibular approach for tracheal intubation in patients with panfacial fractures. *Br J Anaesth.* 2007;98:835-840.
83. Biswas BK, Joshi S, Bhattacharyya P, et al. Percutaneous dilational tracheostomy kit: an aid to submental intubation. *Anesth Analg.* 2006;103:1055.
84. Mallory S, Yap LH, Jones BM, et al. Anaesthetic management in facial bipartition surgery. The experience of one centre. *Anaesthesia.* 2004;59:44-51.
85. Stricker PA, Shaw TL, Desouza DG, et al. Blood loss, replacement, and associated morbidity in infants and children undergoing craniofacial surgery. *Pediatr Anesth.* 2010;20:150-159.
86. Hall CEJ, Shutt LE. Nasotracheal intubation for head and neck surgery. *Anaesthesia.* 2003;58:249-256.
87. Ahmed-Nusrath A, Tong JL, Smith JE. Pathways through the nose for nasal intubation: a comparison of three endotracheal tubes. *Br J Anaesth.* 2008;100:269-274.
88. Davies JR, Reid AP. Preventing damage to the tracheal tube during maxillary osteotomy. *Anaesthesia.* 2003;58:914-915.
89. Moller PL, Sindet-Pederson S, Petersen CT, et al. Onset of acetaminophen analgesia: comparison of oral and intravenous routes after third molar surgery. *Br J Anaesth.* 2005;94:642-648.
90. Van Aken H, Thys L, Veekman L, et al. Assessing analgesia in single and repeated administrations of propacetamol for postoperative pain: comparison with morphine after dental surgery. *Anesth Analg.* 2004;98:159-165.
91. Ansari AA, personal communication, 2010.
92. Rutherford J, Stevenson R. Careful physical examination is essential in the preoperative assessment of children for dental extractions under general anesthesia. *Pediatr Anesth.* 2004;14:920-923.
93. Rama-Maceiras P, Ferreira TA, Molins N, et al. Less postoperative nausea and vomiting after propofol + remifentanil versus propofol + fentanyl anesthesia during plastic surgery. *Acta Anaesth Scand.* 2005;49:305.
94. Chaparro LE, Gallo T, Gonzalez NJ, et al. Effectiveness of combined haloperidol and dexamethasone versus dexamethasone only for postoperative nausea and vomiting in high-risk day surgery patients: a randomized blinded trial. *Eur J Anaesth.* 2010;27:187-191.
95. Farling PA. Thyroid disease. *Br J Anaesth.* 2000;85:15-28.
96. Yumoto E, Minoda R, Hyodo M, et al. Causes of recurrent laryngeal nerve paralysis. *Auris Nasus Larynx.* 2002;29:41-45.
97. Randolph GW, Dralle H, et al. Electrophysiologic Recurrent laryngeal Nerve Monitoring During Thyroid and Parathyroid Surgery: International Standards Guideline Statement. *Laryngoscope.* 2011;121:S1-S16.
98. Mihai R, Farndon JR. Parathyroid disease and calcium metabolism. *Br J Anaesth.* 2000;85:29-43.
99. Mason RA, Fielder CP. The obstructed airway in head and neck surgery. *Anaesthesia.* 1999;54:625-628.
100. Scholz A, Pugh S, Fardy M, et al. The effect of dobutamine on blood flow of free tissue transfer flaps during head and neck reconstructive surgery. *Anaesthesia.* 2009;64:1089-1093.
101. Leiendoeker J, Hocker J, Meybohm P, et al. Postoperative neurocognitive function and microembolus detection in patients undergoing neck dissection: a pilot study. *Eur J Anaesth.* 2010;27:417-424.
102. Strong MS, Vaughn CW, Mahler DL, et al. Cardiac complications of microsurgery of the larynx: etiology, incidence, and prevention. *Laryngoscope.* 1974;84:908-920.