

## Chapter 69

### REFERENCES

1. Alspach D, Falleroni M. Monitoring patients during procedures conducted outside the operating room. *Int Anesthesiol Clin.* 2004;42:95-111.
2. American Society of Anesthesiologists. *Continuum of Depth of Sedation: Definition of General Anesthesia and Levels of Sedation/Analgesia.* Approved by the ASA House of Delegates on October 27, 2004, and amended on October 21, 2009.
3. American Society of Anesthesiologists. *Position on Monitored Anesthesia Care.* Approved by the ASA House of Delegates on October 21, 1986, amended on October 25, 2005 and last updated on September 2, 2008.
4. American Society of Anesthesiologists. *Statement on Granting Privileges to Nonanesthesiologist Practitioners for Personally Administering Deep Sedation or Supervising Deep Sedation by Individuals Who Are Not Anesthesia Professionals.* Approved by the ASA House of Delegates on October 18, 2006. Available at: <http://www.asahq.org/publicationsAndServices/standards/39.pdf>. Accessed July 20, 2010.
5. American Society of Anesthesiologists. *Statement on Granting Privileges for Administration of Moderate Sedation to Practitioners Who Are Not Anesthesia Professionals.* Origin: Ad Hoc Committee on Credentialing. Approved by the ASA House of delegates on October 25, 2005 and amended on October 18, 2006. Available at: <http://www.asahq.org/publicationsAndServices/standards/40.pdf>. Accessed July 20, 2010.
6. Kearse LA Jr, Manberg P, Chamoun N, deBros F, Zaslavsky A. Bispectral analysis of the electroencephalogram correlates with patient movement to skin incision during propofol/nitrous oxide anesthesia. *Anesthesiology.* 1994;81:1365-1370.
7. Pomfrett CJ. Heart rate variability, BIS and depth of anaesthesia'. *Br J Anaesth.* 1999;82:659-662.
8. Glass PS, Bloom M, Kearse L, Rosow C, Sebel P, Manberg P. Bispectral analysis measures sedation and memory effects of propofol, midazolam, isoflurane, and alfentanil in healthy volunteers. *Anesthesiology.* 1997;86:836-847.
9. Kearse LA Jr, Rosow C, Zaslavsky A, Connors P, Dershwitz M, Denman W. Bispectral analysis of the electroencephalogram predicts conscious processing of information during propofol sedation and hypnosis. *Anesthesiology.* 1998;88:25-34.
10. Liu J, Singh H, White PF. Electroencephalographic bispectral index correlates with intraoperative recall and depth of propofol-induced sedation. *Anesth Analg.* 1997;84:185-189.
11. Ibrahim AE, Taraday JK, Kharasch ED. Bispectral index monitoring during sedation with sevoflurane, midazolam, and propofol. *Anesthesiology.* 2001;95:1151-1159.
12. Liu J, Singh H, White PF. Electroencephalogram bispectral analysis predicts the depth of midazolam-induced sedation. *Anesthesiology.* 1996;84:64-69.
13. Leslie K, Sessler DI, Schroeder M, Walters K. Propofol blood concentration and the Bispectral Index predict suppression of learning during propofol/epidural anesthesia in volunteers. *Anesth Analg.* 1995;81:1269-1274.
14. Gan TJ, Glass PS, Windsor A, Payne F, Rosow C, Sebel P, Manberg P. Bispectral index monitoring allows faster emergence and improved recovery from propofol, alfentanil, and nitrous oxide anesthesia. BIS Utility Study Group. *Anesthesiology.* 1997;87:808-815.
15. Bower AL, Ripepi A, Dilger J, Boparai N, Brody FJ, Ponsky JL. Bispectral index monitoring of sedation during endoscopy. *Gastrointest Endosc.* 2000;52:192-196.
16. Gill M, Green SM, Krauss B. A study of the Bispectral Index Monitor during procedural sedation and analgesia in the emergency department. *Ann Emerg Med.* 2003;41:234-241.
17. Liu J, Singh H, White PF. Electroencephalogram bispectral analysis predicts the depth of midazolam-induced sedation. *Anesthesiology.* 1996;84:64-69.
18. Iselin-Chaves IA, Flaishon R, Sebel PS, et al. The effect of the interaction of propofol and alfentanil on recall, loss of consciousness, and the Bispectral Index. *Anesth Analg.* 1998;87:949-955.
19. Mi WD, Sakai T, Kudo T, Kudo M, Matsuki A. Performance of bispectral index and auditory evoked potential monitors in detecting loss of consciousness during anaesthetic induction with propofol with and without fentanyl. *Eur J Anaesthesiol.* 2004;21:807-811.
20. Dahaba AA. Different conditions that could result in the bispectral index indicating an incorrect hypnotic state. *Anesth Analg.* 2005;101:765-773.
21. Suzuki M, Edmonds HL Jr, Tsueda K, Malkani AL, Roberts CS: Effect of ketamine on bispectral index and levels of sedation. *J Clin Monit Comput.* 1998;14:373.
22. Morley AP, Chung DC, Wong AS, Short TG. The sedative and electroencephalographic effects of regional anaesthesia. *Anaesthesia.* 2000;55:864-869.
23. Pollock JE, Neal JM, Liu SS, et al. Sedation during spinal anesthesia. *Anesthesiology.* 2000;93:728-734.
24. American Society of Anesthesiologists. *Statement on Nonoperating Room Anesthetizing Locations.* Approved by the ASA House of Delegates on October 15, 2003 and amended on October 22, 2008.

25. Carlsson U, Grattidge P. Sedation for upper gastrointestinal endoscopy: a comparative study of propofol and midazolam. *Endoscopy*. 1995;27:240-243.
26. K K, N K, E A. The sedative features of propofol and midazolam for upper gastrointestinal endoscopy: a comparative study. *Hellenic J Gastroenterol*. 1997;10:64-67.
27. McQuaid KR, Laine L. A systematic review and meta-analysis of randomized, controlled trials of moderate sedation for routine endoscopic procedures. *Gastrointest Endosc*. 2008;67:910-923.
28. Fudickar A, Bein B, Tonner PH. Propofol infusion syndrome in anaesthesia and intensive care medicine. *Curr Opin Anaesthesiol*. 2006;19:404-410.
29. Jung JS, Park JS, Kim SO, et al. Prophylactic antiemetic effect of midazolam after middle ear surgery. *Otolaryngol Head Neck Surg*. 2007;137:753-756.
30. Lee Y, Wang JJ, Yang YL, Chen A, Lai HY. Midazolam vs ondansetron for preventing postoperative nausea and vomiting: a randomised controlled trial. *Anaesthesia*. 2007;62:18-22.
31. Riad W, Altaf R, Abdulla A, Oudan H. Effect of midazolam, dexamethasone and their combination on the prevention of nausea and vomiting following strabismus repair in children. *Eur J Anaesthesiol*. 2007;24:697-701.
32. Fields HL, Heinricher MM, Mason P. Neurotransmitters in nociceptive modulatory circuits. *Annu Rev Neurosci*. 1991;14:219-245.
33. Trafton JA, Abbadie C, Marchand S, Mantyh PW, Basbaum AI. Spinal opioid analgesia: how critical is the regulation of substance P signaling? *J Neurosci*. 1999;19:9642-9653.
34. Stein C. The control of pain in peripheral tissue by opioids. *N Engl J Med*. 1995;332:1685-1690.
35. Gupta A, Bodin L, Holmstrom B, Berggren L. A systematic review of the peripheral analgesic effects of intraarticular morphine. *Anesth Analg*. 2001;93:761-770.
36. Goulson DT, Fragneto RY. Anesthesia for gastrointestinal endoscopic procedures. *Anesthesiol Clin*. 2009;27:71-85.
37. Keeffe EB, O'Connor KW. 1989 A/S/G/E survey of endoscopic sedation and monitoring practices. *Gastrointest Endosc*. 1990;36(Suppl):S13-S18.
38. Chudnofsky CR, Wright SW, Dronen SC, et al. The safety of fentanyl use in the emergency department. *Ann Emerg Med*. 1989;18:635-639.
39. Thomas S, Benevelli W, Brown D, et al. Safety of fentanyl for analgesia in adults undergoing air medical transport from trauma scenes. *Air Med J*. 1996;15:57-59.
40. Thomas SH. Fentanyl in the prehospital setting. *Am J Emerg Med*. 2007;25:842-843.
41. Glass PS, Gan TJ, Howell S, Ginsberg B. Drug interactions: volatile anesthetics and opioids. *J Clin Anesth*. 1997;9(Suppl):18S-22S.
42. Phillips WJ, Halpin J, Jones J, McKenzie K. Remifentanyl for procedural sedation in the emergency department. *Ann Emerg Med*. 2009;53:163.
43. Dunn MJ, Mitchell R, Souza CD, Drummond G. Evaluation of propofol and remifentanyl for intravenous sedation for reducing shoulder dislocations in the emergency department. *Emerg Med J*. 2006;23:57-58.
44. Litman RS. Conscious sedation with remifentanyl during painful medical procedures. *J Pain Symptom Manage*. 2000;19:468-471.
45. Ahmad S. Office based—is my anesthetic care any different? Assessment and management. *Anesthesiol Clin*. 2010;28:369-384.
46. Bailey PL, Streisand JB, East KA, et al. Differences in magnitude and duration of opioid-induced respiratory depression and analgesia with fentanyl and sufentanil. *Anesth Analg*. 1990;70:8-15.
47. Monk JP, Beresford R, Ward A. Sufentanil. A review of its pharmacological properties and therapeutic use. *Drugs*. 1988;36:286-313.
48. J D. Clinical experience with repeated ketamine administration for procedures requiring anesthesia. 1969.
49. Strayer RJ, Nelson LS. Adverse events associated with ketamine for procedural sedation in adults. *Am J Emerg Med*. 2008;26:985-1028.
50. Willman EV, Andolfatto G. A prospective evaluation of “ketofol” (ketamine/propofol combination) for procedural sedation and analgesia in the emergency department. *Ann Emerg Med*. 2007;49:23-30.
51. Lilburn JK, Dundee JW, Nair SG, Fee JP, Johnston HM. Ketamine sequelae. Evaluation of the ability of various premedicants to attenuate its psychic actions. *Anaesthesia*. 1978;33:307-311.
52. Lilburn JK, Dundee JW, Nair SG. Attenuation of psychic sequelae from ketamine [proceedings]. *Br J Clin Pharmacol*. 1977;4:641P-642P.
53. Belleville JP, Ward DS, Bloor BC, Maze M. Effects of intravenous dexmedetomidine in humans. I. Sedation, ventilation, and metabolic rate. *Anesthesiology*. 1992;77:1125-1133.
54. Ramsay MA, Luteran DL. Dexmedetomidine as a total intravenous anesthetic agent. *Anesthesiology*. 2004;101:787-790.
55. Abdelmalak B, Makary L, Hoban J, Doyle DJ. Dexmedetomidine as sole sedative for awake intubation in management of the critical airway. *J Clin Anesth*. 2007;19:370-373.
56. Hall JE, Uhrich TD, Barney JA, Arain SR, Ebert TJ. Sedative, amnestic, and analgesic properties of small-dose dexmedetomidine infusions. *Anesth Analg*. 2000;90:699-705.
57. Ebert TJ, Hall JE, Barney JA, Uhrich TD, Colincio MD. The effects of increasing plasma concentrations of dexmedetomidine in humans. *Anesthesiology*. 2000;93:382-394.
58. Panzer O, Moitra V, Sladen RN. Pharmacology of sedative-analgesic agents: dexmedetomidine, remifentanyl, ketamine, volatile anesthetics, and the role of peripheral mu antagonists. *Crit Care Clin*. 2009;25:451-469, vii.
59. Scher CS, Gitlin MC. Dexmedetomidine and low-dose ketamine provide adequate sedation for awake fiberoptic intubation. *Can J Anaesth*. 2003;50:607-610.
60. Bloor BC, Ward DS, Belleville JP, Maze M. Effects of intravenous dexmedetomidine in humans. II. Hemodynamic changes. *Anesthesiology*. 1992;77:1134-1142.
61. Talke P, Chen R, Thomas B, et al. The hemodynamic and adrenergic effects of perioperative dexmedetomidine infusion after vascular surgery. *Anesth Analg*. 2000;90:834-839.
62. Muller S, Borowics SM, Fortis EA, et al. Clinical efficacy of dexmedetomidine alone is less than propofol for conscious sedation during ERCP. *Gastrointest Endosc*. 2008;67:651-659.
63. Arain SR, Ebert TJ. The efficacy, side effects, and recovery characteristics of dexmedetomidine versus propofol when used for intraoperative sedation. *Anesth Analg*. 2002;95:461-466, table of contents.
64. Lusedra [package insert]. Research Triangle Park, NC: Eisai Corp.; 2008.
65. Glass PS. Intravenous infusion techniques: how to do it and why we should do it. *Can J Anaesth*. 1998;45:R117-R1137.

66. Newson C, Joshi GP, Victory R, White PF. Comparison of propofol administration techniques for sedation during monitored anesthesia care. *Anesth Analg*. 1995;81:486-491.
67. Schuttler J, Schwilden H, Stoekel H. Pharmacokinetics as applied to total intravenous anaesthesia. Practical implications. *Anaesthesia*. 1983;38(Suppl):53-56.
68. Egan TD. Target-controlled drug delivery: progress toward an intravenous "vaporizer" and automated anesthetic administration. *Anesthesiology*. 2003;99:1214-1219.
69. van den Nieuwenhuyzen MC, Engbers FH, Vuyk J, Burm AG. Target-controlled infusion systems: role in anaesthesia and analgesia. *Clin Pharmacokinet*. 2000;38:181-190.
70. American Society of Anesthesiologists Task Force on Postanesthetic Care. Practice guidelines for postanesthetic care: a report by the American Society of Anesthesiologists Task Force on Postanesthetic Care. *Anesthesiology*. 2002;96:742-752.
71. Bhananker SM, Posner KL, Cheney FW, et al. Injury and liability associated with monitored anesthesia care: a closed claims analysis. *Anesthesiology*. 2006;104:228-234.
72. Nuttall GA, Bowersox MR, Douglass SB, et al. Morbidity and mortality in the use of electroconvulsive therapy. *J ECT*. 2004;20:237-241.
73. Ding Z, White PF. Anesthesia for electroconvulsive therapy. *Anesth Analg*. 2002;94:1351-1364.
74. Gazdag G, Kocsis N, Tolna J, Ivanyi Z. Etomidate versus propofol for electroconvulsive therapy in patients with schizophrenia. *J ECT*. 2004;20:225-229.
75. Locala JA, Irefin SA, Malone D, Cywinski JB, Samuel SW, Naugle R. The comparative hemodynamic effects of methohexital and remifentanyl in electroconvulsive therapy. *J ECT*. 2005;21:12-15.
76. Andersen FA, Arslan D, Holst-Larsen H. Effects of combined methohexital-remifentanyl anaesthesia in electroconvulsive therapy. *Acta Anaesthesiol Scand*. 2001;45:830-833.
77. Zhang Y, White PF, Thornton L, Perdue L, Downing M. The use of nicardipine for electroconvulsive therapy: a dose-ranging study. *Anesth Analg*. 2005;100:378-381.
78. Tang WK, Ungvari GS. Asystole during electroconvulsive therapy: a case report. *Aust N Z J Psychiatry*. 2001;35:382-385.
79. Freese KJ. Can patients safely undergo electroconvulsive therapy while receiving monoamine oxidase inhibitors? *Convuls Ther*. 1985;1:190-194.
80. Dolenc TJ, Habl SS, Barnes RD, Rasmussen KG. Electroconvulsive therapy in patients taking monoamine oxidase inhibitors. *J ECT*. 2004;20:258-261.
81. Chokhavatia S, Nguyen L, Williams R, Kao J, Heavner JE. Sedation and analgesia for gastrointestinal endoscopy. *Am J Gastroenterol*. 1993;88:393-396.
82. ASGE Guidelines. Complications of upper GI endoscopy. *Curr Opin Anaesthesiol*. 2000;13:55-784.
83. Hutchinson RC, Kenny GN. Sedation for endoscopy. *Curr Opin Anaesthesiol*. 2000;13:415-419.
84. Cohen LB, Hightower CD, Wood DA, Miller KM, Aisenberg J. Moderate level sedation during endoscopy: a prospective study using low-dose propofol, meperidine/fentanyl, and midazolam. *Gastrointest Endosc*. 2004;59:795-803.
85. Kazama T, Takeuchi K, Ikeda K, et al. Optimal propofol plasma concentration during upper gastrointestinal endoscopy in young, middle-aged, and elderly patients. *Anesthesiology*. 2000;93:662-669.
86. Eckardt J, Petersen HO, Hakami-Kermani A, Olsen KE, Jorgensen OD, Licht PB. Endobronchial ultrasound-guided transbronchial needle aspiration of undiagnosed intrathoracic lesions. *Interact Cardiovasc Thorac Surg*. 2009;9:232-235.
87. Cameron SE, Andrade RS, Pambuccian SE. Endobronchial ultrasound-guided transbronchial needle aspiration cytology: a state of the art review. *Cytopathology*. 2010;21:6-26.
88. Eberhardt R, Becker HD, Herth FJ. [Endobronchial ultrasound for diagnosis of the mediastinum]. *Chirurg*. 2008;79(1):50-55.
89. Yasufuku K, Nakajima T, Fujiwara T, et al. Role of endobronchial ultrasound-guided transbronchial needle aspiration in the management of lung cancer. *Gen Thorac Cardiovasc Surg*. 2008;56:268-276.
90. Cox G, Miller JD, McWilliams A, Fitzgerald JM, Lam S. Bronchial thermoplasty for asthma. *Am J Respir Crit Care Med*. 2006;173:965-969.
91. Silvestri GA, Vincent BD, Wahidi MM. A phase 3, randomized, double-blind study to assess the efficacy and safety of fospropofol disodium injection for moderate sedation in patients undergoing flexible bronchoscopy. *Chest*. 2009;135(1):41-47.
92. Abdelmalak B, Gutenberg L, Lorenz RR, Smith M, Farag E, Doyle DJ. Dexmedetomidine supplemented with local anesthesia for awake laryngoplasty. *J Clin Anesth*. 2009;21:442-443.
93. Doyle DJ, Abdelmalak B, Machuzak M, Gildea TR. Anesthesia and airway management for removing pulmonary self-expanding metallic stents. *J Clin Anesth*. 2009;21:529-532.
94. Weber CR, Griffin JM. Evaluation of dexamethasone for reducing postoperative edema and inflammatory response after orthognathic surgery. *J Oral Maxillofac Surg*. 1994;52:35-39.
95. Montgomery MT, Hogg JP, Roberts DL, Redding SW. The use of glucocorticosteroids to lessen the inflammatory sequelae following third molar surgery. *J Oral Maxillofac Surg*. 1990;48(2):179-187.
96. Gersema L, Baker K. Use of corticosteroids in oral surgery. *J Oral Maxillofac Surg*. 1992;50:270-277.
97. Hughes R, McGuire G, Montanera W, Wong D, Carmichael FJ. Upper airway edema after carotid endarterectomy: the effect of steroid administration. *Anesth Analg*. 1997;84:475-478.
98. Abdelmalak B, Ryckman JV, AlHaddad S, Sprung J. Respiratory arrest after successful neodymium:yttrium-aluminum-garnet laser treatment of subglottic tracheal stenosis. *Anesth Analg*. 2002;95:485-486, table of contents.
99. Hung WT, Liao SM, Su JM. Laryngeal mask airway in patients with tracheal stents who are undergoing non-airway related interventions: report of three cases. *J Clin Anesth*. 2004;16:214-216.
100. Conlay LA. Special concerns in the cardiac catheterization lab. *Int Anesthesiol Clin*. 2003;41:63-67.
101. Jaffe RA, Samuels SI. Implantation of cardioverter-defibrillator (ICD). In Jaffe RA, Samuels SI, eds. *Anesthesiologist's Manual of Surgical Procedures*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2004:1165.
102. Jaffe RA, Samuels SI. Direct current (DC) cardioversion. In Jaffe RA, Samuels SI, eds. *Anesthesiologist's Manual of Surgical Procedures*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2004:1163.
103. Watkinson AF, Francis IS, Torrie P, et al. Commentary: the role of anaesthesia in interventional radiology. *Br J Radiol*. 2002;75:105-106.
104. Jaffe RA, Samuels SI. Radiofrequency ablation. In Jaffe RA, Samuels SI, eds. *Anesthesiologist's Manual of Surgical*

- Procedures*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2004:1185.
105. Goldberg SN, Grassi CJ, Cardella JF, et al. Image-guided tumor ablation: standardization of terminology and reporting criteria. *J Vasc Interv Radiol*. 2005;16:765-778.
  106. Poggi G, Riccardi A, Quaretti P, et al. Complications of percutaneous radiofrequency thermal ablation of primary and secondary lesions of the liver. *Anticancer Res*. 2007;27:2911-2916.
  107. Manninen PH, Gignac EM, Gelb AW, Lownie SP. Anesthesia for interventional neuroradiology. *J Clin Anesth*. 1995;7:448-452.
  108. Osborn IP. Anesthetic considerations for interventional neuro-radiology. *Int Anesthesiol Clin*. 2003;41:69-77.
  109. The Arteriovenous Malformation Study Group. Arteriovenous malformations of the brain in adults. *N Engl J Med*. 1999;340:1812-1818.
  110. Martin NA, Khanna R, Doberstein C, Bentson J. Therapeutic embolization of arteriovenous malformations: the case for and against. *Clin Neurosurg*. 2000;46:295-318.
  111. Chi OZ, Liu X, Weiss HR. The effects of propranolol on heterogeneity of rat cerebral small vein oxygen saturation. *Anesth Analg*. 1999;89:690-695.
  112. Kraye S. Anaesthesia for interventional neuroradiology. *Curr Opin Anaesthesiol*. 2000;13:421-427.
  113. Van de Velde M. Interventional neuroradiology. *Curr Opin Anaesthesiol*. 2003;16:417-420.
  114. Yadav JS, Wholey MH, Kuntz RE, et al. Protected carotid-artery stenting versus endarterectomy in high-risk patients. *N Engl J Med*. 2004;351:1493-1501.
  115. Ellis JE, Pai SS. Provascular stents in the radiology suite-an anesthesiologist is needed. *J Cardiothorac Vasc Anesth*. 2005;19:801-804.
  116. Ong JP, Sands M, Younossi ZM. Transjugular intrahepatic portosystemic shunts (TIPS): a decade later. *J Clin Gastroenterol*. 2000;30:14-28.
  117. Kelhoffer ER, Osborn IP. The gastroenterology suite and TIPS. *Int Anesthesiol Clin*. 2003;41:51-61.
  118. Shellock F. *Reference manual for magnetic resonance safety: 2003 Edition*. Salt Lake City, UT: Amirsys Inc., 2003.
  119. Dempsey MF, Condon B. Thermal injuries associated with MRI. *Clin Radiol*. 2001;56:457-465.
  120. American Society of Anesthesiologists. Practice advisory on anesthetic care for magnetic resonance imaging: a report by the American Society of Anesthesiologists Task Force on Anesthetic Care for Magnetic Resonance Imaging. *Anesthesiology*. 2009;110:459-479.
  121. Osborn IP. Magnetic resonance imaging anesthesia: new challenges and techniques. *Curr Opin Anaesthesiol*. 2002;15:443-448.
  122. Gooden CK. Anesthesia for magnetic resonance imaging. *Curr Opin Anaesthesiol*. 2004;17:339-342.
  123. Aisenberg J, Brill JV, Ladabaum U, Cohen LB. Sedation for gastrointestinal endoscopy: new practices, new economics. *Am J Gastroenterol*. 2005;100:996-1000.