

Chapter 82

REFERENCES

1. Emergency Cardiac Care Committee, Subcommittees and Task Forces, American Heart Association: 2010 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2010;122:S640-S946.
2. Nadkarni VM, Larkin GL, Peberdy, MA, et al. First documented rhythm and clinical outcome from in-hospital cardiac arrest among children and adults. *JAMA*. 2006;295:50.
3. Olsson GI, Hallen B. Cardiac arrest during anaesthesia. A computer-aided study of 250,543 anaesthetics. *Acta Anaesthesiol Scand*. 1988;32:653.
4. Weaver WD, Hill D, Fahrenbruch CE, et al. Use of the automatic external defibrillator in the management of out-of-hospital cardiac arrest. *N Engl J Med*. 1988;319:661.
5. Eisenberg MS, Bergner L, Hallstrom A. Cardiac resuscitation in the community: importance of rapid provision and implications for program planning. *JAMA*. 1979;241:1905.
6. Brooks DK. *Resuscitation: Care of the Critically Ill*. London: Edward Arnold; 1986.
7. Wood Library Museum. *Resuscitation: An Historical Perspective*. Park Ridge, IL: Wood Library Museum; 1976.
8. Elam JO, Brown ES, Elder JD Jr. Artificial respiration by mouth to mask method: a study of the respiratory gas exchange of paralyzed patients ventilated by operator's expired air. *N Engl J Med*. 1954;250:749.
9. Gordon AS, Frye CS, Gittelson L, et al. Mouth-to-mouth versus manual artificial respiration for children and adults. *JAMA*. 1958;167:320.
10. Safar P, Escarraga LA, Elam JO. A comparison of the mouth-to-mouth and mouth-to-airway methods of artificial respiration with the chest pressure arm-lift methods. *N Engl J Med*. 1958;258:671.
11. Safar P. Failure of manual respiration. *J Appl Physiol*. 1959;14:84.
12. Hooker DR, Kouwenhoven WB, Langworthy OR. The effects of alternating current on the heart. *Am J Physiol*. 1933;103:444.
13. Beck CS, Pritchard WH, Feil HS. Ventricular fibrillation of long duration abolished by electric shock. *JAMA*. 1947;135:985.
14. Zoll PM, Linenthal AJ, Gibson W, et al. Termination of ventricular fibrillation in man by an externally applied electric shock. *N Engl J Med*. 1956;254:727.
15. Kouwenhoven WB, Milnor WR, Knickerbocker GG, et al. Closed chest defibrillation of the heart. *Surgery*. 1957;42:550.
16. Kouwenhoven WB, Jude JR, Knickerbocker GG. Closed-chest cardiac massage. *JAMA*. 1960;173:1064.
17. Redding JS, Pearson JW. Evaluation of drugs for cardiac resuscitation. *Anesthesiology*. 1963;24:203.
18. Weisfeldt ML, Becker LB. Resuscitation after cardiac arrest: a 3-phase time-sensitive model. *JAMA*. 2002;288:3035.
19. Bossaert L, Van Hoeyweghen R, The Cerebral Resuscitation Study Group. Bystander cardiopulmonary resuscitation (CPR) in out-of-hospital cardiac arrest. *Resuscitation*. 1989;17(Suppl):S55.
20. Van Hoeyweghen RJ, Bossaert LL, Mullie A, et al. Quality and efficiency of bystander CPR. *Resuscitation*. 1993;26:47.
21. Berg RA, Kern KB, Sanders AB, et al. Bystander cardiopulmonary resuscitation: is ventilation necessary? *Circulation*. 1993;88:1907.
22. Berg RA, Wilcoxson D, Hilwig RW, et al. The need for ventilatory support during bystander cardiopulmonary resuscitation. *Ann Emerg Med*. 1995;24:342.
23. Berg RA, Kern KB, Hilwig RW, et al. Assisted ventilation does not improve outcome in a porcine model of single-rescuer bystander CPR. *Circulation*. 1997;95:1635.
24. Assar D, Chamberlain D, Colquhoun M, et al. Randomized controlled trials of staged teaching for basic life support: 1. skill acquisition at bronze stage. *Resuscitation*. 2000;45:7.
25. Kern KB, Hilwig R, Berg RA, et al. Importance of continuous chest compressions during cardiopulmonary resuscitation: improved outcome during a simulated single lay-rescuer scenario. *Circulation*. 2002;105:645.
26. Ewy GA, Zuercher M, Hilwig RW, et al. Improved neurological outcome with continuous chest compressions compared with 30:2 compressions-to-ventilations cardiopulmonary resuscitation in a realistic swine model of out-of-hospital cardiac arrest. *Circulation*. 2007;116:2525.
27. Hallstrom A, Cobb L, Johnson E, Copass M. Cardiopulmonary resuscitation by chest compressions alone or with mouth-to-mouth ventilation. *N Engl J Med*. 2000;342:1546.
28. Harrison RR, Maull KI, Keenan RL, et al. Mouth-to-mask ventilation: a superior method of rescue breathing. *Ann Emerg Med*. 1982;11:74.
29. Hess D, Baran C. Ventilatory volumes using mouth-to-mouth, mouth-to-mask, and bag-valve mask techniques. *Am J Emerg Med*. 1985;3:292.
30. International Liaison Committee on Resuscitation and the American Heart Association. 2005 International consensus on cardiopulmonary resuscitation (CPR) and emergency cardiovascular care (ECC) science with treatment recommendations. *Circulation*. 2005;112:III-1-III-136.

31. Babbs CF. New versus old theories of blood flow during CPR. *Crit Care Med.* 1980;8:191.
32. Ralston SH, Voorhees WD, Babbs CF. Intrapulmonary epinephrine during prolonged CPR. Improved regional blood flow and resuscitation in dogs. *Ann Emerg Med.* 1984;13:79.
33. Crile G, Dolley DH. Experimental research into resuscitation of dogs killed by anesthetics and asphyxia. *J Exp Med.* 1906;8:713.
34. Pearson JW, Redding JS. Influence of peripheral vascular tone on cardiac resuscitation. *Anesth Analg.* 1965;44:746.
35. Redding JS. Abdominal compression in cardiopulmonary resuscitation. *Anesth Analg.* 1971;50:668.
36. Yakaitis RW, Otto CW, Blitt CD. Relative importance of α and β adrenergic receptors during resuscitation. *Crit Care Med.* 1979;7:293.
37. Otto CW, Yakaitis RW, Blitt CD. Mechanism of action of epinephrine in resuscitation from asphyxial arrest. *Crit Care Med.* 1981;9:321.
38. Michael JR, Guerci AD, Koehler RC, et al. Mechanisms by which epinephrine augments cerebral and myocardial perfusion during cardiopulmonary resuscitation in dogs. *Circulation.* 1984;69:822.
39. Otto CW, Yakaitis RW. The role of epinephrine in CPR: a reappraisal. *Ann Emerg Med.* 1984;13:840.
40. Sanders AB, Ewy GA, Taft TV. Prognostic and therapeutic importance of the aortic diastolic pressure in resuscitation from cardiac arrest. *Crit Care Med.* 1984;12:871.
41. Niemann JT, Criley JM, Rosborough JP, et al. Predictive indices of successful cardiac resuscitation after prolonged arrest and experimental cardiopulmonary resuscitation. *Ann Emerg Med.* 1985;14:521.
42. Paradis NA, Martin GB, Rivers EP, et al. Coronary perfusion pressure and the return of spontaneous circulation in human cardiopulmonary resuscitation. *JAMA.* 1990;263:1106.
43. Weil MH, Rackow EC, Trevino R, et al. Difference in acid-base state between venous and arterial blood during cardiopulmonary resuscitation. *N Engl J Med.* 1986;315:153.
44. Babbs CF, Tacker WA. Cardiopulmonary resuscitation with interposed abdominal compression. *Circulation.* 1986;74 (Suppl IV):IV-37.
45. Mateer JR, Stueven HA, Thompson BM, et al. Pre-hospital IAC-CPR versus standard CPR. Paramedic resuscitation of cardiac arrests. *Am J Emerg Med.* 1985;3:143.
46. Sack JB, Kesselbrenner MB, Bregman D. Survival from in-hospital cardiac arrest with interposed abdominal counterpulsation during cardiopulmonary resuscitation. *JAMA.* 1992;267:379.
47. Sack JB, Kesselbrenner MB, Jarrad A. Interposed abdominal compression-cardiopulmonary resuscitation and resuscitation outcome during asystole and electromechanical dissociation. *Circulation.* 1992;86:1692.
48. Lurie KG, Lindo C, Chin J. CPR. The P stands for plumber's helper [letter]. *JAMA.* 1990;264:1661.
49. Linder KH, Pfenniger EG, Lurie KG, et al. Effects of active compression-decompression resuscitation on myocardial and cerebral blood flow in pigs. *Circulation.* 1993;88:1254.
50. Shultz JJ, Coffeen P, Sweeney M, et al. Evaluation of standard and active compression-decompression CPR in an acute human model of ventricular fibrillation. *Circulation.* 1994;89:684.
51. Lafuente-Lafuente C, Melero-Bascones M. Active compression-decompression for cardiopulmonary resuscitation. *Cochrane Database Syst Rev.* 2004;CD002751.
52. Steen S, Liao Q, Pierre L, et al. Evaluation of LUCAS, a new device for automatic mechanical compression and active decompression resuscitation. *Resuscitation.* 2002;55:285.
53. Lurie KG, Mulligan KA, McKnite S, et al. Optimizing standard cardiopulmonary resuscitation with an inspiratory impedance threshold valve. *Chest.* 1998;113:1084.
54. Cabrini L, Beccaria P, Landoni G, et al. Impact of impedance threshold devices on cardiopulmonary resuscitation: a systematic review and meta-analysis of randomized controlled studies. *Crit Care Med.* 2008;36:1625.
55. Criley JM, Niemann JT, Rosborough JP, et al. Modifications of cardiopulmonary resuscitation based on the cough. *Circulation.* 1986;74(Suppl IV):IV-42.
56. Halperin HR, Guerci AD, Chandra N, et al. Vest inflation without simultaneous ventilation during cardiac arrest in dogs: improved survival from prolonged cardiopulmonary resuscitation. *Circulation.* 1986;74:1407.
57. Timerman S, Cardoso LF, Ramires JA, Halperin H. Improved hemodynamic performance with a novel chest compression device during treatment of in-hospital cardiac arrest. *Resuscitation.* 2004;61:273.
58. Hallstrom A, Rea TD, Sayre MR, et al. Manual chest compression vs use of an automated chest compression device during resuscitation following out-of-hospital cardiac arrest: a randomized trial. *JAMA.* 2006;295:2620.
59. Bircher N, Safar P, Stewart R. A comparison of standard, MAST-augmented, and open-chest CPR in dogs: a preliminary investigation. *Crit Care Med.* 1980;8:147.
60. DeBehnke DJ, Angelos MG, Leasure JE. Comparison of standard external CPR, open-chest CPR, and cardiopulmonary bypass in a canine myocardial infarct model. *Ann Emerg Med.* 1991;20:754.
61. Sanders AB, Kern KB, Ewy GA, et al. Improved resuscitation from cardiac arrest with open-chest massage. *Ann Emerg Med.* 1984;13:672.
62. Sanders AB, Kern KB, Atlas M, et al. Importance of the duration of inadequate coronary perfusion pressure on resuscitation from cardiac arrest. *J Am Coll Cardiol.* 1985;6:113.
63. Kern KB, Sanders AB, Ewy GA. Open-chest cardiac massage after closed-chest compression in a canine model: when to intervene. *Resuscitation.* 1987;15:51.
64. Kern KB, Sanders AB, Janas W, et al. Limitations of open-chest cardiac massage after prolonged, untreated cardiac arrest in dogs. *Ann Emerg Med.* 1991;20:761.
65. Kern KB, Sanders AB, Badylak SF, et al. Long term survival with open-chest cardiac massage after ineffective closed-chest compression in a canine preparation. *Circulation.* 1987;75:498.
66. Mooney MR, Arom KV, Joyce LD, et al. Emergency cardiopulmonary bypass support in patients with cardiac arrest. *J Thorac Cardiovasc Surg.* 1991;101:450.
67. Angelos M, Safar P, Reich H. A comparison of cardiopulmonary resuscitation with cardiopulmonary bypass after prolonged cardiac arrest in dogs: reperfusion pressures and neurologic recovery. *Resuscitation.* 1991;21:121.
68. Kalenda Z. The capnogram as a guide to the efficacy of cardiac massage. *Resuscitation.* 1978;6:259.
69. Sanders AB, Atlas M, Ewy GA, et al. Expired Pco₂ as an index of coronary perfusion pressure. *Am J Emerg Med.* 1985;3:147.

70. Sanders AB, Atlas M, Ewy GA, et al. Expired PCO₂ as an index of coronary perfusion pressure. *Am J Emerg Med.* 1985;3:147.
71. Sanders AB, Ewy GA, Bragg S, et al. Expired PCO₂ as a prognostic indicator of successful resuscitation from cardiac arrest. *Ann Emerg Med.* 1985;14:948.
72. Trevino RP, Bisera J, Weil MH, et al. End-tidal CO₂ as a guide to successful cardiopulmonary resuscitation: a preliminary report. *Crit Care Med.* 1985;13:910.
73. Sanders AB, Kern KB, Otto CW, et al. End-tidal carbon dioxide monitoring during cardiopulmonary resuscitation: a prognostic indicator for survival. *JAMA.* 1989;262:1347.
74. Kerber RE, Sarnat W. Factors influencing the success of ventricular defibrillation in man. *Circulation.* 1979;60:226.
75. Yakaitis RW, Ewy GA, Otto CW, et al. Influence of time and therapy on ventricular defibrillation in dogs. *Crit Care Med.* 1980;8:157.
76. Weaver WD, Copass MD, Bufe D, et al. Improved neurologic recovery and survival after early defibrillation. *Circulation.* 1984;69:943.
77. Weaver WD, Cobb LA, Hallstrom AP, et al. Factors influencing survival after out-of-hospital cardiac arrest. *J Am Coll Cardiol.* 1986;7:752.
78. Valenzuela TD, Roe DJ, Nichol G, et al. Outcomes of rapid defibrillation by security officers after cardiac arrest in casinos. *N Engl J Med.* 2000;343:1206.
79. Caffrey SL, Willoughby PJ, Pepe PE, et al. Public use of automated external defibrillators. *N Engl J Med.* 2002;347:1242.
80. The Public Access Defibrillation Trial Investigators. Public-access defibrillation and survival after out-of-hospital cardiac arrest. *N Engl J Med.* 2004;351:637.
81. Cobb LA, Fahrenbruch CE, Walsh TR, et al. Influence of CPR prior to defibrillation in out-of-hospital ventricular fibrillation. *JAMA.* 1999;281:1182.
82. Wik L, Hansen TB, Fylling F, et al. Delaying defibrillation to give basic cardiopulmonary resuscitation to patients with out-of-hospital ventricular fibrillation. *JAMA.* 2003;289:1389.
83. Weaver SC, Cobb LA, Dennis D, et al. Amplitude of ventricular fibrillation waveform and outcome after cardiac arrest. *Ann Intern Med.* 1985;102:53.
84. Eftestol T, Wik L, Sunde K, et al. Effects of cardiopulmonary resuscitation on predictors of ventricular fibrillation defibrillation success during out-of-hospital cardiac arrest. *Circulation.* 2004;110:10.
85. Otto CW, Yakaitis RW, Ewy GA. Effects of epinephrine on defibrillation in ischemic ventricular fibrillation. *Am J Emerg Med.* 1985;3:285.
86. Bardy GH, Marchlinski FE, Sharma AD, et al. Multicenter comparison of truncated biphasic shocks and standard damped sine wave monophasic shocks for transthoracic ventricular defibrillation. *Circulation.* 1996;94:2507.
87. Kerber RE, Grayzel J, Hoyt R, et al. Transthoracic resistance in human defibrillation: influence of body weight, chest size, serial shocks, paddle size and paddle contact pressure. *Circulation.* 1981;63:676.
88. Kerber RE, McPherson D, Charbonnier R, et al. Automatic impedance-based energy adjustment for defibrillation: experimental studies. *Circulation.* 1985;71:136.
89. Lerman BB, DeMarco JP, Haines DE. Current-based versus energy-based ventricular defibrillation: a prospective study. *J Am Coll Cardiol.* 1988;12:1259.
90. Kerber RE, Kieso RA, Kienzle MG, et al. Current-based transthoracic defibrillation. *Am J Cardiol.* 1996;78:1113.
91. Dahl CF, Ewy GA, Warner ED, Thomas ED. Myocardial necrosis from direct current countershock. *Circulation.* 1974;50:956.
92. Warner ED, Dahl CF, Ewy GA. Myocardial injury from transthoracic defibrillator countershock. *Arch Pathol.* 1975;99:55.
93. Weaver WD, Cobb LA, Copass MK, et al. Ventricular defibrillation—a comparative trial using 175-J and 320-J shocks. *N Engl J Med.* 1982;307:1101.
94. Geddes LA, Tacker WA, Rosborough JP, et al. Electrical dose for ventricular defibrillation of large and small animals using precordial electrodes. *J Clin Invest.* 1974;53:310.
95. Gutgesell HP, Tacker WA, Geddes LA, et al. Energy dose for defibrillation in children. *Pediatrics.* 1976;58:898.
96. Kerber RE, Sarnat W. Factors influencing the success of ventricular defibrillation in man. *Circulation.* 1979;60:226.
97. Kerber RE, Jensen SR, Gascho JA, et al. Determinants of defibrillation: prospective analysis of 183 patients. *Am J Cardiol.* 1983;52:739.
98. Kudenchuk PJ, Cobb LA, Copass MK, et al. Amiodarone for resuscitation after out of hospital cardiac arrest due to ventricular fibrillation. *N Engl J Med.* 1999;341:871.
99. Dorian P, Cass D, Schwartz B, et al. Amiodarone as compared with lidocaine for shock-resistant ventricular fibrillation. *N Engl J Med.* 2002;346:884.
100. Zoll PM, Zoll RH, Falk RH, et al. External noninvasive temporary cardiac pacing: clinical trials. *Circulation.* 1985;71:937.
101. Ewy GA, Dahl CF, Zimmermann M, et al. Ventricular fibrillation masquerading as ventricular standstill. *Crit Care Med.* 1981;9:841.
102. Liu Y, Rosenthal RE, Haywood Y, et al. Normoxic ventilation after cardiac arrest reduces oxidation of brain lipids and improves neurological outcome. *Stroke.* 1998;29:1679.
103. Zwemer CF, Whitesall SE, D'Alecy LG. Cardiopulmonary-cerebral resuscitation with 100% oxygen exacerbates neurological dysfunction following nine minutes of normothermic cardiac arrest in dogs. *Resuscitation.* 1994;27:159.
104. Lipinski CA, Hicks SD, Callaway CW. Normoxic ventilation during resuscitation and outcome from asphyxial cardiac arrest in rats. *Resuscitation.* 1999;42:221.
105. Redding JS, Pearson JW. Resuscitation from ventricular fibrillation (drug therapy). *JAMA.* 1968;203:255.
106. Schleiens CL, Dean JM, Koehler RC, et al. Effect of epinephrine on cerebral and myocardial perfusion in an infant animal preparation of cardiopulmonary resuscitation. *Circulation.* 1986;73:809.
107. Schleiens CL, Koehler RC, Gervais H, et al. Organ blood flow and somatosensory evoked potentials during and after cardiopulmonary resuscitation with epinephrine or phenylephrine. *Circulation.* 1989;79:1332.
108. Otto CW, Yakaitis RW, Redding JS, Blitt CD. Comparison of dopamine, dobutamine, and epinephrine in CPR. *Crit Care Med.* 1981;9:640.
109. Lindner KH, Prengel AW, Pfenniger EG, et al. Vasopressin improves vital organ blood flow during closed-chest cardiopulmonary resuscitation in pigs. *Circulation.* 1995;91:215.
110. Berg RA, Otto CW, Kern KB, et al. High dose epinephrine results in greater early mortality following resuscitation from prolonged cardiac arrest in pigs: a prospective, randomized study. *Crit Care Med.* 1994;22:282.

111. Brillman JC, Sanders AB, Otto CW, et al. A comparison of epinephrine and phenylephrine for resuscitation and neurologic outcome of cardiac arrest in dogs. *Ann Emerg Med.* 1987;16:11.
112. Olson DW, Thakur R, Stueven HA. Randomized study of epinephrine versus methoxamine in prehospital ventricular fibrillation. *Ann Emerg Med.* 1989;18:250.
113. Silvast T, Saarnivaara L, Kinnunen A, et al. Comparison of adrenaline and phenylephrine in out-of-hospital CPR. A double-blind study. *Acta Anaesthesiol Scand.* 1985;29:610.
114. Wenzel V, Krismer AC, Arntz HR, et al. A comparison of vasopressin and epinephrine for out-of-hospital cardiopulmonary resuscitation. *N Engl J Med.* 2004;350:105.
115. Stiell IG, Wells GA, Field B, et al. Advanced cardiac life support in out-of-hospital cardiac arrest. *N Engl J Med.* 2004;351:647.
116. Olasveengen TM, Sunde K, Brunborg C, et al. Intravenous drug administration during out-of-hospital cardiac arrest: a randomized trial. *JAMA.* 2009;302:2222.
117. Berg RA, Otto CW, Kern KB, et al. A randomized, blinded trial of high-dose epinephrine versus standard-dose epinephrine in a swine model of pediatric asphyxial cardiac arrest. *Crit Care Med.* 1996;24:1695.
118. Vandycke C, Martens P. High dose versus standard dose epinephrine in cardiac arrest—a meta-analysis. *Resuscitation.* 2000;45:161.
119. Lindner KH, Brinkmann A, Pfenninger EG, et al. Effect of vasopressin on hemodynamic variables, organ blood flow, and acid-base status in a pig model of cardiopulmonary resuscitation. *Anesth Analg.* 1993;77:427.
120. Lindner KH, Prengel AW, Pfenninger EG, et al. Vasopressin improves vital organ blood flow during closed-chest cardiopulmonary resuscitation in pigs. *Circulation.* 1995;91:215.
121. Wenzel V, Lindner KH, Krismer AC, et al. Repeated administration of vasopressin, but not epinephrine, maintains coronary perfusion pressure after early and late administration during cardiopulmonary resuscitation in pigs. *Circulation.* 1999;99:1379-1384.
122. Prengel AW, Lindner KH, Keller A, Lurie KG. Cardiovascular function during the postresuscitation phase after cardiac arrest in pigs: a comparison of epinephrine versus vasopressin. *Crit Care Med.* 1996;24:2014.
123. Lindner KH, Dirks B, Strohmenger HU, et al. Randomized comparison of epinephrine and vasopressin in patients with out-of-hospital ventricular fibrillation. *Lancet.* 1997;349:535.
124. Stiell IG, Hebert PC, Wells GA, et al. Vasopressin versus epinephrine for in-hospital cardiac arrest: a randomized controlled trial. *Lancet.* 2001;358:105.
125. Aung K, Htay T. Vasopressin for cardiac arrest: a systematic review and meta-analysis. *Arch Intern Med.* 2005;165:17.
126. Coon GA, Clinton JE, Ruiz E. Use of atropine for bradysystolic prehospital cardiac arrest. *Ann Emerg Med.* 1981;10:462.
127. Stueven HA, Tonsfeldt DJ, Thompson BM, et al. Atropine in asystole: human studies. *Ann Emerg Med.* 1984;13:815.
128. Stueven HA, Thompson BM, Aprahamian C, et al. Calcium chloride: reassessment of use in asystole. *Ann Emerg Med.* 1984;13:820.
129. Stueven HA, Thompson BM, Aprahamian C, et al. The effectiveness of calcium chloride in refractory electromechanical dissociation. *Ann Emerg Med.* 1985;14:626.
130. Federiuk CS, Sanders AB, Kern KB, et al. The effect of bicarbonate on resuscitation from cardiac arrest. *Ann Emerg Med.* 1991;20:1173.
131. Guerci AD, Chandra N, Johnson E, et al. Failure of sodium bicarbonate to improve resuscitation from ventricular fibrillation in dogs. *Circulation.* 1986;74(Suppl IV):IV-75.
132. Bishop RL, Weisfeldt ML. Sodium bicarbonate administration during cardiac arrest: effect on arterial pH, Pco₂, and osmolality. *JAMA.* 1976;235:506.
133. Weil MH, Grundler W, Yamaguchi M, et al. Arterial blood gases fail to reflect acid-base status during cardiopulmonary resuscitation. a preliminary report. *Crit Care Med.* 1985;13:884.
134. Mattar JA, Weil MH, Shubin H, et al. Cardiac arrest in the critically ill. II. Hyperosmolal states following cardiac arrest. *Am J Med.* 1974;56:162.
135. Sanders AB, Otto CW, Kern KB, et al. Acid-base balance in a canine model of cardiac arrest. *Ann Emerg Med.* 1988;17:667.
136. Kette F, Weil MH, von Planta M, et al. Buffer agents do not reverse intramyocardial acidosis during cardiac resuscitation. *Circulation.* 1990;81:1660.
137. Hedges JR, Barsan WB, Doan LA, et al. Central versus peripheral intravenous routes in cardiopulmonary resuscitation. *Am J Emerg Med.* 1984;2:385.
138. Josefson LM. A new method of treatment— intraosseal injections. *Acta Med Scand.* 1934;81:550.
139. Berg RA. Emergency infusion of catecholamines into bone marrow. *Am J Dis Child.* 1984;138:810.
140. Morris RE, Schonfeld N, Haftel AJ. Treatment of hemorrhagic shock with intraosseous administration of crystalloid fluid in the rabbit model. *Ann Emerg Med.* 1987;16:1321.
141. Spivey WH, Lathers CM, Malone DR, et al. Comparison of intraosseous, central, and peripheral routes of sodium bicarbonate administration during CPR in pigs. *Ann Emerg Med.* 1985;14:1135.
142. Andropoulos DB, Soifer SJ, Schreiber MD. Plasma epinephrine concentrations after intraosseous and central venous injection during cardiopulmonary resuscitation in the lamb. *J Pediatr.* 1990;166:312.
143. McDonald JL. Serum lidocaine levels during cardiopulmonary resuscitation after intravenous and endotracheal administration. *Crit Care Med.* 1985;13(11):914.
144. Ralston SH, Tacker WA, Showen L, et al. Endotracheal versus intravenous epinephrine during electromechanical dissociation with CPR in dogs. *Ann Emerg Med.* 1985;14:1044.
145. Hornchen U, Schuttler J, Stoeckel H, et al. Endobronchial instillation of epinephrine during cardiopulmonary resuscitation. *Crit Care Med.* 1987;15:1037.
146. Abramson NS, Safar P, Detre KM, et al. Randomized clinical study of cardiopulmonary-cerebral resuscitation: thiopental loading in comatose cardiac arrest survivors. *N Engl J Med.* 1986;314:397.
147. Brain Resuscitation Clinical Trial II Study Group. A randomized clinical study of a calcium-entry blocker (lidoflazine) in the treatment of comatose survivors of cardiac arrest. *N Engl J Med.* 1991;324:1225.
148. Holzer M, The Hypothermia After Cardiac Arrest Study Group. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med.* 2002;346:549.
149. Bernard SA, Gray TW, Buist MD, et al. Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med.* 2002;346:557.

150. Booth CM, Boone RH, Tomlinson G, et al. Is this patient dead, vegetative, or severely neurologically impaired? Assessing outcome for comatose survivors of cardiac arrest. *JAMA*. 2004;291:870.
151. Longstreth WT, Inui TS, Cobb LA, Copass MK. Neurologic recovery after out-of-hospital cardiac arrest, *Ann Intern Med*. 1983;98:588.
152. Zandbergen EG, Hijdra A, Koelman JH, et al. Prediction of poor outcome within the first 3 days of postanoxic coma. *Neurology*. 2006;66:62.
153. Edgren E, Hedstrand U, Nordin M, et al. Prediction of outcome after cardiac arrest. *Crit Care Med*. 1987;15:820.
154. Young GB, Doig G, Ragazzoni A. Anoxic-ischemic encephalopathy: clinical and electrophysiological associations with outcome. *Neurocrit Care*. 2005;2:159.