

Chapter 84

REFERENCES

1. Moller JT, Cluitmans P, Rasmussen LS, et al. Long-term postoperative cognitive dysfunction in the elderly ISPOCD1 study. ISPOCD investigators. International Study of Post-Operative Cognitive Dysfunction. *Lancet*. 1998;351(9106):857-861.
2. Newman MF, Grocott HP, Mathew JP, et al. Report of the sub-study assessing the impact of neurocognitive function on quality of life five years after cardiac surgery. Neurologic Outcome Research Group and Cardiothoracic Anesthesia Research Endeavors (CARE) Investigators of the Duke Heart Center. *Stroke*. 2001;32(12):2874-2881.
3. Rasmussen LS. Defining postoperative cognitive dysfunction. *Eur J Anaesthesiol*. 1998;15(6):761-764.
4. Silverstein JH, Timberger M, Reich DL, Uysal S. Central nervous system dysfunction after noncardiac surgery and anesthesia in the elderly. *Anesthesiology*. 2007;106(3):622-628.
5. Steinmetz J, Christensen KB, Lund T, Lohse N, Rasmussen LS. Long-term consequences of postoperative cognitive dysfunction. *Anesthesiology*. 2009;110(3):548-555.
6. Hudetz JA, Patterson KM, Byrne AJ, Pagel PS, Warltier DC. Postoperative delirium is associated with postoperative cognitive dysfunction at one week after cardiac surgery with cardiopulmonary bypass. *Psychol Rep*. 2009;105(3 Pt 1):921-932.
7. Rudolph JL, Marcantonio ER, Culley DJ, et al. Delirium is associated with early postoperative cognitive dysfunction. *Anaesthesia*. 2008;63(9):941-947.
8. Bekker AY, Weeks EJ. Cognitive function after anaesthesia in the elderly. *Best Pract Res Clin Anaesthesiol*. 2003;17(2):259-272.
9. Koster S, Hensens AG, van der Palen J. The long-term cognitive and functional outcomes of postoperative delirium after cardiac surgery. *Ann Thorac Surg*. 2009;87(5):1469-1474.
10. Bitsch M, Foss N, Kristensen B, Kehlet H. Pathogenesis of and management strategies for postoperative delirium after hip fracture: a review. *Acta Orthop Scand*. 2004;75(4):378-389.
11. Leung JM, Sands LP, Mullen EA, Wang Y, Vaurio L. Are pre-operative depressive symptoms associated with postoperative delirium in geriatric surgical patients? *J Gerontol A Biol Sci Med Sci*. 2005;60(12):1563-1568.
12. Litaker D, Locala J, Franco K, Bronson DL, Tannous Z. Preoperative risk factors for postoperative delirium. *Gen Hosp Psychiatry*. 2001;23(2):84-89.
13. Cole MG. Delirium in elderly patients. *Am J Geriatr Psychiatry*. 2004;12(1):7-21.
14. Fong HK, Sands LP, Leung JM. The role of postoperative analgesia in delirium and cognitive decline in elderly patients: a systematic review. *Anesth Analg*. 2006;102(4):1255-1266.
15. Marcantonio ER, Goldman L, Orav EJ, Cook EF, Lee TH. The association of intraoperative factors with the development of postoperative delirium. *Am J Med*. 1998;105(5):380-384.
16. Smith PJ, Attix DK, Weldon BC, Greene NH, Monk TG. Executive function and depression as independent risk factors for postoperative delirium. *Anesthesiology*. 2009;110(4):781-787.
17. Franco K, Litaker D, Locala J, Bronson D. The cost of delirium in the surgical patient. *Psychosomatics*. 2001;42(1):68-73.
18. Monk TG, Weldon BC, Garvan CW, et al. Predictors of cognitive dysfunction after major noncardiac surgery. *Anesthesiology*. 2008;108(1):18-30.
19. Hanning CD, Blokland A, Johnson M, Perry EK. Effects of repeated anaesthesia on central cholinergic function in the rat cerebral cortex. *Eur J Anaesthesiol*. 2003;20(2):93-97.
20. Phillips-Bute B, Mathew JP, Blumenthal JA, et al. Association of neurocognitive function and quality of life 1 year after coronary artery bypass graft (CABG) surgery. *Psychosom Med*. 2006;68(3):369-375.
21. Holm K, Foreman M. Analysis of measures of functional and cognitive ability for aging adults with cardiac and vascular disease. *J Cardiovasc Nurs*. 2006;21(5 Suppl 1):S40-S45.
22. Ryan CM. Diabetes, aging, and cognitive decline. *Neurobiol Aging*. 2005;26(Suppl 1):21-25.
23. Ritchie K. Mild cognitive impairment: an epidemiological perspective. *Dialogues Clin Neurosc*. 2004;6(4):401-408.
24. Di Bonito P, Di Fraia L, Di Gennaro L, et al. Impact of impaired fasting glucose and other metabolic factors on cognitive function in elderly people. *Nutr Metab Cardiovasc Dis*. 2007;17(3):203-208.
25. Panza F, Solfrizzi V, Colacicco AM, et al. Cerebrovascular disease in the elderly: lipoprotein metabolism and cognitive decline. *Aging Clin Exp Res*. 2006;18(2):144-148.
26. Strittmatter WJ, Roses AD. Apolipoprotein E and Alzheimer disease. *Proc Natl Acad Sci USA*. 1995;92(11):4725-4727.
27. Strittmatter WJ, Saunders AM, Goedert M, et al. Isoform-specific interactions of apolipoprotein E with microtubule-associated protein tau: implications for Alzheimer disease. *Proc Natl Acad Sci USA*. 1994;91(23):11183-11186.
28. Bondi MW, Salmon DP, Monsch AU, et al. Episodic memory changes are associated with the APOE-epsilon 4 allele in non-demented older adults. *Neurology*. 1995;45(12):2203-2206.

29. Caselli RJ, Osborne D, Reiman EM, et al. Preclinical cognitive decline in late middle-aged asymptomatic apolipoprotein E-e4/4 homozygotes: a replication study. *J Neurol Sci*. 2001;189(1-2):93-98.
30. den Heijer T, Oudkerk M, Launer LJ, van Duijn CM, Hofman A, Breteler MM. Hippocampal, amygdalar, and global brain atrophy in different apolipoprotein E genotypes. *Neurology*. 2002;59(5):746-748.
31. Flory JD, Manuck SB, Ferrell RE, Ryan CM, Muldoon MF. Memory performance and the apolipoprotein E polymorphism in a community sample of middle-aged adults. *Am J Med Genet*. 2000;96(6):707-711.
32. Greenwood PM, Lambert C, Sunderland T, Parasuraman R. Effects of apolipoprotein E genotype on spatial attention, working memory, and their interaction in healthy, middle-aged adults: results from the National Institute of Mental Health's BIOCARD study. *Neuropsychology*. 2005;19(2):199-211.
33. Schuff N, Amend DL, Knowlton R, Norman D, Fein G, Weiner MW. Age-related metabolic changes and volume loss in the hippocampus by magnetic resonance spectroscopy and imaging. *Neurobiol Aging*. 1999;20(3):279-285.
34. Breteler MM, van Swieten JC, Bots ML, et al. Cerebral white matter lesions, vascular risk factors, and cognitive function in a population-based study: the Rotterdam Study. *Neurology*. 1994;44(7):1246-1252.
35. Haan MN, Shemanski L, Jagust WJ, Manolio TA, Kuller L. The role of APOE epsilon4 in modulating effects of other risk factors for cognitive decline in elderly persons. *JAMA*. 1999;282(1):40-46.
36. Slooter AJ, van Duijn CM, Bots ML, et al. Apolipoprotein E genotype, atherosclerosis, and cognitive decline: the Rotterdam Study. *J Neural Transm Suppl*. 1998;53:17-29.
37. Tardiff BE, Newman MF, Saunders AM, et al. Preliminary report of a genetic basis for cognitive decline after cardiac operations. The Neurologic Outcome Research Group of the Duke Heart Center. *Ann Thorac Surg*. 1997;64(3):715-720.
38. McDonagh DL, Mathew JP, White WD, et al. Cognitive function after major noncardiac surgery, apolipoprotein E4 genotype, and biomarkers of brain injury. *Anesthesiology*. 2010;112(4):852-859.
39. Mathew JP, Podgoreanu MV, Grocott HP, et al. Genetic variants in P-selectin and C-reactive protein influence susceptibility to cognitive decline after cardiac surgery. *J Am Coll Cardiol*. 2007;49(19):1934-1942.
40. Milgram NW, Siwak-Tapp CT, Araujo J, Head E. Neuroprotective effects of cognitive enrichment. *Ageing Res Rev*. 2006;5(3):354-369.
41. Newman MF, Kirchner JL, Phillips-Bute B, et al. Longitudinal assessment of neurocognitive function after coronary-artery bypass surgery. Neurological Outcome Research Group and CARE Investigators. *N Engl J Med*. 2001;344(6):395-402.
42. Newman MF, Wolman R, Kanchuger M, et al. Multicenter preoperative stroke risk index for patients undergoing coronary artery bypass graft surgery. Multicenter Study of Perioperative Ischemia (McSPI) Research Group. *Circulation*. 1996;94(Suppl II):II-74-II-80.
43. Williams-Russo P, Sharrock NE, Mattis S, Szatrowski TP, Charlson ME. Cognitive effects after epidural vs general anesthesia in older adults. A randomized trial. *JAMA*. 1995;274(1): 44-50.
44. Mahanna EP, Blumenthal JA, White WD, et al. Defining neuropsychological dysfunction after coronary artery bypass grafting. *Ann Thorac Surg*. 1996;61(5):1342-1347.
45. Rasmussen LS, Larsen K, Houx P, Skovgaard LT, Hanning CD, Moller JT. The assessment of postoperative cognitive function. *Acta Anaesthesiol Scand*. 2001;45(3):275-289.
46. Murkin JM. The role of CPB management in neurobehavioral outcomes after cardiac surgery. *Ann Thorac Surg*. 1995;59(5):1308-1311.
47. Murkin JM, Stump DA, Blumenthal JA, McKhann G. Defining dysfunction: group means versus incidence analysis—a statement of consensus. *Ann Thorac Surg*. 1997;64(3):904-905.
48. Newman S, Arrowsmith J, Grocott H, Murkin J, Stump D, Wingerhoets G. The Outcomes 2006 Statement of Consensus on Cognitive Difficulties Following Coronary Artery Bypass Surgery. Available: <http://www.outcomeskeywest.com/portals/16/Consensus2006.pdf>. Accessed on December 13, 2011.
49. Ahlgren E, Lundqvist A, Nordlund A, Aren C, Rutberg H. Neurocognitive impairment and driving performance after coronary artery bypass surgery. *Eur J Cardiothorac Surg*. 2003;23(3):334-340.
50. Whelihan WM, DiCarlo MA, Paul RH. The relationship of neuropsychological functioning to driving competence in older persons with early cognitive decline. *Arch Clin Neuropsych*. 2005;20(2):217-228.
51. Cosgrove DM, Loop FD, Lytle BW, et al. Primary myocardial revascularization. Trends in surgical mortality. *J Thorac Cardiovasc Surg*. 1984;88(5 Pt 1):673-684.
52. Califf R, Harrell F, Lee K, et al. Changing efficacy of coronary revascularization: implications for patient selection. *Circulation*. 1988;78(Suppl 1):I-185-I-191.
53. Gersh B, Kornmal R, Schaff H, et al. Comparison of coronary artery bypass surgery and medical therapy in patients 65 years of age or older. *N Engl J Med*. 1985;313(4):217-224.
54. Hammermeister KE, Burchfiel C, Johnson R, Grover FL. Identification of patients at greatest risk for developing major complications at cardiac surgery. *Circulation*. 1990;82(5 Suppl):IV-380-IV-389.
55. Jones EL, Weintraub WS, Carver JM, Guyton RA, Cohen CL. Coronary bypass surgery: is the operation different today? *J Thorac Cardiovasc Surg*. 1991;101(1):108-115.
56. Mohan R, Amsel BJ, Walter PL. Coronary artery bypass grafting in the elderly: a review of studies on patients older than 64, 69, or 74 years. *Cardiology*. 1992;80(3-4):215-225.
57. Wolman RL, Nussmeier NA, Aggarwal A, et al. Cerebral injury after cardiac surgery: identification of a group at extraordinary risk. *Stroke*. 1999;30(3):514-522.
58. Cleveland JC, Shroyer LW, Chen A, Peterson E, Grover FL. Off-pump coronary artery bypass grafting significantly decreases risk-adjusted mortality and morbidity. *Ann Thorac Surg*. 2001;72(4):1282-1289.
59. van Dijk D, Spoor M, Hijman R, et al. Cognitive and cardiac outcomes 5 years after off-pump vs on-pump coronary artery bypass graft surgery. *JAMA*. 2007;297(7):701-708.
60. Jensen BØ, Rasmussen LS, Steinbruchel DA. Cognitive outcomes in elderly high-risk patients 1 year after off-pump versus on-pump coronary artery bypass grafting. A randomized trial. *Eur J Cardiothorac Surg*. 2008;34(5):1016-1021.
61. Liu YH, Wang DX, Li LH, et al. The effects of cardiopulmonary bypass on the number of cerebral microemboli and the incidence of cognitive dysfunction after coronary artery bypass graft surgery. *Anesth Analg*. 2009;109(4):1013-1022.
62. Bonke B. Postoperative cognitive dysfunction in the elderly. *Lancet*. 1998;351(9119):1889.

63. Kanchuger MS, Sweeney MN, Grossi E, Marschall KE. Aortic atheromatous disease, atherectomy and outcome in patients undergoing cardiac surgery. *Anesth Analg*. 1996;82:SCA31.
64. Roach GW, Kanchuger M, Mora Mangano C, et al. Adverse cerebral outcomes after coronary bypass surgery. *N Engl J Med*. 1996;335(25):1857-1863.
65. Wareing TH, Davila-Roman VG, Barzilai B, Murphy SF, Kouchoukos NT. Management of the severely atherosclerotic ascending aorta during cardiac operations: a strategy for detection and treatment. *J Thorac Cardiovasc Surg*. 1992;103(3):453-462.
66. Moody DM, Brown WR, Challa VR, Stump DA, Reboussin DM, Legault C. Brain microemboli associated with cardiopulmonary bypass: a histologic and magnetic resonance imaging study. *Ann Thorac Surg*. 1995;59(5):1304-1307.
67. Floyd TF, Shah PN, Price CC, et al. Clinically silent cerebral ischemic events after cardiac surgery: their incidence, regional vascular occurrence, and procedural dependence. *Ann Thorac Surg*. 2006;81(6):2160-2166.
68. Kadoi Y, Goto F. Factors associated with postoperative cognitive dysfunction in patients undergoing cardiac surgery. *Surg Today*. 2006;36(12):1053-1057.
69. Evered LA, Silbert BS, Scott DA. Postoperative cognitive dysfunction and aortic atheroma. *Ann Thorac Surg*. 2010;89(4):1091-1097.
70. Gilman S. Cerebral disorders after open heart operations. *N Eng J Med*. 1965;272:489-498.
71. Caplan LR, Hennerici M. Impaired clearance of emboli (washout) is an important link between hypoperfusion, embolism, and ischemic stroke. *Arch Neurol*. 1998;55(11):1475-1482.
72. Gold JP, Charlson ME, Williams-Russo P, et al. Improvement of outcomes after coronary artery bypass. A randomized trial comparing intraoperative high versus low mean arterial pressure. *J Thorac Cardiovasc Surg*. 1995;110(5):1302-1311.
73. Hartman GS, Yao FS, Bruefach M 3rd, et al. Severity of aortic atheromatous disease diagnosed by transesophageal echocardiography predicts stroke and other outcomes associated with coronary artery surgery: a prospective study. *Anesth Analg*. 1996;83(4):701-708.
74. Newman MF, Croughwell ND, Blumenthal JA, et al. Effect of aging on cerebral autoregulation during cardiopulmonary bypass. Association with postoperative cognitive dysfunction. *Circulation*. 1994;90(5 Pt 2):II243-II249.
75. Tuman KJ, McCarthy RJ, Najafi H, Ivankovich AD. Differential effects of advanced age on neurologic and cardiac risks of coronary artery operations. *J Thorac Cardiovasc Surg*. 1992;104(6):1510-1517.
76. Newman MF, Kramer DC, Croughwell ND, et al. Differential age effects of mean arterial pressure and rewarming on cognitive dysfunction after cardiac surgery. *Anesth Analg*. 1995;81(2):236-242.
77. de Lange F, Jones WL, Mackensen GB, Grocott HP. The effect of limited rewarming and postoperative hypothermia on cognitive function in a rat cardiopulmonary bypass model. *Anesth Analg*. 2008;106(3):739-745.
78. Grigore AM, Grocott HP, Mathew JP, et al. The rewarming rate and increased peak temperature alter neurocognitive outcome after cardiac surgery. *Anesth Analg*. 2002;94(1):4-10, table of contents.
79. Grocott HP, Mackensen GB, Grigore AM, et al. Postoperative hyperthermia is associated with cognitive dysfunction after coronary artery bypass graft surgery. *Stroke*. 2002;33(2):537-541.
80. Grocott HP, Newman MF, El-Moalem H, Bainbridge D, Butler A, Laskowitz DT. Apolipoprotein E genotype differentially influences the proinflammatory and anti-inflammatory response to cardiopulmonary bypass. *J Thorac Cardiovasc Surg*. 2001;122(12):622-623.
81. Westaby S, Saatvedt K, White S, Katsumata T, van Oeveren W, Halligan PW. Is there a relationship between cognitive dysfunction and systemic inflammatory response after cardiopulmonary bypass? *Ann Thorac Surg*. 2001;71(2):667-672.
82. Mathew JP, Shernan SK, White WD, et al. Preliminary report of the effects of complement suppression with pexelizumab on neurocognitive decline after coronary artery bypass graft surgery. *Stroke*. 2004;35(10):2335-2339.
83. Ebmeier K, Rose E, Steele D. Cognitive impairment and fMRI in major depression. *Neurotox Res*. 2006;10(2):87-92.
84. Steed L, Kong R, Stygall J, et al. The role of apolipoprotein E in cognitive decline after cardiac operation. *Ann Thorac Surg*. 2001;71(3):823-826.
85. Mathew JP, Rinder CS, Howe JG, et al. Platelet PIA2 polymorphism enhances the risk of neurocognitive decline after cardiopulmonary bypass. Multicenter Study of Perioperative Ischemia (McSPI) Research Group. *Ann Thorac Surg*. 2001;71(2):663-666.
86. Ridker PM, Hennekens CH, Schmitz C, Stampfer MJ, Lindpaintner K. PLA1/A2 polymorphism of platelet glycoprotein IIIa and risks of myocardial infarction, stroke, and venous thrombosis. *Lancet*. 1997;349(9049):385-388.
87. Weiss EJ, Bray PF, Tayback M, et al. A polymorphism of a platelet glycoprotein receptor as an inherited risk factor for coronary thrombosis. *N Engl J Med*. 1996;334(17):1090-1094.
88. Monk TG, Saini V, Weldon BC, Sigl JC. Anesthetic management and one-year mortality after noncardiac surgery. *Anesth Analg*. 2005;100(1):4-10.
89. Schneck HJ, Ruprecht J. Central anticholinergic syndrome (CAS) in anesthesia and intensive care. *Acta Anaesthesiol Belg*. 1989;40(3):219-228.
90. Fodale V, Santamaria LB. The inhibition of central nicotinic nACh receptors is the possible cause of prolonged cognitive impairment after anesthesia. *Anesth Analg*. 2003;97(4):1207.
91. Fodale V, Santamaria LB. Drugs of anesthesia, central nicotinic receptors and post-operative cognitive dysfunction. *Acta Anaesthesiol Scand*. 2003;47(9):1180.
92. Praticò C, Quattrone D, Lucanto T, et al. Drugs of anesthesia acting on central cholinergic system may cause post-operative cognitive dysfunction and delirium. *Med Hypotheses*. 2005;65(5):972-982.
93. Abe K, Taguchi K, Kato M, et al. Characterization of muscarinic receptor subtypes in the rostral ventrolateral medulla and effects on morphine-induced antinociception in rats. *Eur J Pharmacol*. 2003;465(3):237-249.
94. Murasaki O, Kaibara M, Nagase Y, et al. Site of action of the general anesthetic propofol in muscarinic M1 receptor-mediated signal transduction. *J Pharmacol Exp Ther*. 2003;307(3):995-1000.
95. Yamanoue T, Brum JM, Estafanous FG, Khairallah PA, Ferrario CM. Fentanyl attenuates porcine coronary arterial contraction through M3-muscarinic antagonism. *Anesth Analg*. 1993;76(2):382-390.
96. Nordberg A, Wahlström G. Cholinergic mechanisms in physical dependence on barbiturates, ethanol and benzodiazepines. *J Neural Transm Gen Sect*. 1992;88(3):199-221.

97. Eckenhoff RG, Johnansson JS, Wei H, et al. Inhaled anesthetic enhancement of amyloid-beta oligomerization and cytotoxicity. *Anesthesiology*. 2004;101(3):703-709.
98. Xie Z, Dong Y, Maeda U, et al. The common inhalation anesthetic isoflurane induces apoptosis and increases amyloid beta protein levels. *Anesthesiology*. 2006;104(5):988-994.
99. Xie Z, Dong Y, Maeda U, et al. Isoflurane-induced apoptosis: a potential pathogenic link between delirium and dementia. *J Gerontol A Biol Sci Med Sci*. 2006;61(12):1300-1306.
100. Xie Z, Dong Y, Maeda U, et al. The inhalation anesthetic isoflurane induces a vicious cycle of apoptosis and amyloid beta-protein accumulation. *J Neurosci*. 2007;27(6):1247-1254.
101. Mandal PK, Williams JP, Mandal R. Molecular understanding of Abeta peptide interaction with isoflurane, propofol, and thiopental: NMR spectroscopic study. *Biochemistry*. 2007;46(3):762-771.
102. Marcantonio ER, Rudolph JL, Culley D, Crosby G, Alsop D, Inouye SK. Serum biomarkers for delirium. *J Gerontol A Biol Sci Med Sci*. 2006;61(12):1281-1286.
103. Abildstrom H, Christiansen M, Siersma VD, Rasmussen LS. Apolipoprotein E genotype and cognitive dysfunction after noncardiac surgery. *Anesthesiology*. 2004;101(4):855-861.
104. Rasmussen, LS, Johnson, T, Kuipers, HM, et al. Does anaesthesia cause postoperative cognitive dysfunction? A randomised study of regional versus general anaesthesia in 438 elderly patients. *Acta Anaesthesiologica Scandinavica*, 47:260-266.
105. Everend, L, Scott, D, Silert, , Maruff, P. Postoperative cognitive dysfunction is independent of type of surgery and anesthetic. *Anesth Analg*. 2011;112(5):1179-1185.
106. Goto T, Yoshitake A, Baba T, Shibata Y, Sakata R, Uozumi H. Cerebral ischemic disorders and cerebral oxygen balance during cardiopulmonary bypass surgery: preoperative evaluation using magnetic resonance imaging and angiography. *Anesth Analg*. 1997;84(1):5-11.